Supplementary Table 1. Pre-transplant patient demographics and tumour characteristics in the overall patient cohort (n = 435), patients with cirrhosis and hepatocellular carcinoma undergoing LDLT

Patient demographics $(n = 435)$		Values
Age (years)	Median (range)	55 (20-73)
Gender	Male Female	348 (80%) 87 (20%)
Primary disease	Hepatitis C Hepatitis B Ethanol NAFLD/NASH Cryptogenic Others	196 (45%) 113 (26%) 44 (10%) 39 (9%) 30 (7%) 13 (3%)
Non-tumour MELD score Neutrophil lymphocyte ratio (NLR) Platelet lymphocyte ration (PLR)	Median (range) Median (IQR) Median (IQR)	10 (6-38) 2.4 (1.7-3.7) 435 (292-642)
Tumour characteristics		
Pre-LT AFP ng/mL	Median (IQR), range	24 (7-169), 1-17,500
Number of tumours	Single 2-3 > 3	221 (55%) 117 (29%) 65 (16%)
Maximum tumour diameter	Mean ±SD	4.15 ± 2.95
Milan Criteria UCSF Criteria	In/Out In/Out	213 (49%)/222 (51%) 278 (57%)/187 (43%)
FDG-18 PET (available in 335 patients)	PET avid Non PET avid	168 (50.1%) 167 (49.9%)