

Review

Open Access



Trends in the evolution to robot-assisted minimally invasive thoracoscopic esophagectomy

Jiajia Liu, Satoru Motoyama, Yusuke Sato, Akiyuki Wakita, Yuta Kawakita, Yushi Nagaki, Hiromu Fujita, Kazuhiro Imai, Yoshihiro Minamiya

Esophageal Surgery, Akita University Hospital, Akita 010-0043, Japan.

Correspondence to: Dr. Satoru Motoyama, Esophageal Surgery, Akita University Hospital, 1-1-1 Hondo, Akita 010-8543, Japan.
E-mail: motoyama@doc.med.akita-u.ac.jp

How to cite this article: Liu J, Motoyama S, Sato Y, Wakita A, Kawakita Y, Nagaki Y, Fujita H, Imai K, Minamiya Y. Trends in the evolution to robot-assisted minimally invasive thoracoscopic esophagectomy. *Mini-invasive Surg* 2020;4:44.
<http://dx.doi.org/10.20517/2574-1225.2020.20>

Received: 13 Feb 2020 **First Decision:** 1 Apr 2020 **Revised:** 30 Apr 2020 **Accepted:** 24 Jun 2020 **Published:** 19 Jul 2020

Academic Editor: Itasu Ninomiya **Copy Editor:** Cai-Hong Wang **Production Editor:** Jing Yu

Abstract

Much effort has been made to improve outcomes and/or minimize the invasiveness of esophagectomy for thoracic esophageal cancer. This has led to the evolution from open esophagectomy to thoracoscopic minimally invasive esophagectomy (MIE), and from MIE to robot-assisted minimally invasive esophagectomy (RAMIE). RAMIE is being applied clinically to overcome the limitations of MIE. In this article, we review the trends in the evolution from thoracoscopic MIE to RAMIE. It has now been demonstrated that RAMIE is both safe and feasible, and may decrease morbidity and mortality rates associated with esophagectomy and improve oncological outcomes. On the other hand, there are still many problems that need to be solved.

Keywords: Esophagectomy, esophageal cancer, robot-assisted esophagectomy, thoracoscopic esophagectomy

INTRODUCTION

Esophageal cancer is the 6th highest cause of cancer mortality worldwide due, in large part, to its high potential for metastasis^[1]. The most reliable curative treatment is surgery entailing radical resection of the esophagus with extended lymphadenectomy in the mediastinum, abdomen, and neck. However, esophagectomy is associated with high postoperative morbidity (about 40%) and mortality (about 3.4%)^[2,3]. To improve outcomes, patients are often treated with multimodal treatments such as neoadjuvant



© The Author(s) 2020. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, sharing, adaptation, distribution and reproduction in any medium or format, for any purpose, even commercially, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.



Table 1. RAMIE history

Year	
1960s	Start of development of a remote operation system
1998	DVSS enters clinical trials, first commercial sale
2000	DVSS obtains Food and Drug Administration clearance
2001	Performance of the first transatlantic surgery (robotic cholecystectomy)
2003	The first transhiatal RAMIE
2004	The first transthoracic RAMIE

DVSS: Da Vinci Surgical System; RAMIE: robot-assisted minimally invasive esophagectomy

chemotherapy or chemoradiotherapy, and there is much surgical effort towards improving operative techniques^[4,5]. This has led to the evolution from open esophagectomy (OE) to thoracoscopic minimally invasive esophagectomy (MIE)^[6], and from MIE to robot-assisted minimally invasive esophagectomy (RAMIE)^[7]. Despite the many advantages of MIE, there are several associated limitations. RAMIE, which has advantages in terms of an enhanced three-dimensional magnified view, tremorless action, and articulated instruments, is being applied clinically to overcome the limitations of MIE^[8]. In this article, we review the trends in the evolution from thoracoscopic esophagectomy to MIE and RAMIE.

HISTORY OF RAMIE

In the 1960s, the US Army and NASA began research on surgical robots with the aim of developing a remote operative system. It took nearly 30 years to complete the first fully functional surgical robot system. Called the Da Vinci Surgical System (DVSS), it has been clinically applied in the USA since 1997. In 1998, DVSS entered clinical trials and became commercially available in the USA. In 2000, DVSS was approved by the USA Food and Drug Administration. In 2001, a French surgeon, Jacques Marescaux, successfully performed the first transatlantic robotic-assisted cholecystectomy while working in the USA^[9]. In 2003, Talamini *et al.*^[10] reported the first series of transhiatal RAMIE. This was 8 years after the first transhiatal conventional MIE was reported by DePaula *et al.*^[11] in 1995. In 2004, Kernstine *et al.*^[12] reported the first series of transthoracic RAMIEs, which was 12 years after the first transthoracic conventional MIE was reported by Cuschieri *et al.*^[6] in 1992. Since then, RAMIE has been performed worldwide in many institutions. Moreover, given its many unique advantages, further clinical application of RAMIE is now being widely investigated. The history of RAMIE is summarized in Table 1.

CHARACTERISTICS OF THE OPERATIVE APPROACHES TO ESOPHAGECTOMY

MIE was introduced to improve outcomes and/or reduce the invasiveness of OE, and it has produced satisfactory results. In 2003, Luketich *et al.*^[13] reported the first large series of total MIEs and reported impressively low incidence of morbidity and mortality among 222 patients. Total MIE is performed by starting with a transthoracic MIE, followed by laparoscopic surgery to mobilize the stomach and perform upper abdominal lymphadenectomy. Transthoracic MIE provides improved magnified vision, less chest wall injury and relatively easy access to the upper thoracic structures, while laparoscopic surgery has less abdominal wall injury and less blood loss due to the pneumonic pressure. The first published randomized control trial in 2012, the TIME trial, is considered to be the cornerstone of MIE studies^[14]. Between 2009 and 2017, eight meta-analyses were published, comparing postoperative and oncologic outcomes of MIE and OE^[15]. MIE was generally found to be superior to OE in terms of intraoperative blood loss, acute immunological response, postoperative pulmonary infections, length of hospital stay, postoperative pain scores, and quality of life. Furthermore, the lymph node dissection (LND) yield and 3-year survival were equivalent^[14,16,17]. However, the two-dimension view, reduced eye-hand coordination, narrow operative field, restricted freedom of movement of operative instruments, moving targets, and nearby vital structures are all limitations such that MIE remains a highly complex procedure to be mastered by the surgeon^[8,18]. For example, the learning curve for an intrathoracic anastomosis was 119 cases when the incidence of

Table 2. Characteristics of each approach to esophagectomy

	OE	MIE	RAMIE
Difficulty level of technique	Relatively easy	Highly complex	Easier than MIE
Special points	Conventional operative method	Better vision	Zoomed-in enhanced three-dimensional vision
	with a lot of history	A two-dimensional view	Better overview
	Gold standard method	Reduced eye-hand coordination	Increased range of movement
		Restricted range of movement	Tremorless actions
			Flexible endo-wrists
Ergonomic conditions	Normal	Worst	Best
Blood loss	More	Less	Least
Operative time	Shorter	Longer	Longer
Postoperative pain score	High	Lower	Lower
Postoperative respiratory complications	More	Less	Less
Difficulty and exactness of upper mediastinum lymph node dissection	Difficult to access	More challenging maneuver than OE	Easier than MIE
	Equivalent	Equivalent	More exact
Postoperative recurrent laryngeal nerve paralysis	Equivalent	Equivalent	Reduced
Intrathoracic hand-sewn anastomosis	Difficult	The most difficult	Easy compared to MIE
Acute immunological response	More	Less	Same as total MIE
Functional recovery	Slowest	Fast	Same as total MIE
Length of hospital stay	Longest	Short	Same as total MIE
Mortality	Equivalent	Equivalent	Equivalent
Cost	Equivalent	Equivalent	Highest
Survival	Equivalent	Equivalent	Equivalent

OE: open esophagectomy; MIE: minimum invasive esophagectomy; RAMIE: robot-assisted minimally invasive esophagectomy

anastomotic leakage was the determining parameter (the anastomotic leakage rate dropped from 18.8% to 4.5%)^[18]. The learning phase of MIE was also considered to be a likely explanation for the higher re-operation rates as compared to OE in multiple population-based studies^[19-22]. This may explain the findings from a survey amongst esophageal surgeons in 2014, which showed that only 43% of the respondents reported MIE as their preferred approach^[23]. Indeed, due to its high technical complexity, MIE has not been adopted as the standard approach for esophageal cancer. These issues are summarized in Table 2.

A hybrid MIE (HMIE), which combines laparoscopy with a conventional thoracotomy, or combines a thoracoscopy with a conventional laparotomy, has been suggested as an alternative to total MIE^[24]. Messenger *et al.*^[25] reported that patients undergoing HMIE showed less mortality at both 30 (3.3% vs. 5.7%) and 90 days (6.9% vs. 10%) when compared to OE. In addition, Mariette *et al.*^[26] reported a randomized phase III trial (MIRO trial), which found that HMIE had a lower incidence of perioperative complications (36% vs. 64%), especially pulmonary complications (18% vs. 30%), with equivalent 3-year survival (67% vs. 55%) when compared to OE. Studies comparing HMIE with total MIE are scarce. In one study, however, Bonavina *et al.*^[27] compared a series of 80 total MIE versus 80 HMIE patients and found no differences in early postoperative complications or mortality. In addition, Grimminger *et al.*^[28] reported a series of 75 patients (HMIE 25, total MIE 25, RAMIE 25), which showed comparable morbidity and short-term outcomes in the three groups, although the total minimally invasive approaches appear to be associated with a lower incidence of complications such as pneumonia and wound infections. Those studies showed that although HMIE is a transitional operative method between OE and total MIE, because of its relatively lower difficulty level, somewhat reduced invasiveness and satisfactory clinical outcomes, it is a valuable operative method worth being performed.

To overcome the disadvantages of total MIE and HMIE, a robotic surgical system was developed and applied clinically. Transhiatal RAMIE was first introduced in 2003^[11], and transthoracic RAMIE

was introduced a year later^[12]. Although RAMIE is still under development, it is now described as a promising minimally invasive operative method with short-term and long-term clinical outcomes that are equivalent to (or perhaps better than) those achieved with OE and MIE [Table 2]^[29]. In a US report, 32.1% of esophageal cancer patients were treated with MIE. Of these, 19.6% were RAMIE^[30]. In that report, no differences in postoperative mortality or disease-free survival was noted between MIE and RAMIE^[30]. Nevertheless, given the many unique advantages of the robot, it is expected to decrease the morbidity and mortality rates of surgery for esophageal cancer and to improve oncological outcomes. Results of the recently published ROBOT trial showed improved clinical outcomes with reduced surgical and cardiopulmonary complication rates, reduced pain and improved functional outcomes with RAMIE as compared to OE^[31]. Moreover, RAMIE was associated with less intraoperative blood loss, lower postoperative pain scores, faster functional recovery, and better quality of life when compared to OE^[31]. Lymph node yield and overall survival did not differ between the two approaches, indicating that RAMIE offers short-term benefits while maintaining the high oncological standards. Needless to say, evidence remains weak due to limited RCT results, and more RCT studies are still needed.

Additionally, Yun *et al.*^[32] showed that RAMIE is also safe and feasible for use with patients who have received neoadjuvant chemoradiotherapy for locally advanced esophageal cancer, with postoperative mortality and morbidity rates comparable to that in OE. Another recently published study compared the clinical benefits of RAMIE with conventional OE. They showed that RAMIE could be a better surgical option for selected esophageal squamous cell carcinoma patients, offering both short-term and long-term benefits^[33]. Although both the short-term and long-term outcomes of RAMIE appear equivalent to MIE in most studies, one paper showed that RAMIE for esophageal cancer patients with node-positive disease in the superior mediastinum is associated with increased mortality (7.5%) and morbidity^[34].

LYMPH NODE DISSECTION IN RAMIE

The number of lymph nodes removed is a key factor contributing to the improved survival of esophageal cancer patients^[35]. LND along the recurrent laryngeal nerve (RLN) is considered beneficial; however, RLN LND is frequently complicated by RLN palsy (20%-80%), which is especially common on the left side. Early meta-analysis studies showed that, unfortunately, MIE does not reduce the rates of postoperative RLN palsy following RLN LND^[36-38]. On the other hand, RAMIE has several advantages for LND, especially RLN LND [Table 2]. The ROBOT trial showed that a mean of 27 and 25 lymph nodes were harvested in RAMIE and OE, respectively (not significantly different)^[31], which demonstrated that robotic surgery is at least comparable to open surgery for retrieving a sufficient number of lymph nodes. Although most early studies have found that the lymph node yield with RAMIE and MIE are similar^[39-41], in two recent series in which RAMIE and MIE were applied to upper mediastinal LND, markedly larger numbers of lymph nodes were harvested with RAMIE (median 37-49 vs. 19-21)^[42,43]. In addition, when Motoyama *et al.*^[44] compared the number of lymph nodes dissected from around the left RLN, they found that significantly more lymph nodes were dissected with RAMIE than MIE (median 6 vs. 4). This indicates that a robot-assisted surgical system may enable more extensive dissection of lymph nodes around the left RLN. Similarly, Park *et al.*^[42] demonstrated that the total number of dissected lymph nodes was significantly greater in the RAMIE group (37.3 ± 17.1 vs. 28.7 ± 11.8 ; $P = 0.003$), and intergroup differences were significant for the number of lymph nodes dissected from both the upper mediastinum (RAMIE: 10.7 ± 9.7 vs. MIE: 6.3 ± 9.3 , $P = 0.032$) and abdomen (RAMIE: 12.2 ± 8.7 vs. MIE: 7.8 ± 7.1 , $P = 0.007$). The five-year overall survival did not differ between the two groups (RAMIE: 69% vs. MIE: 59%, $P = 0.737$). Deng *et al.*^[45] showed that RAMIE may have an advantage for lymphadenectomy (mean: 20.6 ± 8.8 vs. 17.9 ± 7.7 ; $P = 0.048$) over MIE without increasing the risk of major postoperative complications. A recent propensity-matched analysis of patients undergoing modified Ivor Lewis esophagectomy also showed that the median total lymph node yield was 27 (range 13-84) in the RAMIE group compared to 23 in the MIE group (range 11-48). With a P -value of 0.053, their results suggest a trend towards improved lymphadenectomy with RAMIE^[46]. These studies

Table 3. Intrathoracic anastomosis in RAMIE

Intrathoracic anastomosis methods	Merits	Limitations
Hand-sewn	Can take full advantage of robot-assisted hand-sewing. Can be performed when the length is insufficient for staple anastomosis	Operative field is not satisfactory in the posterior wall anastomosis
Overlap (linear stapler × 1) + Hand-sewn	No need for additional mini-thoracotomy. Lower occurrence of stenosis. Can save stapler. Can take full advantage of robot-assisted hand-sewing	Cannot completely remove tissue poorly supplied with blood. Need a longer tubular stomach and esophageal end than circular stapler
Function (linear stapler × 2)	No need for additional mini-thoracotomy	Need a longer tubular stomach and esophageal end. Cannot completely remove tissue poorly supplied with blood
Triangular stapling (linear stapler × 3)	A reportedly lower rate of anastomotic complications. Lower occurrence of stenosis	The need to intrathoracically staple three times in three directions is a technical challenge
Circular stapler	Relatively easy to perform. Can completely remove tissue poorly supplied with blood	Need an additional mini-thoracotomy. Need an extra circular stapler. Higher occurrence of stenosis

RAMIE: robot-assisted minimally invasive esophagectomy

demonstrate that RAMIE may be more effective for extensive LND than MIE or OE. Recurrent nerve palsy is a complication that is especially associated with lymph node dissection in the superior mediastinum. In the ROBOT trial, the recurrent nerve palsy rate was 9%^[31]. However, Park *et al.*^[47] showed a significant learning curve on RLN palsy rates, which dropped from 55% to 0% after performing 20 cases in their study. The length of the learning curve for RAMIE has been reported to be 20-70 cases^[8,18].

ROBOTIC INTRATHORACIC ANASTOMOSIS

The robotic intrathoracic anastomosis can be hand-sewn or performed with linear or circular staplers. Although complete hand-sewing takes full advantage of robot assistance, it appears posterior wall anastomoses are technically challenging because of the deep and narrow operative field^[48]. Wang *et al.*^[49] showed side-to-side anastomosis to be a promising approach with the advantages of there being no need for additional mini-thoracotomy and a lower incidence of stenosis. In their report, the authors also emphasized the usefulness of the barbed knotless suture. Another recent study reported similar satisfactory outcomes with end-to-side anastomosis^[50]. Those authors concluded that end-to-side anastomosis requires a shorter length of the esophageal end, and section with poor blood supply was removed by a second stapler, which may ensure a good blood supply to the anastomosis. Triangular stapling is another anastomotic technique, which is reportedly associated with a lower rate of anastomotic complications^[51]. However, stapling three times in three directions would seem to present a great technical challenge intrathoracically. Recently, Han *et al.*^[52] reviewed diverse ways of intrathoracic anastomosis. Among these anastomotic methods, mortality was equivalent, but the anastomotic leak rates differed. Further large clinical trials are still needed. In general, each method has its merits and demerits. Surgeons should determine the anastomotic method of every single case with the final aim of maximizing patient benefits. The methods used for anastomosis in RAMIE are summarized in Table 3.

TRANSTHORACIC VS. TRANSHIATAL RAMIE

As with MIE, different variations of RAMIE have been established. Transthoracic RAMIE is one of the most commonly used approaches. It has a wide operative field, and after posterior and middle mediastinal LND, superior mediastinal LND can be performed in this operative field. However, destruction of the thoracic wall and pleura are unavoidable and differential lung ventilation is still needed. In 2003, Talamini *et al.*^[10] reported the first series of transhiatal RAMIE. Conventional transhiatal MIE has been proven as a less

Table 4. Transthoracic vs. transhiatal route in RAMIE

RAMIE	Merits	Limitations
Transthoracic	Wide operative field. Superior mediastinum lymph node dissection can be performed in the same operative field	Thoracic wall and pleura destruction are unavoidable. Differential lung ventilation is still needed in most case
Transhiatal	No need for thoracic wall destruction. No pleurotomy. No need of differential lung ventilation. No need for a change in body position. Almost no postoperative respiratory complications	Narrow operative field. Need a decent experience for mediastinum lymph node dissection under mediastinoscopy

RAMIE: robot-assisted minimally invasive esophagectomy

invasive operative method but oncologically inferior to radical esophagectomy^[53]. Although lymph node dissection of the lower mediastinal field is considered to be equivalent to radical esophagectomy, when it comes to the middle mediastinal field, it shows shortages because conventional endoscopic devices suffer from the paralleled right- and left-hand in the deep narrow operative fields. Meanwhile, the robot has articulated instruments and enhanced three-dimensional magnified view, which can move freely in the deep narrow cavity. It has been proven that RAMIE can overcome the limitations of the conventional transhiatal MIE and can dissect lymph nodes equivalent to radical esophagectomy^[54]. Yoshimura *et al.*^[55] showed that transhiatal RAMIE is associated with fewer pulmonary complications (0%) and better postoperative quality of life. However, it requires two LND steps. Posterior and middle mediastinal LND is performed using transhiatal RAMIE, followed by cervical mediastinoscopy for superior mediastinal LND. Mori *et al.*^[56] showed that the radicality of transmediastinal esophagectomy is equivalent to that of transthoracic esophagectomy in terms of the number of harvested lymph nodes and the pathology of surgical margins. Similarly, postoperative pneumonia did not occur in the transhiatal group. Although short-term and long-term outcomes were reported to not be inferior, due to the narrow operative field with the transhiatal procedure and mediastinoscopy, transhiatal RAMIE appears to be a more complex procedure. RAMIE operative routes are summarized in Table 4.

OPERATIVE POSITIONS IN RAMIE

Acute lung injury occurs in 25%-30% of patients after transthoracic esophagectomy, and single lung ventilation has been implicated in its pathogenesis^[57]. Until recently, RAMIE has been performed with the patient in the left lateral decubitus position in a setting of single-lung ventilation. Full lateral decubitus position with a cephalic parallel approach was reported to save some operative time (381 ± 57.7 min)^[58]. However, this approach requires total lung collapse and is therefore, often accompanied by serious pulmonary complications. To overcome the disadvantages of differential ventilation, Palanivelu *et al.*^[59] performed MIE with patients in a prone position. With their large patient cohort, they found that the prone position takes advantage of gravity to displace the lung from the dorsal thoracic structures and the esophagus, and that it has lower respiratory complications and shorter operative times due to the excellent exposure of the operative field and the better ergonomics for the surgeon. Sometimes, the vertebral column may obstruct the view of the operative field. Ruurda *et al.*^[60] reviewed the application of the prone position in RAMIE, with the patient cart of the robot system standing on the patient's side and extending its arms in a direction crossing the longitudinal axis of the patient. In the subsequent abdominal phase, the patient cart must be repositioned in front of the patient's head. This patient cart repositioning is time-consuming^[58]. On the other hand, urgent conversion to a classic thoracotomy, if needed, is probably more difficult with the prone position^[61]. As a solution to overcome this problem, whilst retaining the benefits of the prone position, a relatively complicated position, a modified semi-prone position has been adopted by surgeons around the world^[62]. Operative positions are summarized in Table 5.

Table 5. Operative positions in RAMIE

	Merits	Limitations
Left lateral decubitus position	Similar to open resection and does not require repositioning in the case of a conversion to open surgery	Need single-lung ventilation with more postoperative pulmonary complications.
Prone position	Takes advantage of gravity to displace the lung from the dorsal thoracic structures and the esophagus. Excellent exposure of the operative field. Allows for double-lung ventilation with less postoperative pulmonary complications	Need repositioning in the case of a conversion to open surgery
Semi-prone position	Has benefits of both the prone position and left lateral decubitus position	Relatively complicated

RAMIE: robot-assisted minimally invasive esophagectomy

Table 6. Prospects of RAMIE

1	Tactile function
2	Forceps tip shape change function
3	Automatic forceps switch function
4	Flexible camera
5	Artificial intelligence
6	Miniaturized operating robot body and wrist
7	Break the monopoly of the Da Vinci system

RAMIE: robot-assisted minimally invasive esophagectomy

PROSPECTS FOR RAMIE

Although RAMIE has a number of advantages that can overcome the shortcomings of MIE, there are still many problems that need to be resolved [Table 6]. For example, to perform surgery more safely, if possible we would like to add tactile function to the robot. To shorten the operative time, a forceps tip with shape changing function, automatic forceps switching function, and flexible camera are expected. Artificial intelligence is another exciting feature that is being developed. To reduce interference, we are looking forward to the development and manufacture of an operating robot with a miniaturized body and wrists. In addition, to break the monopoly of the Da Vinci system, many surgical robot companies worldwide are working on the development and manufacture of new robot surgery systems, which could bring lower costs.

DECLARATIONS

Authors' contributions

Made substantial contributions to the conception and design of the study and performed data analysis and interpretation: Liu J, Motoyama S

Performed data acquisition, as well as provided administrative, technical, and material support: Sato Y, Wakita A, Kawakita Y, Nagaki Y, Fujita H, Imai K, Minamiya Y

Availability of data and materials

Not applicable.

Financial support and sponsorship

None.

Conflicts of interest

All authors declared that there are no conflicts of interest.

Ethical approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Copyright

© The Author(s) 2020.

REFERENCES

1. Torre LA, Siegel RL, Ward EM, Jemal A. Global cancer incidence and mortality rates and trends--an update. *Cancer Epidemiol Biomarkers Prev* 2016;25:16-27.
2. Hulscher JB, van Sandick JW, de Boer AG, Wijnhoven BP, Tijssen JG, et al. Extended transthoracic resection compared with limited transhiatal resection for adenocarcinoma of the esophagus. *N Engl J Med* 2002;347:1662-9.
3. Takeuchi H, Miyata H, Gotoh M, Kitagawa Y, Baba H, et al. A risk model for esophagectomy using data of 5354 patients included in a Japanese nationwide web-based database. *Ann Surg* 2014;260:259-66.
4. Chen MF, Yang YH, Lai CH, Chen PC, Chen WC. Outcome of patients with esophageal cancer: a nationwide analysis. *Ann Surg Oncol* 2013;20:3023-30.
5. Kato H, Nakajima M. Treatments for esophageal cancer: a review. *Gen Thorac Cardiovasc Surg* 2013;61:330-5.
6. Cuschieri A, Shimi S, Banting S. Endoscopic oesophagectomy through a right thoracoscopic approach. *J R Coll Surg Edinb* 1992;37:7-11.
7. van Hillegersberg R, Boone J, Draaisma WA, Broeders IA, Giezeman MJ, et al. First experience with robot-assisted thoracoscopic esophagolymphadenectomy for esophageal cancer. *Surg Endosc* 2006;20:1435-9.
8. Kingma BF, de Maat MFG, van der Horst S, van der Sluis PC, Ruurda JP, et al. Robot-assisted minimally invasive esophagectomy (RAMIE) improves perioperative outcomes: a review. *J Thorac Dis* 2019;11:S735-42.
9. Marescaux J, Leroy J, Gagner M, Rubino F, Mutter D, et al. Transatlantic robot-assisted telesurgery. *Nature* 2001;413:379-80.
10. Talamini MA, Chapman S, Horgan S, Melvin WS; Academic robotics group. A prospective analysis of 211 robotic-assisted surgical procedures. *Surg Endosc* 2003;17:1521-4.
11. DePaula AL, Hashiba K, Ferreira EA, de Paula RA, Grecco E. Laparoscopic transhiatal esophagectomy with esophagogastroplasty. *Surg Laparosc Endosc* 1995;5:1-5.
12. Kernstine KH, DeArmond DT, Karimi M, Van Natta TL, Campos JH, et al. The robotic, 2-stage, 3-field esophagolymphadenectomy. *J Thorac Cardiovasc Surg* 2004;127:1847-9.
13. Luketich JD, Alvelo-Rivera M, Buenaventura PO, Christie NA, McCaughan JS, et al. Minimally invasive esophagectomy: outcomes in 222 patients. *Ann Surg* 2003;238:486-94; discussion 494-5.
14. Biere SS, van Berge Henegouwen MI, Maas KW, Bonavina L, Rosman C, et al. Minimally invasive versus open oesophagectomy for patients with oesophageal cancer: a multicentre, open-label, randomised controlled trial. *Lancet* 2012;379:1887-92.
15. van der Sluis PC, Schizas D, Liakakos T, van Hillegersberg R. Minimally Invasive Esophagectomy. *Dig Surg* 2020;37:93-100.
16. Maas KW, Biere SS, van Hoogstraten IM, van der Peet DL, Cuesta MA. Immunological changes after minimally invasive or conventional esophageal resection for cancer: a randomized trial. *World J Surg* 2014;38:131-7.
17. Straatman J, van der Wielen N, Cuesta MA, Daams F, Roig Garcia J, et al. Minimally invasive versus open esophageal resection: three-year follow-up of the previously reported randomized controlled trial: the TIME trial. *Ann Surg* 2017;266:232-6.
18. van Workum F, Stenstra MHBC, Berkelmans GHK, Slaman AE, van Berge Henegouwen MI, et al. Learning curve and associated morbidity of minimally invasive esophagectomy: a retrospective multicenter study. *Ann Surg* 2019;269:88-94.
19. Takeuchi H, Miyata H, Ozawa S, Udagawa H, Osugi H, et al. Comparison of short-term outcomes between open and minimally invasive esophagectomy for esophageal cancer using a nationwide database in Japan. *Ann Surg Oncol* 2017;24:1821-7.
20. Sihag S, Kosinski AS, Gaissert HA, Wright CD, Schipper PH. Minimally invasive versus open esophagectomy for esophageal cancer: a comparison of early surgical outcomes from the society of thoracic surgeons national database. *Ann Thorac Surg* 2016;101:1281-8; discussion 1288-9.
21. Seesing MFJ, Gisbertz SS, Goense L, van Hillegersberg R, Kroon HM, et al. A propensity score matched analysis of open versus minimally invasive transthoracic esophagectomy in the netherlands. *Ann Surg* 2017;266:839-46.
22. Mamidanna R, Bottle A, Aylin P, Faiz O, Hanna GB. Short-term outcomes following open versus minimally invasive esophagectomy for cancer in England: a population-based national study. *Ann Surg* 2012;255:197-203.
23. Haverkamp L, Seesing MF, Ruurda JP, Boone J, V Hillegersberg R. Worldwide trends in surgical techniques in the treatment of esophageal and gastroesophageal junction cancer. *Dis Esophagus* 2017;30:1-7.
24. Briez N, Piessen G, Bonnetain G, Brigand C, Carrere N, et al. Open versus laparoscopically-assisted oesophagectomy for cancer: a multicentre randomised controlled phase III trial - the MIRO trial. *BMC Cancer* 2011;11:310.
25. Messager M, Pasquer A, Duhamel A, Caranhac G, Piessen G, et al; FREGAT working group FRENCH. Laparoscopic gastric mobilization reduces postoperative mortality after esophageal cancer surgery: a french nationwide study. *Ann Surg* 2015;262:817-22; discussion 822-3.
26. Mariette C, Markar SR, Dabakuyo-Yonli TS, Meunier B, Pezet D, et al; Fédération de Recherche en Chirurgie (FRENCH) and French

- Eso-Gastric Tumors (FREGAT) Working Group. Hybrid minimally invasive esophagectomy for esophageal cancer. *N Engl J Med* 2019;380:152-62.
27. Bonavina L, Scolari F, Aiolfi A, Bonitta G, Sironi A, et al. Early outcome of thoracoscopic and hybrid esophagectomy: propensity-matched comparative analysis. *Surgery* 2016;159:1073-81.
 28. Grimminger PP, Tagkalos E, Hadzijušević E, Corvinus F, Babic B, et al. Change from hybrid to fully minimally invasive and robotic esophagectomy is possible without compromises. *Thorac Cardiovasc Surg* 2019;67:589-96.
 29. Biebl M, Andreou A, Chopra S, Denecke C, Pratschke J. Upper gastrointestinal surgery: robotic surgery versus laparoscopic procedures for esophageal malignancy. *Visc Med* 2018;34:10-5.
 30. Weksler B, Sullivan JL. Survival after esophagectomy: a propensity-matched study of different surgical approaches. *Ann Thorac Surg* 2017;104:1138-46.
 31. van der Sluis PC, van der Horst S, May AM, Schippers C, Brosens LAA, et al. Robot-assisted minimally invasive thoracoscopic esophagectomy versus open transthoracic esophagectomy for resectable esophageal cancer: a randomized controlled trial. *Ann Surg* 2019;269:621-30.
 32. Yun JK, Lee IS, Gong CS, Kim BS, Kim HR, et al. Clinical utility of robot-assisted transthoracic esophagectomy in advanced esophageal cancer after neoadjuvant chemoradiation therapy. *J Thorac Dis* 2019;11:2913-23.
 33. Yun JK, Chong BK, Kim HJ, Lee IS, Gong CS, et al. Comparative outcomes of robot-assisted minimally invasive versus open esophagectomy in patients with esophageal squamous cell carcinoma: a propensity score-weighted analysis. *Dis Esophagus* 2020;33:doz071.
 34. van der Horst S, de Maat MFG, van der Sluis PC, Ruurda JP, van Hillegersberg R. Extended thoracic lymph node dissection in robotic-assisted minimal invasive esophagectomy (RAMIE) for patients with superior mediastinal lymph node metastasis. *Ann Cardiothorac Surg* 2019;8:218-25.
 35. Visser E, van Rossum PSN, Ruurda JP, van Hillegersberg R. Impact of lymph node yield on overall survival in patients treated with neoadjuvant chemoradiotherapy followed by esophagectomy for cancer: a population-based cohort study in the Netherlands. *Ann Surg* 2017;266:863-9.
 36. Nagpal K, Ahmed K, Vats A, Yakoub D, James D, et al. Is minimally invasive surgery beneficial in the management of esophageal cancer? A meta-analysis. *Surg Endosc* 2010;24:1621-9.
 37. Sgourakis G, Gockel I, Radtke A, Musholt TJ, Timm S, et al. Minimally invasive versus open esophagectomy: meta-analysis of outcomes. *Dig Dis Sci* 2010;55:3031-40.
 38. Guo W, Ma X, Yang S, Zhu X, Qin W, et al. Combined thoracoscopic-laparoscopic esophagectomy versus open esophagectomy: a meta-analysis of outcomes. *Surg Endosc* 2016;30:3873-81.
 39. Weksler B, Sharma P, Moudgill N, Chojnacki KA, Rosato EL. Robot-assisted minimally invasive esophagectomy is equivalent to thoracoscopic minimally invasive esophagectomy. *Dis Esophagus* 2012;25:403-9.
 40. Suda K, Ishida Y, Kawamura Y, Inaba K, Kanaya S, et al. Robot-assisted thoracoscopic lymphadenectomy along the left recurrent laryngeal nerve for esophageal squamous cell carcinoma in the prone position: technical report and short-term outcomes. *World J Surg* 2012;36:1608-16.
 41. Zhang Y, Han Y, Gan Q, Xiang J, Jin R, et al. Early outcomes of robot-assisted versus thoracoscopic-assisted Ivor Lewis esophagectomy for esophageal cancer: a propensity score-matched study. *Ann Surg Oncol* 2019;26:1284-91.
 42. Park S, Hwang Y, Lee HJ, Park IK, Kim YT, et al. Comparison of robot-assisted esophagectomy and thoracoscopic esophagectomy in esophageal squamous cell carcinoma. *J Thorac Dis* 2016;8:2853-61.
 43. Park SY, Kim DJ, Do YW, Suh J, Lee S. The oncologic outcome of esophageal squamous cell carcinoma patients after robot-assisted thoracoscopic esophagectomy with total mediastinal lymphadenectomy. *Ann Thorac Surg* 2017;103:1151-7.
 44. Motoyama S, Sato Y, Wakita A, Kawakita Y, Nagaki Y, et al. Extensive lymph node dissection around the left laryngeal nerve achieved with robot-assisted thoracoscopic esophagectomy. *Anticancer Res* 2019;39:1337-42.
 45. Deng HY, Luo J, Li SX, Li G, Alai G, et al. Does robot-assisted minimally invasive esophagectomy really have the advantage of lymphadenectomy over video-assisted minimally invasive esophagectomy in treating esophageal squamous cell carcinoma? A propensity score-matched analysis based on short-term outcomes. *Dis Esophagus* 2019;32:doz110.
 46. Tagkalos E, Goense L, Hoppe-Lotichius M, Ruurda JP, Babic B, et al. Robot-assisted minimally invasive esophagectomy (RAMIE) compared to conventional minimally invasive esophagectomy (MIE) for esophageal cancer: a propensity-matched analysis. *Dis Esophagus* 2020;33:doz060.
 47. Park SY, Kim DJ, Kang DR, Haam SJ. Learning curve for robotic esophagectomy and dissection of bilateral recurrent laryngeal nerve nodes for esophageal cancer. *Dis Esophagus* 2017;30:1-9.
 48. Cerfolio RJ, Wei B, Hawn MT, Minnich DJ. Robotic esophagectomy for cancer: early results and lessons learned. *Semin Thorac Cardiovasc Surg* 2016;28:160-9.
 49. Wang F, Zhang H, Zheng Y, Wang Z, Geng Y, et al. Intrathoracic side-to-side esophagogastrostomy with a linear stapler and barbed suture in robot-assisted Ivor Lewis esophagectomy. *J Surg Oncol* 2019;120:1142-7.
 50. Wang Z, Zhang H, Wang F, Wang Y. Robot-assisted esophagogastric reconstruction in minimally invasive Ivor Lewis esophagectomy. *J Thorac Dis* 2019;11:1860-6.
 51. Noshiro H, Urata M, Ikeda O, Iwasaki H, Nabae T, et al. Triangulating stapling technique for esophagogastrostomy after minimally invasive esophagectomy. *Surgery* 2013;154:604-10.
 52. Han Y, Zhang Y, Li H. Application of different intrathoracic esophagogastric anastomosis in surgery for esophageal cancer. *Zhonghua Wei*

- Chang Wai Ke Za Zhi 2018;21:995-1000. (in Chinese)
53. Montenovio MI, Chambers K, Pellegrini CA, Oelschlager BK. Outcomes of laparoscopic-assisted transhiatal esophagectomy for adenocarcinoma of the esophagus and esophago-gastric junction. *Dis Esophagus* 2011;24:430-6.
 54. DeLong JC, Kelly KJ, Jacobsen GR, Sandler BJ, Horgan S, et al. The benefits and limitations of robotic assisted transhiatal esophagectomy for esophageal cancer. *J Vis Surg* 2016;2:156.
 55. Yoshimura S, Mori K, Yamagata Y, Aikou S, Yagi K, et al. Quality of life after robot-assisted transmediastinal radical surgery for esophageal cancer. *Surg Endosc* 2018;32:2249-54.
 56. Mori K, Yamagata Y, Aikou S, Nishida M, Kiyokawa T, et al. Short-term outcomes of robotic radical esophagectomy for esophageal cancer by a nontransthoracic approach compared with conventional transthoracic surgery. *Dis Esophagus* 2016;29:429-34.
 57. Baudouin SV. Lung injury after thoracotomy. *Br J Anaesth* 2003;91:132-42.
 58. Zhu Y, Ma L, Liu L, Lin Y. Application of full lateral decubitus position with cephalic parallel approach in robotic-assisted minimally invasive esophagectomy. *J Thorac Dis* 2019;11:3250-6.
 59. Palanivelu C, Prakash A, Senthilkumar R, Senthilnathan P, Parthasarathi R, et al. Minimally invasive esophagectomy: thoracoscopic mobilization of the esophagus and mediastinal lymphadenectomy in prone position--experience of 130 patients. *J Am Coll Surg* 2006;203:7-16.
 60. Ruurda JP, van der Sluis PC, van der Horst S, van Hilllegersberg R. Robot-assisted minimally invasive esophagectomy for esophageal cancer: a systematic review. *J Surg Oncol* 2015;112:257-65.
 61. Markar SR, Wiggins T, Antonowicz S, Zacharakis E, Hanna GB. Minimally invasive esophagectomy: lateral decubitus vs. prone positioning; systematic review and pooled analysis. *Surg Oncol* 2015;24:212-9.
 62. Seesing MFJ, Goense L, Ruurda JP, Luyer MDP, Nieuwenhuijzen GAP, et al. Minimally invasive esophagectomy: a propensity score-matched analysis of semiprone versus prone position. *Surg Endosc* 2018;32:2758-65.