


AGEING AND NEURODEGENERATIVE DISEASES



Review of evidence implicating the plasminogen activator system in blood-brain barrier dysfunction associated with Alzheimer's disease

Mei-Yun Tang, Fredric A. Gorin, Pamela J. Lein

Editor-in-Chief



Wei-Dong Le

Prof. Le is currently the deputy director of the Academic Committee and the director of the Institute of Neurology, Sichuan Academy of Medical Science-Sichuan Provincial People's Hospital. He is also a life-time Professor of Neurology and the director of Center for Translational Research on Neurological Diseases, the First Affiliated Hospital of Dalian Medical University. He has been engaged in clinical practice, teaching and research of neurology for almost 40 years, including 24 years of work in Baylor College of Medicine as a Professor of Neurology and Director of the Parkinson's Disease Research Center; and 10 years of working as a Professor of Neuroscience and Head of Neurogenomic Lab in the Institute of Health Science of Chinese Academy of Science jointed with Shanghai Jiao-Tong University School of Medicine.

Honorary Editors-in-Chief



Ted M. Dawson



David Rubinsztein

Journal Scope

High-quality original articles, reviews, case reports, commentaries are welcomed from ageing and neurodegenerative diseases, particularly on the basic mechanisms of ageing and their roles in the onset and progression of neurodegenerative diseases. The journal aims to report innovative research advances on the following topics:

1. The cellular and molecular mechanisms of ageing and the pathogenesis of neurodegenerative diseases;
2. The associations between neurodegenerative diseases and the biological bases of ageing with a focus on: genomic instability, epigenetic alterations, telomere attrition, protein degradation system failure, mitochondrial dysfunction, cellular senescence, nutrient sensing deregulation, stem cell exhaustion, intercellular communication impairment, etc.;
3. Translational research into prevention and treatment of age-related neurodegenerative diseases;
4. Mechanistic bases for epidemiological observations in aging-related neurodegenerative diseases.



Journal Homepage

EDITORIAL BOARD

Editor-in-Chief

Wei-Dong Le

Sichuan Academy of Medical Science-
Sichuan Provincial Hospital, China

Honorary Editors-in-Chief

Ted M. Dawson

Johns Hopkins University School of
Medicine, USA

David Rubinsztein

University of Cambridge, UK

Advisory Editors

Shu-Min Duan

Zhejiang University School of Medicine,
China

Kwok-Fai So

Jinan University, China

Erik Ch. Wolters

Neuroplast BV, Netherlands

Moussa B.H. Youdim

Technion-Israel Institute of Technology,
Israel

Xu Zhang

Chinese Academy of Sciences, China

Associate Editors

Mark Hallett

National Institutes of Health, USA

Heinz Reichmann

University of Dresden, Germany

Weihong Song

Wenzhou Medical University, China

Hua-Xi Xu

Medical College of Xiamen University,
China

Kristine Yaffe

University of California San Francisco,
USA

Youth Associate Editor

Peng Lei

Sichuan University, China

Editorial Board Members

Shilpa Buch

University of Nebraska Medical Center,
USA

Huaibin Cai

National Institutes of Health, USA

Raymond Chuen-Chung Chang

The University of Hong Kong, China

Sheng-Di Chen

Shanghai Jiao Tong University School of
Medicine, China

Veralice Meireles Sales de Bruin

Sleep and Biological Rhythms
Laboratory, Brazil

Hao Deng

Central South University, China

Dong-Sheng Fan

Peking University Third Hospital, China

David I. Finkelstein

The Florey Institute, Australia

Fen-Biao Gao

University of Massachusetts Medical
School, USA

Jau-Shyong Hong

Research Triangle Park, USA

Giuseppe Lanza

Oasi Research Institute - IRCCS, Italy

Hao Li

University of California, USA

Jia-Yi Li

China Medical University, China

Xiao-Jiang Li

Jinan University, China

Michael Maes

Chulalongkorn University, Thailand

Zi-Xu Mao

Emory University, USA

Kalipada Pahan

Rush University Medical Center, USA

Hanming Shen

University of Macau, China

Jun Tan

Guizhou Medical University, China

Yi Tang

Capital Medical University, China

MOURAD TAYEBI

School of Medicine, Australia

Guanghui Wang

Soochow University College of
Pharmaceutical Sciences, China

Yan-Jiang Wang

Third Military Medical University, China

Zhexing Wen

Emory University School of Medicine,
USA

Sara Xapelli

Universidade de Lisboa, Portugal

Ping-Yi Xu

First Affiliated Hospital of Guangzhou
Medical University, China

Qian Yang

Air Force Medical University, China

Feng-Wei Yu

National University of Singapore,
Singapore

Jintai Yu

Fudan University, China

Min-Ming Zhang

Zhejiang University School of Medicine,
China

Zhentao Zhang

Renmin Hospital of Wuhan University,
China

Youth Editorial Board Members

Sheng Chen

University School of Medicine, China

Jifeng Guo

Xiangya Hospital, China

Wei Luo

Zhejiang University, China

Louise Parr-Brownlie

University of Otago, New Zealand

GENERAL INFORMATION

About the Journal

AND is a peer-reviewed and open access multidisciplinary journal that publishes high-quality original articles, reviews, case reports, commentaries, letters to editor, etc. Ageing is a major risk factor for neurodegeneration, and the prevalence of ageing-related neurodegenerative diseases such as Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis, etc. continues to rise with the increased ageing population. Unfortunately, there are no effective treatments available for the age-related neurodegenerative diseases. Thus, to develop successful interventions, it is important to investigate the basic mechanisms of ageing and their roles in the onset and progression of neurodegenerative diseases. Therefore, we plan to launch this new journal, which is aimed to report innovative research advances on the following topics:

1. The cellular and molecular mechanisms of ageing and the pathogenesis of neurodegenerative diseases;
2. The associations between neurodegenerative diseases and the biological bases of ageing with a focus on: genomic instability, epigenetic alterations, telomere attrition, protein degradation system failure, mitochondrial dysfunction, cellular senescence, nutrient sensing deregulation, stem cell exhaustion, intercellular communication impairment, etc.;
3. Translational research into prevention and treatment of age-related neurodegenerative diseases;
4. Mechanistic bases for epidemiological observations in aging-related neurodegenerative diseases;
5. Psychology and Geriatric Mental Illness;
6. Neuropharmacology.

Information for Authors

Manuscripts should be prepared in accordance with Author Instructions.

Please check www.agneudisjournal.com/pages/view/author_instructions for details.

All manuscripts should be submitted online at <https://oaemesas.com/login?JournalId=and>

Copyright

The entire contents of the *AND* are protected under international copyrights. The journal, however, grants to all users a free, irrevocable, worldwide, perpetual right of access to, and a license to copy, use, distribute, perform and display the work publicly and to make and distribute derivative works in any digital medium for any reasonable purpose, subject to proper attribution of authorship and ownership of the rights. The journal also grants the right to make small numbers of printed copies for their personal use under the Creative Commons Attribution 4.0 License.

Copyright is reserved by © The Author(s) 2022.

Permissions

For information on how to request permissions to reproduce articles/information from this journal, please visit www.agneudisjournal.com.

Disclaimer

The information and opinions presented in the journal reflect the views of the authors and not of the journal or its Editorial Board or the Publisher. Publication does not constitute endorsement by the journal. Neither the *AND* nor its publishers nor anyone else involved in creating, producing or delivering the *AND* or the materials contained therein, assumes any liability or responsibility for the accuracy, completeness, or usefulness of any information provided in the *AND*, nor shall they be liable for any direct, indirect, incidental, special, consequential or punitive damages arising out of the use of the *AND*. The *AND*, nor its publishers, nor any other party involved in the preparation of material contained in the *AND* represents or warrants that the information contained herein is in every respect accurate or complete, and they are not responsible for any errors or omissions or for the results obtained from the use of such material. Readers are encouraged to confirm the information contained herein with other sources.

Publisher

OAE Publishing Inc.

245 E Main Street st112, Alhambra, CA 91801, USA

Website: www.oaepublish.com

Contacts

E-mail: editorialoffice@agneudisjournal.com

Website: www.agneudisjournal.com

CONTENTS

Review

A novel neuroprotective cholinesterase-monoamine oxidase inhibitor for treatment of dementia and depression in Parkinson's disease

*Wei Liu, Yuqiang Wang, Moussa B. H. Youdim**

Review of evidence implicating the plasminogen activator system in blood-brain barrier dysfunction associated with Alzheimer's disease

*Mei-Yun Tang, Fredric A. Gorin, Pamela J. Lein**

Recent developments in understanding brain aging: sex differences, mechanisms, and implications in diseases

Jing Yang, Jing Qu, Huan Ma**

Influence of sleep disruption on protein accumulation in neurodegenerative diseases

*Xiying Wang, Rui Wang, Jiada Li**

Review

Open Access



A novel neuroprotective cholinesterase-monoamine oxidase inhibitor for treatment of dementia and depression in Parkinson's disease

Wei Liu¹, Yuqiang Wang², Moussa B. H. Youdim³

¹Guangzhou Magpie Pharmaceuticals Co., Ltd., Nongxin Mansion, Guangzhou 510627, Guangdong, China.

²Institute of New Drug Research, International Cooperative Laboratory of Traditional Chinese Medicine Modernization and Innovative Drug Development of Chinese Ministry of Education, Jinan University College of Pharmacy, Guangzhou 510632, Guangdong, China.

³Eve Topf Centers of Excellence for Neurodegenerative Diseases Research, Faculty of Medicine, Technion-Israel Institute of Technology, Haifa 31096, Israel.

Correspondence to: Prof. Moussa B. H. Youdim, Eve Topf Centers of Excellence for Neurodegenerative Diseases Research, Faculty of Medicine, Technion-Israel Institute of Technology, 1 Efron St. P.O.B. 9697, Haifa 31096, Israel.
E-mail: youdim@tx.technion.ac.il

How to cite this article: Liu W, Wang Y, Youdim MBH. A novel neuroprotective cholinesterase-monoamine oxidase inhibitor for treatment of dementia and depression in Parkinson's disease. *Ageing Neur Dis* 2022;2:1.
<https://dx.doi.org/10.20517/and.2021.09>

Received: 20 Oct 2021 **First Decision:** 6 Dec 2021 **Revised:** 17 Dec 2021 **Accepted:** 5 Jan 2022 **Published:** 17 Jan 2022

Academic Editors: Weidong Le, Guanghui Wang **Copy Editor:** Yue-Yue Zhang **Production Editor:** Yue-Yue Zhang

Abstract

The current novel therapeutic approach suggests that multi-targeted compounds, with diverse biological activities but a single set of bioavailability and pharmacokinetics, will be significantly more advantageous in the treatment of the complex pathology of Parkinson's diseases (PD) than traditional therapies. This review introduces a novel cholinesterase (ChE)-monoamine oxidase (MAO) inhibitor, namely MT-031, which was designed by amalgamating the propargyl moiety of the irreversible selective MAO-B inhibitor and neuroprotective/neurorestorative anti-Parkinsonian drug, rasagiline, into the methylamino position of the ChE inhibitor anti-AD drug, rivastigmine. MT-031 possesses neuroprotective, cognition enhancing, anti-depressant, and anti-inflammatory properties both *in vitro* and *in vivo*. Altogether, these findings suggest that MT-031 may be a potential treatment for combating PD and associated dementia and depression.



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, sharing, adaptation, distribution and reproduction in any medium or format, for any purpose, even commercially, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.



Keywords: Parkinson's disease, dementia, cholinesterase, monoamine oxidase, multi-targeted drugs

INTRODUCTION

With aging and the increasing life span of the population, Parkinson's disease (PD), an age-related neurodegenerative disorder, is receiving increased attention. It is estimated that the number of PD patients will reach more than 12 million by 2040, doubling the cases seen in 2016^[1]. The motor deficits of PD are emphasized in both making the initial diagnosis and in tracking the progression of the disease^[2]. As understanding of the symptoms and pathogenesis deepens, however, it has been suggested that the non-motor features of PD, including cognitive impairment, i.e., dementia, should be more attended to^[3,4]. A previous study indicated that approximately 25.8% of individuals with PD exhibit mild cognitive impairment^[5], and longitudinal studies have documented that up to 70% of these patients will progress to dementia after ten years of symptoms^[3]. In addition to cognitive impairment, other symptoms, e.g., depression, may emerge regularly throughout the development of PD^[6-8], and this symptom may worsen the severity of dementia as the disease progresses. Since dementia in both Alzheimer's disease (AD) and PD patients generally presents with similar features, present treatments for Parkinson's disease dementia (PDD) are mostly derived from drugs utilized in AD, such as cholinesterase inhibitors (ChEIs) and memantine, which was initially developed for the treatment of AD. To date, rivastigmine is the only FDA-approved therapy that is currently licensed for PDD.

It is well known that neurodegenerative diseases, such as AD, PD, amyotrophic lateral sclerosis, and Huntington's disease, are possibly triggered by a group of pathologies, characterized by separate etiologies with distinct morphological and pathophysiological features, including iron accumulation^[9-11], generation of reactive oxygen^[11] and nitrogen species^[12], inflammation^[13-15], mitochondrial (complex I) deficiency^[16], ubiquitin-proteasome system dysfunction^[17], and abnormal protein folding and aggregation^[18,19]. This suggests that the "cocktail of drugs" strategy, i.e., mixing different targeted molecules as drug combinations, may offer theoretically feasible treatment for these diseases. Nonetheless, compared to using a single effective compound, the cocktail strategy increases the risk of side effects and ups the difficulty of managing drug-drug interactions, safe dosing, and metabolic shunt effects^[20,21]. A single drug with multiple targets - one compound conjugating two or more diverse biological properties - thus has a pronounced advantage over single-target drugs or drug cocktails^[22,23]. An attractive example of a multi-targeted drug is ladostigil (TV3326), a cholinesterase (ChE)-monoamine oxidase (MAO) inhibitor, indicated to target various pathogenic mechanisms of neurodegenerative diseases^[24-27]. The underlying principle in the design of ladostigil was to join the carbamate ChE inhibitory moiety of the anti-AD drug, rivastigmine, to the irreversible selective MAO-B inhibitor, rasagiline^[24]. Ladostigil has shown positive results in a phase II clinical trial evaluating its safety and efficacy in patients diagnosed with MCI^[28].

Based on a similar rationale, a novel ChE-MAO inhibitor, namely MT-031 [Figure 1], was designed and synthesized for the treatment of AD. MT-031 amalgamates the propargyl moiety of the irreversible selective MAO-B inhibitor and neuroprotective/neurorestorative drug, rasagiline, into the methylamino position of the ChE inhibitor, rivastigmine^[29]. Since AD and PD share similar pharmacological treatment demands, this review discusses the potential use of this novel multi-targeted drug, MT-031, for dementia and depression in PD.

INHIBITORY EFFECT OF MT-031 ON MAO

Rasagiline (Azilect®) is an anti-Parkinsonian MAO-B inhibitor drug, which presented neuroprotective and neurorescue activities in animal models and neuronal cell models of neurodegeneration^[30] and exerted




Figure 1. The chemical structure of the novel ChE-MAO inhibitor, MT-031[(S)-3-(1-(Methyl(prop-2-yn-1-yl)amino)ethyl)phenyl ethyl(methyl)carbamate], designed by amalgamating the active propargyl moiety of the anti-Parkinsonian drug, rasagiline, a brain selective MAO-B inhibitor, into the “N-methyl” position of the anti-AD drug ChE inhibitor, rivastigmine. AD: Alzheimer’s disease; ChE: cholinesterase; MAO: monoamine oxidase; MAO-B: monoamine oxidase-B.

disease-modifying effects in PD patients^[30-32]. The propargyl moiety of rasagiline has been proven to be an important active functional group for its MAO inhibitory activity^[33,34] and neuroprotective/neurorestorative effects^[35,36]. By retaining the active propargyl moiety, the inhibition of MAO in the brain is associated with neuroprotective effects in the neurodegenerative and age-related disturbances of homeostasis, and the products of the MAO-catalyzed reaction (e.g., aldehydes and hydrogen peroxide) are compelling inducers of lipid peroxidation and the generation of free radicals in the involution of the nervous system^[37,38]. By retaining the propargyl moiety of rasagiline, MT-031 was found to be a selective MAO-A inhibitor (selectivity of MAO-A/B > 500-fold, Table 1); interestingly, this is different from its parent drug, rasagiline, which is a selective MAO-B inhibitor (selectivity of MAO-B/A = 100-fold, Table 1)^[29]. In humans, MAO-A is found within the outer mitochondrial membrane of both neuronal and glial cells, where it participates in the inactivation of dopamine (DA) in the primate and human brain^[39]. As dopamine depletion in the striatum causes the core motor manifestations of PD, a selective MAO-A inhibitor might provide an anti-Parkinsonian benefit^[40,41].

Additionally, depression has also been reported to be one of the most common symptoms of PD, occurring in around 40% of patients with PD, and it is often persistent^[42]. The efficacy of MAO-A inhibitors has been proven effective in the treatment of atypical depression, high levels of anxiety, anergic bipolar depression, and treatment-resistant depression for decades^[43-45]. MAO-A mainly metabolizes serotonin (5-HT) and norepinephrine (NE), and a reduction in the 5-HT major metabolite, 5-hydroxyindoleacetic acid, in the cerebrospinal fluid was reported to be associated with violent and impulsive behavior, including violent suicide attempts^[46]. The antidepressant effects of MAOIs were hypothesized to be based on a deficiency in catecholamines, specifically NE and DA, as well as possibly the indolamine 5-HT^[47]; the mechanisms of action of MAOIs as antidepressants were thus thought to be because they directly resulted in increased levels of neurotransmitter amines at nerve terminals^[48,49]. Selective MAO-B inhibitors may not be effective as antidepressants because MAO-B has no direct effect on either 5-HT or NE metabolism. A dual MAO-A/B inhibitor may rapidly increase DA levels to heighten feelings of pleasure, but abnormal surges in DA are linked to serious side effects^[50-53]. Therefore, a drug with selective MAO-A inhibition could potentially be a safer and more effective treatment for depression in PD patients.

Table 1. The inhibitory effect (IC₅₀) of MT-031 and its parent drugs, rasagiline and rivastigmine, on MAO and ChE *in vitro*

Compound	Inhibition (IC ₅₀ μM ^a)					
	MAO-A	MAO-B	MAO selectivity (A/B)	AChE	BuChE	ChE selectivity (AChE/BuChE)
MT-031	0.71 ± 0.04	> 1000	> 500	58.3 ± 6.3	34.6 ± 8.3	0.59
Rasagiline	0.41	0.0044	0.01	NA	NA	-
Rivastigmine	NA ^b	NA	- ^c	2.07	0.37	0.18

^aIC₅₀, micromolar (μM) concentration at which compound inhibits 50% of the enzyme activity; ^bNA, no activity; ^c-, not tested. MAO: Monoamine oxidase; MAO-A: monoamine oxidase-A; MAO-B: monoamine oxidase-B; ChE: cholinesterase; AChE: acetylcholinesterase; BuChE: butyrylcholinesterase.

Moreover, an important finding is that, following administration of MT-031, there is little inhibition of MAO-A in the liver and small intestine^[29,54]. Irreversible, high degrees of MAO-A inhibition in peripheral tissues is associated with potentiation of tyramine-induced cardiovascular activity^[55], namely the “cheese effect”^[56,57]. These data indicate that MT-031 may produce only limited potentiation of blood pressure in response to oral tyramine, as previously described for rasagiline^[57,58] and other propargyl containing drugs, such as ladostigil^[59], M30^[60], and VAR-10303^[61].

INHIBITORY EFFECT OF MT-031 ON CHE

To date, acetylcholinesterase inhibitors (AChEIs) have been the mainstay of therapeutic approaches for AD. AChEIs are used to increase synaptic levels of acetylcholine (ACh) and block the breakdown of ACh by inhibiting AChE^[62]. Some reports suggest that cortical cholinergic deficits are more pronounced in PDD and that they are strongly correlated with cognitive decline and neuropsychiatric disturbances in PD^[63,64]. The efficacy of the only FDA approved dual AChE and butyrylcholinesterase (BuChE) inhibitor, rivastigmine [Figure 1 and Table 1], one of the parent drugs of MT-031, has been proved in various clinical trials in the treatment of PDD^[65]. Rivastigmine exerts its therapeutic effects by increasing the levels of acetylcholine in the brain via reversible inhibition of its hydrolysis^[66]. It has been proposed that the effects of rivastigmine might reflect an additional property of BuChE inhibition, which is implicated in symptom progression and thus can provide some patients supplementary benefits over AChE selectivity^[67]. In humans, AChE predominates (80%) and BuChE is considered to play a minor role in regulating ACh levels in the healthy brain^[68]. Especially, BuChE activity rises while AChE activity remains unchanged or declines in the AD brain^[68-70], thereby supporting the key role of BuChE in regulating brain acetylcholine levels^[71]. Therefore, both enzymes are likely to be involved in regulating ACh levels and represent legitimate therapeutic targets to ameliorate cholinergic deficits^[72]. MT-031 was found to significantly inhibit both AChE and BuChE activities *in vitro*, although with a lower IC₅₀ than that of its parent drug, rivastigmine [Table 1]^[29]. Accordingly, our previous study showed that MT-031 treatment prevented cognitive deficits induced by scopolamine and improved spatial learning and memory. These results may be attributed to MT-031 being able attenuate scopolamine-induced ChE disturbance by inhibition of ChE activity. In addition, after acute treatment in rats, MT-031 inhibited cortical and hippocampal AChE/BuChE by 50%-70% at doses ranging from 5 to 10 mg/kg^[29]. The high inhibitory effect of ChE activity is very crucial, as the fact that the clinical study of ladostigil (clinicaltrials.gov/ct2/show/NCT01354691) in the treatment of AD did not achieve its primary outcome may be due to its low inhibitory ratio on AChE (ladostigil inhibited an average of 21.3% of AChE)^[28,73]. Furthermore, 24 h after the last dose was given to mice in a chronic administration model, MT-031 still caused dose-dependent antagonism of the spatial memory deficits induced by scopolamine in mice^[54]. These results may suggest that MT-031 is a reversible but long-term ChE inhibitor, and that it is able to increase brain ACh levels sufficiently to compete with scopolamine for the muscarinic receptors subserving memory^[74].

NEUROPROTECTIVE ACTIVITY OF MT-031

One aspect of the neuroprotective activity of MT-031 is that it directly scavenges free radicals over-produced in hydrogen peroxide (H_2O_2)-treated SH-SY5Y cells^[29]. H_2O_2 is a major source of free radicals; it is produced during the redox process and considered to be a messenger in intracellular signaling cascades, including cellular metabolism and proliferation^[75,76]. The predominant sources of H_2O_2 in the brain are spontaneous superoxide dismutation catalyzed by the enzyme superoxide dismutase^[77] and MAO activity^[78]. MAO-A and -B, in particular, catalyze the oxidative deamination of DA, 5-HT, and NE^[39] and yield metabolic products, aldehydes, and reactive oxygen species (ROS) such as H_2O_2 . Therefore, the neuroprotective abilities of MAO inhibitors in the treatment of PD may be through reducing ROS production^[39,79,80]. In addition, several lines of evidence suggest that AChE and BuChE activation may be involved in the apoptosis associated with H_2O_2 ^[81,82]. The link between cholinergic signaling and oxidative stress provides an additional therapeutic target for ChEIs in PD. Indeed, the ChEIs, tacrine^[81], huperzine A^[83], and rivastigmine^[84] were demonstrated to significantly protect cells against H_2O_2 insult. Moreover, MT-031 was found to enhance the mRNA expression levels of neurotrophins, anti-apoptotic molecules (Bcl-2 like 1 and Bcl-2), and an anti-oxidative enzyme (catalase) in the mouse striatum, further demonstrating the significant neuroprotective and anti-oxidative actions of this drug^[54]. Multiple studies with various apoptotic paradigms have shown that Bcl-2 can protect cells against oxidative insults^[85-88]. Measurements of ROS levels including H_2O_2 have shown that Bcl-2 expression is correlated with reduced levels of oxidative stress in cells exposed to oxidative damage. Additionally, increased synaptic ACh levels resulting from AChE inhibition may potentiate the effect of neurotrophins, neuronal growth factor and brain-derived neurotrophic factor, which was previously demonstrated to induce neuroprotection against free radical insults^[89,90].

Increasing evidence suggests that neuroinflammation contributes to the cascade leading to progressive neuronal damage in PD^[15,91]. The major pro-inflammatory cytokines, such as interleukin-1 β (IL-1 β), IL-2, IL-6, IL-17, tumor necrosis factor- α (TNF- α), and interferon- γ (IFN- γ), lead to increased production of inducible oxidative stress, neuronal stress, and further neuronal dysfunction and death in the AD brain^[92-95]. The anti-inflammatory effect of MT-031 was found to be associated with elevation of the levels of one of the major cytokines, IL-10, which limits inflammation by reducing the synthesis of pro-inflammatory cytokines such as IL-1, IL-6, IFN- γ , and TNF- α ^[54]. The anti-inflammatory effect of MT-031 was also demonstrated in proliferated splenocytes activated by anti-CD3, in which MT-031 did not affect the viability of the unstimulated splenocytes, indicating that the anti-proliferative effect was not associated with a protective effect against cytotoxicity^[54]. In addition to proliferation, splenocytes and microglia cells can also be activated to produce cytokines, multi-functional soluble factors with pro- and anti-inflammatory activities^[96,97]. MT-031 suppressed the elevation of IL-17 and INF- γ in anti-CD3-activated splenocytes, possibly by increasing the generation of IL-2, although the exact mechanism needs to be addressed by further study. Inconsistent with the anti-inflammatory effects seen in cell cultures, MT-031 upregulated the mRNA expression levels of the anti-inflammatory cytokine neurotrophic tyrosine kinase receptor and reduced levels of the pro-inflammatory cytokine IL-6 in a scopolamine mouse model^[54].

EFFECTS OF MT-031 ON SCOPOLAMINE-INDUCED DEMENTIA

It has been shown that scopolamine exerts its effects through antagonizing muscarinic acetylcholine receptors^[98,99]. A previous study confirmed that MT-031 treatment prevented cognitive deficits induced by scopolamine and improved spatial learning and memory, as examined in the Y maze task and Morris water maze test^[54]. This effect may be attributed to an increase of amine contents, NE, 5-HT, and DA, as well as to the direct effect on scopolamine-induced ChE disturbance through inhibition of ChE activity. MT-031 exerted a significant inhibitory effect on ChE in the hippocampus and frontal cortex of mice^[54]. This is an




Figure 2. Suggestive schematic illustration for the mechanism of multifunctional brain permeable drug, MT-031, as a potential therapeutic approach of dementia and depression in PD. PD: Parkinson's disease; AChE: acetylcholinesterase; BuChE: butyrylcholinesterase; MAO-A: monoamine oxidase-A; ROS: reactive oxygen species; IL: interleukin; TNF- α : tumor necrosis factor-alpha; IFN- γ : interferon-gamma; TNF- α : tumor necrosis factor-alpha; Ntrk: tyrosine kinase receptor; NGF: neuronal growth factor; BDNF: brain-derived neurotrophic factor; GDNF: glial cell-derived neurotrophic factor; Bcl-2 like 1: B-cell lymphoma 2 like 1.

advantageous property of MT-031, as previous data show that, when ChE inhibitors are less effective in the hippocampus, other brain regions may produce insufficient amounts of ACh to displace scopolamine from receptors, which results in dysfunctional mediation of working memory^[100]. Our data are in line with the reported protective effects of rivastigmine^[101] and ladostigil^[102] in a scopolamine mouse model, suggesting the importance of inhibiting both AChE and BuChE activities in ameliorating cognitive impairments^[65,101]. There are more and more studies that support the idea that multi-targeted brain selective MAO and ChE inhibitors may exert better treatment effects than single ChE inhibitors in the treatment of dementia in neurodegenerative disorders such as AD and PD^[22,26,80].

CONCLUSION AND PERSPECTIVE

Available treatments for PDD are limited in both number and quality, and they only provide symptomatic relief for cognitive impairment. The multi-factorial causes of the disease make the development of new drugs a difficult task. The rational design of incorporating two or more distinct functional pharmacophores into one molecule has been suggested to be feasible^[22,103]. A single target molecule may have greater affinity towards a specific target than a molecule with multiple targets; however, a multi-target strategy creates compounds with a balanced affinity for treating the multifactorial causes of multiple neurodegenerative diseases. To date, none of the cholinesterase inhibitors in the clinic has been proved to possess neuroprotective activity or anti-depressant action. The design of the novel drug candidate, MT-031, was aimed at targeting multiple neurodegenerative processes. MT-031 is a brain selective MAO-A and AChE/BuChE inhibitor and has been found to exert a wide range of neuroprotective activities [Figure 2], including anti-oxidative activity, clearance of ROS accumulation, prevention of neuronal death, and increasing levels of neurotrophic factors. MT-031 also possesses anti-inflammatory capabilities including

preventing cellular proliferation, upregulating anti-inflammatory cytokines, and downregulating pro-inflammatory cytokines^[29,54]. There is evidence that MT-031 inherited the neuroprotective potency described for propargylamine derivatives in neurodegenerative animal models^[29,54,104]. Similar to its other parent compound rivastigmine^[101] at a dose that inhibited ChE in the cortex and hippocampus by approximately 70%, MT-031 was effective in antagonizing the working and reference memory deficits induced by scopolamine^[54]. These miscellaneous pharmacological properties of MT-031 [Figure 2], accompanied by its ability to improve cognitive deficits, make this compound valuable as a novel drug candidate for the treatment of dementia and depression in PD.

DECLARATIONS

Acknowledgments

The authors gratefully acknowledge the support of the Rappaport Family Research Institute, Technion-Israel Institute of Technology (Haifa, Israel). The authors also thank Ms. Linda Wang for editing this manuscript.

Authors' contributions

Wrote the review paper: Liu W

Checked the review paper: Wang Y, Youdim MBH

Availability of data and material

Not applicable.

Financial support and sponsorship

The work was supported by Youdim Pharmaceuticals.

Conflicts of interest

All authors declared that there are no conflicts of interest.

Ethical approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Copyright

© The Author(s) 2022.

REFERENCES

1. Dorsey ER, Elbaz A, Nichols E, et al. Global, regional, and national burden of Parkinson's disease, 1990-2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet Neurol* 2018;17:939-53. DOI PubMed PMC
2. Gibb WR, Lees AJ. The relevance of the Lewy body to the pathogenesis of idiopathic Parkinson's disease. *J Neurol Neurosurg Psychiatry* 1988;51:745-52. DOI PubMed PMC
3. Chaudhuri KR, Martinez-Martin P, Schapira AH, et al. International multicenter pilot study of the first comprehensive self-completed nonmotor symptoms questionnaire for Parkinson's disease: the NMSQuest study. *Mov Disord* 2006;21:916-23. DOI PubMed
4. Zečević I. Clinical practice guidelines based on evidence for cognitive-behavioural therapy in Parkinson's disease comorbidities: a literature review. *Clin Psychol Psychother* 2020;27:504-14. DOI PubMed
5. Aarsland D, Bronnick K, Williams-Gray C, et al. Mild cognitive impairment in Parkinson disease: a multicenter pooled analysis. *Neurology* 2010;75:1062-9. DOI PubMed PMC
6. Hely MA, Reid WG, Adena MA, Halliday GM, Morris JG. The Sydney multicenter study of Parkinson's disease: the inevitability of dementia at 20 years. *Mov Disord* 2008;23:837-44. DOI PubMed
7. Duncan GW, Khoo TK, Yarnall AJ, et al. Health-related quality of life in early Parkinson's disease: the impact of nonmotor symptoms. *Mov Disord* 2014;29:195-202. DOI PubMed

8. Goodarzi Z, Mrklas KJ, Roberts DJ, Jette N, Pringsheim T, Holroyd-Leduc J. Detecting depression in Parkinson disease: a systematic review and meta-analysis. *Neurology* 2016;87:426-37. DOI PubMed PMC
9. Riederer P, Monoranu C, Strobel S, Iordache T, Sian-Hülsmann J. Iron as the concert master in the pathogenic orchestra playing in sporadic Parkinson's disease. *J Neural Transm (Vienna)* 2021;128:1577-98. DOI PubMed PMC
10. Genoud S, Senior AM, Hare DJ, Double KL. Meta-analysis of copper and iron in Parkinson's disease brain and biofluids. *Mov Disord* 2020;35:662-71. DOI PubMed
11. Van Houten B, Woshner V, Santos JH. Role of mitochondrial DNA in toxic responses to oxidative stress. *DNA Repair (Amst)* 2006;5:145-52. DOI PubMed
12. Liguori I, Russo G, Curcio F, et al. Oxidative stress, aging, and diseases. *Clin Interv Aging* 2018;13:757-72. DOI PubMed PMC
13. Ransohoff RM. How neuroinflammation contributes to neurodegeneration. *Science* 2016;353:777-83. DOI PubMed
14. Stephenson J, Nutma E, van der Valk P, Amor S. Inflammation in CNS neurodegenerative diseases. *Immunology* 2018;154:204-19. DOI PubMed PMC
15. Hirsch EC, Standaert DG. Ten unsolved questions about neuroinflammation in Parkinson's disease. *Mov Disord* 2021;36:16-24. DOI PubMed
16. de Moura MB, dos Santos LS, Van Houten B. Mitochondrial dysfunction in neurodegenerative diseases and cancer. *Environ Mol Mutagen* 2010;51:391-405. DOI PubMed
17. Paul S. Dysfunction of the ubiquitin-proteasome system in multiple disease conditions: therapeutic approaches. *Bioessays* 2008;30:1172-84. DOI PubMed
18. Dobson CM. Protein aggregation and its consequences for human disease. *Protein Pept Lett* 2006;13:219-27. DOI PubMed
19. Chiti F, Dobson CM. Protein misfolding, amyloid formation, and human disease: a summary of progress over the last decade. *Annu Rev Biochem* 2017;86:27-68. DOI PubMed
20. Hopkins AL, Mason JS, Overington JP. Can we rationally design promiscuous drugs? *Curr Opin Struct Biol* 2006;16:127-36. DOI PubMed
21. Savelieff MG, Nam G, Kang J, Lee HJ, Lee M, Lim MH. Development of multifunctional molecules as potential therapeutic candidates for Alzheimer's disease, Parkinson's disease, and amyotrophic lateral sclerosis in the last decade. *Chem Rev* 2019;119:1221-322. DOI PubMed
22. Youdim MB. Why do we need multifunctional neuroprotective and neurorestorative drugs for Parkinson's and Alzheimer's diseases as disease modifying agents. *Exp Neurobiol* 2010;19:1-14. DOI PubMed PMC
23. Van der Schyf CJ, Geldenhuys WJ. Multimodal drugs and their future for Alzheimer's and Parkinson's disease. *Int Rev Neurobiol* 2011;100:107-25. DOI PubMed
24. Weinstock M, Bejar C, Wang R, et al. TV3326, a novel neuroprotective drug with cholinesterase and monoamine oxidase inhibitory activities for the treatment of Alzheimer's disease. In: Riederer P, Calne DB, Horowski R, Mizuno Y, Olanow CW, Poewe W, Youdim MBH, editors. *Advances in Research on Neurodegeneration*. Vienna: Springer; 2000. p. 157-69. DOI PubMed
25. Weinstock M, Poltyrev T, Bejar C, Youdim MB. Effect of TV3326, a novel monoamine-oxidase cholinesterase inhibitor, in rat models of anxiety and depression. *Psychopharmacology (Berl)* 2002;160:318-24. DOI PubMed
26. Weinreb O, Amit T, Bar-Am O, Youdim MB. Ladostigil: a novel multimodal neuroprotective drug with cholinesterase and brain-selective monoamine oxidase inhibitory activities for Alzheimer's disease treatment. *Curr Drug Targets* 2012;13:483-94. DOI PubMed
27. Weinstock M, Bejar C, Schorer-Apelbaum D, Panarsky R, Luques L, Shoham S. Dose-dependent effects of ladostigil on microglial activation and cognition in aged rats. *J Neuroimmune Pharmacol* 2013;8:345-55. DOI PubMed
28. Schneider LS, Geffen Y, Rabinowitz J, et al; Ladostigil Study Group. Low-dose ladostigil for mild cognitive impairment: a phase 2 placebo-controlled clinical trial. *Neurology* 2019;93:e1474-84. DOI PubMed PMC
29. Liu W, Lang M, Youdim MBH, et al. Design, synthesis and evaluation of novel dual monoamine-cholinesterase inhibitors as potential treatment for Alzheimer's disease. *Neuropharmacology* 2016;109:376-85. DOI PubMed
30. Youdim MB, Bar Am O, Yorgev-Falach M, et al. Rasagiline: neurodegeneration, neuroprotection, and mitochondrial permeability transition. *J Neurosci Res* 2005;79:172-9. DOI PubMed
31. Olanow CW, Rascol O, Hauser R, et al; ADAGIO Study Investigators. A double-blind, delayed-start trial of rasagiline in Parkinson's disease. *N Engl J Med* 2009;361:1268-78. DOI PubMed
32. Hauser RA, Li R, Pérez A, et al; NINDS NET-PD Investigators. Longer duration of MAO-B inhibitor exposure is associated with less clinical decline in Parkinson's disease: an analysis of NET-PD LS1. *J Parkinsons Dis* 2017;7:117-27. DOI PubMed
33. Knoll J. [History of deprenyl--the first selective inhibitor of monoamine oxidase type B]. *Vopr Med Khim* 1997;43:482-93. PubMed
34. Youdim MB. Rasagiline: an anti-Parkinson drug with neuroprotective activity. *Expert Rev Neurother* 2003;3:737-49. DOI PubMed
35. Weinreb O, Amit T, Bar-Am O, Chillag-Talmor O, Youdim MB. Novel neuroprotective mechanism of action of rasagiline is associated with its propargyl moiety: interaction of Bcl-2 family members with PKC pathway. *Ann N Y Acad Sci* 2005;1053:348-55. DOI PubMed
36. Weinreb O, Amit T, Bar-am O, Sagi Y, Mandel S, Youdim MBH. Involvement of multiple survival signal transduction pathways in the neuroprotective, neurorescue and APP processing activity of rasagiline and its propargyl moiety. In: Riederer P, Reichmann H, Youdim MBH, Gerlach M, editors. *Parkinson's disease and related disorders*. Vienna: Springer; 2006. p. 457-65. DOI PubMed
37. Kumar MJ, Andersen JK. Perspectives on MAO-B in aging and neurological disease: where do we go from here? *Mol Neurobiol* 2004;30:77-90. DOI PubMed
38. Shemyakov SE. Monoamine oxidase activity, lipid peroxidation, and morphological changes in human hypothalamus during aging.

- Bull Exp Biol Med* 2001;131:586-8. DOI PubMed
39. Youdim MB, Edmondson D, Tipton KF. The therapeutic potential of monoamine oxidase inhibitors. *Nat Rev Neurosci* 2006;7:295-309. DOI PubMed
40. Huot P. Monoamine oxidase A inhibition and Parkinson's disease. *Neurodegener Dis Manag* 2020;10:335-7. DOI PubMed
41. Hamadjida A, Nuara SG, Frouni I, et al. Monoamine oxidase A inhibition as monotherapy reverses parkinsonism in the MPTP-lesioned marmoset. *Naunyn Schmiedebergs Arch Pharmacol* 2020;393:2139-44. DOI PubMed
42. van der Hoek TC, Bus BA, Matui P, van der Marck MA, Esselink RA, Tendolkar I. Prevalence of depression in Parkinson's disease: effects of disease stage, motor subtype and gender. *J Neurol Sci* 2011;310:220-4. DOI
43. Meyer JH, Wilson AA, Sagrati S, et al. Brain monoamine oxidase A binding in major depressive disorder: relationship to selective serotonin reuptake inhibitor treatment, recovery, and recurrence. *Arch Gen Psychiatry* 2009;66:1304-12. DOI PubMed
44. Sacher J, Wilson AA, Houle S, et al. Elevated brain monoamine oxidase A binding in the early postpartum period. *Arch Gen Psychiatry* 2010;67:468-74. DOI PubMed
45. Sacher J, Houle S, Parkes J, et al. Monoamine oxidase A inhibitor occupancy during treatment of major depressive episodes with moclobemide or St. John's wort: an [11C]-harmine PET study. *J Psychiatry Neurosci* 2011;36:375-82. DOI PubMed PMC
46. Mann JJ, Currier D. Medication in suicide prevention insights from neurobiology of suicidal behavior. In: Dwivedi Y, editor. The neurobiological basis of suicide. Boca Raton (FL): CRC Press/Taylor & Francis; 2012. PubMed
47. Schildkraut JJ. The catecholamine hypothesis of affective disorders: a review of supporting evidence. *Am J Psychiatry* 1965;122:509-22. DOI PubMed
48. Delgado PL, Charney DS, Price LH, Aghajanian GK, Landis H, Heninger GR. Serotonin function and the mechanism of antidepressant action. Reversal of antidepressant-induced remission by rapid depletion of plasma tryptophan. *Arch Gen Psychiatry* 1990;47:411-8. DOI PubMed
49. Bymaster FP, McNamara RK, Tran PV. New approaches to developing antidepressants by enhancing monoaminergic neurotransmission. *Expert Opin Investig Drugs* 2003;12:531-43. DOI PubMed
50. Lacombe S, Stanislav SW, Marken PA. Pharmacologic treatment of cocaine abuse. *DICP* 1991;25:818-23. DOI PubMed
51. Miczek KA, Haney M. Psychomotor stimulant effects of d-amphetamine, MDMA and PCP: aggressive and schedule-controlled behavior in mice. *Psychopharmacology (Berl)* 1994;115:358-65. DOI PubMed
52. Vestergaard P, Rejnmark L, Mosekilde L. Fracture risk associated with parkinsonism and anti-Parkinson drugs. *Calcif Tissue Int* 2007;81:153-61. DOI PubMed
53. Krzymowski T, Stefanczyk-Krzymowska S. New facts and the concept of physiological regulation of the dopaminergic system function and its disorders. *J Physiol Pharmacol* 2015;66:331-41. PubMed
54. Liu W, Rabinovich A, Nash Y, et al. Anti-inflammatory and protective effects of MT-031, a novel multitarget MAO-A and AChE/BuChE inhibitor in scopolamine mouse model and inflammatory cells. *Neuropharmacology* 2017;113:445-56. DOI PubMed
55. Monoamine oxidase inhibitors. Meyler's side effects of drugs: the international encyclopedia of adverse drug reactions and interactions. Elsevier; 2006. p. 2371-8. DOI
56. Finberg JP, Tenne M. Relationship between tyramine potentiation and selective inhibition of monoamine oxidase types A and B in the rat vas deferens. *Br J Pharmacol* 1982;77:13-21. DOI PubMed PMC
57. Rudzik AD, Eble JN. The potentiation of pressor responses to tyramine by a number of amphetamine-like compounds. *Proc Soc Exp Biol Med* 1967;124:655-7. DOI PubMed
58. Finberg JP, Lamensdorf I, Weinstock M, Schwartz M, Youdim MB. Pharmacology of rasagiline (N-propargyl-1R-aminoindan). *Adv Neurol* 1999;80:495-9. PubMed
59. Weinstock M, Gorodetsky E, Wang R, Gross A, Weinreb O, Youdim M. Limited potentiation of blood pressure response to oral tyramine by brain-selective monoamine oxidase A-B inhibitor, TV-3326 in conscious rabbits. Supported by Teva Pharmaceuticals Ltd (Israel). *Neuropharmacology* 2002;43:999-1005. DOI
60. Gal S, Abassi ZA, Youdim MB. Limited potentiation of blood pressure in response to oral tyramine by the anti-Parkinson brain selective multifunctional monoamine oxidase-AB inhibitor, M30. *Neurotox Res* 2010;18:143-50. DOI PubMed
61. Bar-Am O, Amit T, Kupersmidt L, et al. Neuroprotective and neurorestorative activities of a novel iron chelator-brain selective monoamine oxidase-A/monoamine oxidase-B inhibitor in animal models of Parkinson's disease and aging. *Neurobiol Aging* 2015;36:1529-42. DOI PubMed
62. Tabet N. Acetylcholinesterase inhibitors for Alzheimer's disease: anti-inflammatories in acetylcholine clothing! *Age Ageing* 2006;35:336-8. DOI PubMed
63. Bohnen NI, Kaufer DI, Ivanco LS, et al. Cortical cholinergic function is more severely affected in parkinsonian dementia than in Alzheimer disease: an in vivo positron emission tomographic study. *Arch Neurol* 2003;60:1745-8. DOI PubMed
64. Hilker R, Thomas AV, Klein JC, et al. Dementia in Parkinson disease: functional imaging of cholinergic and dopaminergic pathways. *Neurology* 2005;65:1716-22. DOI PubMed
65. Kandiah N, Pai MC, Senanarong V, et al. Rivastigmine: the advantages of dual inhibition of acetylcholinesterase and butyrylcholinesterase and its role in subcortical vascular dementia and Parkinson's disease dementia. *Clin Interv Aging* 2017;12:697-707. DOI PubMed PMC
66. Weinstock M. Selectivity of cholinesterase inhibition: clinical implications for the treatment of Alzheimer's disease. *CNS Drugs* 1999;12:307-23. DOI
67. Giacobini E, Spiegel R, Enz A, Veroff AE, Cutler NR. Inhibition of acetyl- and butyryl-cholinesterase in the cerebrospinal fluid of patients with Alzheimer's disease by rivastigmine: correlation with cognitive benefit. *J Neural Transm (Vienna)* 2002;109:1053-65.

[DOI PubMed](#)

68. Mesulam M, Guillozet A, Shaw P, Quinn B. Widely spread butyrylcholinesterase can hydrolyze acetylcholine in the normal and Alzheimer brain. *Neurobiol Dis* 2002;9:88-93. [DOI PubMed](#)
69. Perry EK, Perry RH, Blessed G, Tomlinson BE. Changes in brain cholinesterases in senile dementia of Alzheimer type. *Neuropathol Appl Neurobiol* 1978;4:273-7. [DOI PubMed](#)
70. Mesulam M, Guillozet A, Shaw P, Levey A, Duysen E, Lockridge O. Acetylcholinesterase knockouts establish central cholinergic pathways and can use butyrylcholinesterase to hydrolyze acetylcholine. *Neuroscience* 2002;110:627-39. [DOI PubMed](#)
71. Greig NH, Utsuki T, Yu Q, et al. A new therapeutic target in Alzheimer's disease treatment: attention to butyrylcholinesterase. *Curr Med Res Opin* 2001;17:159-65. [DOI PubMed](#)
72. Nordberg A, Ballard C, Bullock R, Darreh-Shori T, Somogyi M. A review of butyrylcholinesterase as a therapeutic target in the treatment of Alzheimer's disease. *Prim Care Companion CNS Disord* 2013;15:PCC. [DOI PubMed PMC](#)
73. Darreh-Shori T, Almkvist O, Guan ZZ, et al. Sustained cholinesterase inhibition in AD patients receiving rivastigmine for 12 months. *Neurology* 2002;59:563-72. [DOI PubMed](#)
74. Müller T. Rivastigmine in the treatment of patients with Alzheimer's disease. *Neuropsychiatr Dis Treat* 2007;3:211-8. [DOI PubMed PMC](#)
75. Rhee SG. Redox signaling: hydrogen peroxide as intracellular messenger. *Exp Mol Med* 1999;31:53-9. [DOI PubMed](#)
76. Stone JR, Yang S. Hydrogen peroxide: a signaling messenger. *Antioxid Redox Signal* 2006;8:243-70. [DOI PubMed](#)
77. Fridovich I. Superoxide radical and superoxide dismutases. *Annu Rev Biochem* 1995;64:97-112. [DOI PubMed](#)
78. Nicotra A. Monoamine oxidase expression during development and aging. *NeuroToxicology* 2004;25:155-65. [DOI PubMed](#)
79. Riederer P. Monoamine oxidase-B inhibition in Alzheimer's disease. *NeuroToxicology* 2004;25:271-7. [DOI PubMed](#)
80. Youdim MB, Buccafusco JJ. CNS targets for multi-functional drugs in the treatment of Alzheimer's and Parkinson's diseases. *J Neural Transm (Vienna)* 2005;112:519-37. [DOI PubMed](#)
81. Xiao XQ, Lee NT, Carlier PR, Pang Y, Han YF. Bis(7)-tacrine, a promising anti-Alzheimer's agent, reduces hydrogen peroxide-induced injury in rat pheochromocytoma cells: comparison with tacrine. *Neuroscience Letters* 2000;290:197-200. [DOI PubMed](#)
82. Schallreuter KU, Elwary S. Hydrogen peroxide regulates the cholinergic signal in a concentration dependent manner. *Life Sci* 2007;80:2221-6. [DOI PubMed](#)
83. Xiao XQ, Yang JW, Tang XC. Huperzine A protects rat pheochromocytoma cells against hydrogen peroxide-induced injury. *Neurosci Lett* 1999;275:73-6. [DOI PubMed](#)
84. Mortazavian SM, Parsaei H, Mousavi SH, Tayarani-Najarian Z, Ghorbani A, Sadeghnia HR. Acetylcholinesterase inhibitors promote angiogenesis in chick chorioallantoic membrane and inhibit apoptosis of endothelial cells. *Int J Alzheimers Dis* 2013;2013:121068. [DOI PubMed PMC](#)
85. Tyurina YY, Tyurin VA, Carta G, Quinn PJ, Schor NF, Kagan VE. Direct evidence for antioxidant effect of Bcl-2 in PC12 rat pheochromocytoma cells. *Arch Biochem Biophys* 1997;344:413-23. [DOI PubMed](#)
86. Maruyama W, Akao Y, Youdim MB, Davis BA, Naoi M. Transfection-enforced Bcl-2 overexpression and an anti-Parkinson drug, rasagiline, prevent nuclear accumulation of glyceraldehyde-3-phosphate dehydrogenase induced by an endogenous dopaminergic neurotoxin, N-methyl(R)salsolinol. *J Neurochem* 2001;78:727-35. [DOI PubMed](#)
87. Godley BF, Jin GF, Guo YS, Hurst JS. Bcl-2 overexpression increases survival in human retinal pigment epithelial cells exposed to H₂O₂. *Exp Eye Res* 2002;74:663-9. [DOI PubMed](#)
88. Tran VV, Chen G, Newgard CB, Hohmeier HE. Discrete and complementary mechanisms of protection of beta-cells against cytokine-induced and oxidative damage achieved by bcl-2 overexpression and a cytokine selection strategy. *Diabetes* 2003;52:1423-32. [DOI PubMed](#)
89. Jackson GR, Apffel L, Werrbach-Perez K, Perez-Polo JR. Role of nerve growth factor in oxidant-antioxidant balance and neuronal injury. I. Stimulation of hydrogen peroxide resistance. *J Neurosci Res* 1990;25:360-8. [DOI PubMed](#)
90. Mattson MP, Lovell MA, Furukawa K, Markesbery WR. Neurotrophic factors attenuate glutamate-induced accumulation of peroxides, elevation of intracellular Ca²⁺ concentration, and neurotoxicity and increase antioxidant enzyme activities in hippocampal neurons. *J Neurochem* 1995;65:1740-51. [DOI PubMed](#)
91. Hirsch EC, Hunot S, Hartmann A. Neuroinflammatory processes in Parkinson's disease. *Parkinsonism Relat Disord* 2005;11 Suppl 1:S9-S15. [DOI PubMed](#)
92. Clark BD, Collins KL, Gandy MS, Webb AC, Auron PE. Genomic sequence for human prointerleukin 1 beta: possible evolution from a reverse transcribed prointerleukin 1 alpha gene. *Nucleic Acids Res* 1986;14:7897-914. [DOI PubMed PMC](#)
93. Dinarello CA, van der Meer JW. Treating inflammation by blocking interleukin-1 in humans. *Semin Immunol* 2013;25:469-84. [DOI PubMed PMC](#)
94. Zhang YY, Fan YC, Wang M, Wang D, Li XH. Atorvastatin attenuates the production of IL-1 β , IL-6, and TNF- α in the hippocampus of an amyloid β 1-42-induced rat model of Alzheimer's disease. *Clin Interv Aging* 2013;8:103-10. [DOI PubMed PMC](#)
95. Jabbari Azad F, Talaei A, Rafatpanah H, Yousefzadeh H, Jafari R, et al. Association between Cytokine production and disease severity in Alzheimer's disease. *Iran J Allergy Asthma Immunol* 2014;13:433-9. [PubMed](#)
96. Feghali CA, Wright TM. Cytokines in acute and chronic inflammation. *Front Biosci* 1997;2:d12-26. [DOI PubMed](#)
97. Feng LL, Wu XF, Liu HL, et al. Vatacillinol, a resveratrol tetramer, exerts more preferable immunosuppressive activity than its precursor in vitro and in vivo through multiple aspects against activated T lymphocytes. *Toxicol Appl Pharmacol* 2013;267:167-73. [DOI PubMed](#)
98. Henderson Z, Sherriff FE. Distribution of choline acetyltransferase immunoreactive axons and terminals in the rat and ferret

- brainstem. *J Comp Neurol* 1991;314:147-63. DOI PubMed
99. Mahmoodi G, Ahmadi S, Pourmotabbed A, Oryan S, Zarrindast MR. Inhibitory avoidance memory deficit induced by scopolamine: interaction of cholinergic and glutamatergic systems in the ventral tegmental area. *Neurobiol Learn Mem* 2010;94:83-90. DOI PubMed
100. Nielsen JA, Mena E, Williams IH, Nocerini MR, Liston D. Correlation of brain levels of 9-amino-1,2,3,4-tetrahydroacridine (THA) with neurochemical and behavioral changes. *Eur J Pharmacol* 1989;173:53-64. DOI PubMed
101. Bejar C, Wang R, Weinstock M. Effect of rivastigmine on scopolamine-induced memory impairment in rats. *Eur J Pharmacol* 1999;383:231-40. DOI PubMed
102. Weinstock M, Gorodetsky E, Poltyrev T, Gross A, Sagi Y, Youdim M. A novel cholinesterase and brain-selective monoamine oxidase inhibitor for the treatment of dementia comorbid with depression and Parkinson's disease. *Prog Neuropsychopharmacol Biol Psychiatry* 2003;27:555-61. DOI PubMed
103. Buccafusco JJ, Terry AV Jr. Multiple central nervous system targets for eliciting beneficial effects on memory and cognition. *J Pharmacol Exp Ther* 2000;295:438-46. PubMed
104. Bar-Am O, Amit T, Youdim MB, Weinreb O. Neuroprotective and neurorestorative potential of propargylamine derivatives in ageing: focus on mitochondrial targets. *J Neural Transm (Vienna)* 2016;123:125-35. DOI PubMed

Review

Open Access



Review of evidence implicating the plasminogen activator system in blood-brain barrier dysfunction associated with Alzheimer's disease

Mei-Yun Tang¹, Fredric A. Gorin^{1,2}, Pamela J. Lein¹

¹Department of Molecular Biosciences, School of Veterinary Medicine, University of California, Davis, CA 95616, USA.

²Department of Neurology, School of Medicine, University of California, Davis, CA 95616, USA.

Correspondence to: Dr. Pamela J. Lein, Department of Molecular Biosciences, School of Veterinary Medicine, University of California, 1089 Veterinary Medicine Drive, Davis, CA 95616, USA. E-mail: pjlein@ucdavis.edu

How to cite this article: Tang MY, Gorin FA, Lein PJ. Review of evidence implicating the plasminogen activator system in blood-brain barrier dysfunction associated with Alzheimer's disease. *Ageing Neur Dis* 2022;2:2.
<https://dx.doi.org/10.20517/and.2022.05>

Received: 17 Aug 2021 **First Decision:** 9 Nov 2021 **Revised:** 23 Nov 2021 **Accepted:** 17 Jan 2022 **Published:** 29 Jan 2022

Academic Editors: Athanassios P. Kyritsis, Jin-Bin Xu, Weidong Le **Copy Editor:** Xi-Jun Chen **Production Editor:** Xi-Jun Chen

Abstract

Elucidating the pathogenic mechanisms of Alzheimer's disease (AD) to identify therapeutic targets has been the focus of many decades of research. While deposition of extracellular amyloid-beta plaques and intraneuronal neurofibrillary tangles of hyperphosphorylated tau have historically been the two characteristic hallmarks of AD pathology, therapeutic strategies targeting these proteinopathies have not been successful in the clinics. Neuroinflammation has been gaining more attention as a therapeutic target because increasing evidence implicates neuroinflammation as a key factor in the early onset of AD disease progression. The peripheral immune response has emerged as an important contributor to the chronic neuroinflammation associated with AD pathophysiology. In this context, the plasminogen activator system (PAS), also referred to as the vasculature's fibrinolytic system, is emerging as a potential factor in AD pathogenesis. Evolving evidence suggests that the PAS plays a role in linking chronic peripheral inflammatory conditions to neuroinflammation in the brain. While the PAS is better known for its peripheral functions, components of the PAS are expressed in the brain and have been demonstrated to alter neuroinflammation and blood-brain barrier (BBB) permeation. Here, we review plasmin-dependent and -independent mechanisms by which the PAS modulates the BBB in AD pathogenesis and discuss therapeutic implications of these observations.



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, sharing, adaptation, distribution and reproduction in any medium or format, for any purpose, even commercially, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.



Keywords: Bradykinin, matrix metalloproteinase, neuroinflammation, neurovascular unit, plasmin, tissue-type plasminogen activator

INTRODUCTION

Alzheimer's disease (AD) is recognized as the most common cause of dementia in the elderly, and over 6 million Americans are currently living with this disorder. In the United States, AD is the sixth leading single cause of death and the second most common contributing cause of death. The hallmark neuropathologic characteristic of AD is abnormal extracellular protein accumulation in the brain, notably the extracellular deposition of amyloid- β (A β) peptide generated from the improper cleavage of amyloid precursor protein (APP) that gives rise to A β monomers that aggregate into oligomeric A β fibrils and plaques, and intraneuronal neurofibrillary tangles (NF) comprised largely of hyperphosphorylated tau. These proteinopathies are associated with the loss of synapses and subsequent neuronal cell loss in the entorhinal cortex, hippocampus, and frontal cortex^[1-3], and currently, the biomarkers most commonly used in human AD studies are beta-amyloid 42, tau, and phospho-tau proteins in the cerebrospinal fluid. More recently, blood p-tau181 has been reported as being a useful biomarker for distinguishing AD from other dementias^[4]. Thus, it has been widely posited that A β plaques and/or abnormal hyperphosphorylated tau protein accumulation are causally linked to the behavioral and neurologic symptoms of AD. However, therapeutic strategies for decreasing A β plaque load^[5,6], reducing A β production with BACE-1 inhibitors^[7], or inhibiting hyperphosphorylated tau aggregation^[8], have been largely unsuccessful in clinical trials over the past several years^[3]. These failed clinical trials coupled with observations of age-related increases in A β deposition in cognitively intact individuals as well as evidence that A β plaque load does not closely correspond with cognitive decline in AD patients^[1,9] and neurofibrillary tangles are associated with severe cognitive impairment characteristic of late stages of AD^[10,11], have prompted research into alternative pathogenic mechanisms of AD.

It is now recognized that the extracellular deposition of A β and hyperphosphorylated tau triggers pro-inflammatory responses in microglia and astrocytes^[12-14]. The neuroinflammatory response in AD has been described in detail in several recent reviews^[14,15], and it appears that neuroinflammation plays an important role in the early progression of AD^[16,17]. Multiple investigators have shown that A β monofibrils, oligomers, and plaques activate gene expression of pro-inflammatory mediators in microglia and astrocytes^[13,16,18,19]. While microglial phagocytosis of amyloid may be neuroprotective in the early stages of AD by promoting A β clearance^[20,21], microglial activation in later stages may promote the progression of AD^[1,16]. Network-based integrative analysis of whole-genome gene-expression profiling and genotypic data obtained from late-onset AD and non-demented control brains identified the immune/microglia module as the molecular system most strongly associated with the pathophysiology of AD, and in particular, late-onset AD^[22]. Microglial activation is thought to promote AD progression by (1) complement-mediated phagocytosis of synaptic structures to promote synapse loss; and/or (2) release of nitric oxide (NO) and proinflammatory cytokines, including TNF- α , IL-6, and IL-1 β , that act as soluble synaptotoxic factors and induce "A1" neurotoxic astrocytes^[23-26]. In support of these proposed mechanisms, microglial activation has been linked to increased synaptic loss and neurodegeneration in AD^[2,24,27], and pharmacologic inhibition of microglial proliferation in the APP/PS1 mouse effectively shifted microglia to an anti-inflammatory phenotype that was associated with decreased synaptic degeneration and improved memory^[28]. In Alzheimer mouse models, early synaptic loss is associated with C1q complement tightly bound to AB plaques surrounded by neuronal atrophy from microglial phagocytosis^[29]. Mononuclear phagocytes enter the central nervous system (CNS) signaled by chemokines (CXCL1), while the innate immune system also appears to contribute to the neuroinflammatory response to activated microglia in AD models^[30].

While the initial focus on the role of the immune response in AD pathogenesis has been on the brain's intrinsic neuroinflammatory response, attention is now being directed to multiple systemic inflammatory disorders that accelerate or in some instances may be the primary trigger for neuroinflammatory responses that initiate and/or promote AD and other dementias^[31-34]. Some of the observations that have stimulated this shift in focus include reports that young children chronically exposed to high levels of air pollution were found to have neuropathological hallmarks of AD upon incidental autopsy^[35,36], and evidence that type 2 diabetes/ metabolic syndrome and inflammatory bowel disease are associated with increased risk of developing AD^[15,37,38]. The causal factors linking peripheral inflammatory conditions to AD are likely multifactorial and have not yet been clearly delineated; however, several mechanisms are emerging. Peripheral inflammatory conditions have been shown to (1) generate inflammatory cytokines that facilitate access of peripheral inflammatory lymphocytes into the CNS, most notably TNF α , IL-1 β , and IL-6; (2) cause dysfunction of the blood-brain barrier (BBB); and (3) activate the plasminogen activator system (PAS), which has direct effects on the CNS and further facilitates BBB dysfunction. The remainder of this review will investigate the role of the PAS in mediating inflammatory crosstalk between the periphery and the brain and its potential role in AD pathogenesis.

PLASMINOGEN ACTIVATOR SYSTEM

The plasminogen activator system (PAS) is comprised of a group of serine proteases, inhibitors, and binding proteins that control the activity of the serine protease plasmin [Figure 1]^[39]. Plasmin plays a key role in the fibrinolysis cascade, catalyzing the final degradation of fibrin and various extracellular matrix proteins^[40,41]. The zymogen plasminogen (PlG) is converted to activated plasmin by plasmin activators, which include tissue-type plasminogen activator (tPA) and urokinase-type plasminogen activator (uPA). tPA is primarily involved in intravascular fibrinolysis, activating plasminogen that is bound to polymerized fibrin. In contrast, uPA is secreted as a pro-enzyme whose active form is primarily localized on cell surfaces where it binds to the uPA receptor (uPAR). Plasminogen conversion by tPA and uPA in both the periphery and the CNS is tightly regulated by serine protease inhibitors (serpins). Serpins represent a superfamily of proteins with similar structures. Most relevant to this discussion are plasminogen activator inhibitor type 1 (PAI-1) and neuroserpin (NSP). PAI-1 irreversibly inhibits uPA or tPA by undergoing a large conformational change upon binding uPA or tPA that disrupts the active site of the plasmin activator and of PAI-1. In contrast, NSP preferentially inhibits tPA by forming an unstable complex that can release active tPA^[42]. Reflecting the need for stringent regulation of the plasminogen cascade, free forms of activated plasmin activators, PAI-1, and NSP exist at very low concentrations with half lives in the order of minutes^[43,44].

PAS in the periphery

The peripheral PAS plays a central role in mediating fibrinolysis, extracellular migration, cell signaling, cellular migration, and tumor growth, which has been reviewed in detail elsewhere^[45,46]. The PAS converts inactive plasminogen to plasmin, a trypsin-like serine protease, via the catalytic activity of PA^[41]. Plasminogen is primarily present in platelets in the plasma and liver. However, in mice, plasminogen mRNA has been found in the adrenal, kidney, brain, testis, heart, lung, uterus, spleen, thymus, and gut^[40,47]. In the periphery, PAI-1 serves as the main suppressor of plasma fibrinolytic activity^[40]. In the bloodstream, PAI-1 exists on its own in an active form, or as part of a complex with tPA or vitronectin, a glycoprotein that can convert PAI-1 into its active form. Elevated levels of PAI-1 are associated with metabolic syndrome and associated with increased risk of atherothrombosis and stroke^[48,49].

PAS in the CNS

In the CNS, plasminogen is expressed at low levels by neurons in the hippocampus, cortex, cerebellum, as well as neuroendocrine tissues, but it is primarily transported to the brain via systemic circulation^[12,50,51]. Plasminogen has been localized to the extracellular space, while the plasmin activators, tPA and uPA, have




Figure 1. Schematic diagram of the molecular mechanisms of the plasminogen activator system. PAI-1: Plasminogen activator inhibitor-1; NSP: neuroserpin; uPA: urokinase-type plasminogen activator; tPA: tissue-type plasminogen activator; PLG: plasminogen; PLM: plasmin. Created with BioRender.com.

been localized to not only the extracellular space, but also to neurons and astrocytes. Both plasmin activators have been shown to modulate synaptic function when released into the synaptic cleft^[52-54]. Membrane depolarization induces the rapid release of tPA from cerebral cortical neurons, which modulates neuronal plasticity, learning, stress-induced anxiety, and visual cortex plasticity^[55]. tPA and uPA activities have been localized to well-defined areas of the brain^[56-59] and shown to participate in intracellular signaling that is independent of plasminogen activation (see below). tPA is the principal plasmin activator in the CNS with PAI-1 regulating its activity primarily in the extracellular space. NSP is primarily localized in neurons in the developing brain with very low levels detected in the mature CNS^[60], where it preferentially binds to and inhibits tPA^[61]. Interestingly, mutations of NSP are associated with rare familial dementia characterized by neuronal inclusion bodies that are biochemically comprised of polymers of NSP^[62].

Plasmin activity has been shown to be upregulated in axonal growth and synaptic pruning, suggesting a role in brain development and regeneration that is not yet well understood^[50]. While both tPA and uPA can mediate plasminogen activation in the CNS, plasminogen activation is primarily controlled by the tight regulation between tPA and PAI-1^[51]. uPA has a low baseline expression in specific neurons and astrocytes in the normal brain, but is upregulated in pathologically inflammatory environments, such as multiple sclerosis and epilepsy^[50,51]. Endothelial cells of microvessels in the brain contribute to the production of tPA, but tPA can also be expressed by glial cells, neurons, and infiltrating leukocytes, implicating a broad spectrum of tPA involvement in the brain. While tPA in the mature brain is detected primarily in neurons, its enzymatic activity is primarily restricted to the hippocampus, amygdala and hypothalamus^[63,64]. The discrepancy between the expression of tPA mRNA and its areas of enzymatic activity is consistent with its trafficking and transport to mossy fiber tracts^[63,64].

The plasmin activators, tPA and uPA have been shown to play an important role in CNS function and dysfunction with some of their functions being independent of plasminogen^[65,66]. Extracellular tPA participates in cerebellar motor learning^[67], remodeling in various nonneural tissues^[67], and neuronal

regeneration following ischemic injury^[68]. tPA also participates in the regulation of BBB permeability^[69,70]. Neuronal uPA is present in lower levels than tPA, participating in neurogenesis in the developing brain^[71]. Its release in the mature central nervous system triggers astrocytic activation^[53] and, like tPA, uPA promotes axonal and synaptic recovery following different forms of injury^[72]. Both tPA and uPA are found in pre-synaptic vesicles that are released by calcium-dependent mechanisms^[52,54,55].

The PAS is altered in AD

There has been longstanding interest in the role of the PAS in AD beginning with early reports that active plasmin efficiently digests A β peptides^[73-77] both *in vitro* and in rodent AD models^[19,73,74,76-81]. In the AD brain, tPA is highly expressed in regions of AD plaques, and in AD models where tPA is genetically inactivated, there is an increased accumulation of A β , synaptic dysfunction and memory deficits^[78]. However, the enzymatic ability of brain tPA and uPA to activate plasmin *in vivo* is thought to be prevented by irreversible binding to high levels of extracellular PAI-1 secreted by immune-activated microglia and astrocytes^[18]. PAI-1 is minimally expressed in the normal brain or cerebral vasculature, but does increase with senescence^[82-84]. Brain levels of PAI-1 are also markedly increased in APP/PS1 mice^[66] and the serum levels of PAI-1 are positively correlated with cognitive impairment in AD patients^[85]. Consistent with the hypothesis that PAI-1 promotes AD pathology, genetic knockdown or small molecule inhibitors of PAI-1 reduced plaque formation in AD rodent models, and the small molecule PAI-1 inhibitor, PAZ-417, was shown to significantly improve hippocampal LTP and cognitive function in AD mice^[73,74,86,87]. This finding was confirmed recently in an APP/PS1 AD mouse model using another small molecule PAI-1 inhibitor^[86].

Whether tPA primarily plays a beneficial or detrimental role in AD progression is debated. Several studies have demonstrated that tPA activation of plasmin enzymatically reduces A β accumulation^[78]. Conversely, tPA has been shown to mediate excitotoxic neurodegeneration by activating plasmin and causing subsequent laminin degradation^[66,78]. Independent of plasmin activation, tPA causes GSK3 activation, tau hyperphosphorylation, microtubule destabilization, and neurotoxicity in rodent hippocampal neurons^[88]. It has also been shown to mediate amyloid-induced microglial activation^[89]. Based on such observations, it has been proposed that tPA contributes to neurotoxicity, microglial activation, and tau hyperphosphorylation as part of a feed-forward inflammatory pathway^[73,88,89].

PAI-1 expression has been reported to be increased in the plasma^[85,90,91] and brain tissues of AD patients^[76]. PAI-1 expression is not detected in normal healthy human brains but is sporadically present in aged brains^[84,92], and possibly linked to cerebrovascular disease. PAI-1 is the primary regulator of tPA in the CNS extracellular space and is a proinflammatory biomarker. Cytokines upregulate PAI-1 expression in microglia and astrocytes in human and animal models of AD^[18,93]. The PAI-1 promoter is activated by TNF- α via an NF κ B 5' upstream element and directly activated by TGF- β 1 via SMAD2/3 promoter binding sites^[82,94,95]. When PAI-1 is complexed with low density lipoprotein receptor-related protein-1 (LRP-1), it signals changes in microglial morphology and motility that are consistent with microglial activation^[96-98]. In patients with AD, plasminogen activator activity is reduced while PAI-1 and NSP are upregulated^[99]. However, there are contradictory findings regarding measurements of PAI-1 and tPA in the CSF and serum of patients with AD^[76,92,100].

Congophilic amyloid angiopathy (CAA) is a vascular complication of AD in which A β 40 plaques accumulate within the brain endothelium of cerebral arteries, arterioles and capillaries^[101]. CAA can result in intracranial hemorrhages, cognitive impairment, or subacute inflammatory encephalopathy. tPA activation of endothelial NMDA receptors has been shown to regulate neurovascular coupling via nitric oxide-mediated regulation of cerebral blood flow. Elevated levels of brain PAI-1 impairs this tPA-dependent

neurovascular coupling in Tg2576 AD mice, and pharmacologic inhibition of PAI-1 was shown to improve cognition in this animal model by selectively restoring neurovascular function while not affecting cortical amyloid plaques^[102].

PAS modulates BBB integrity in AD

There is increasing evidence identifying BBB leakage as an early sign of cognitive dysfunction, as well as evidence linking BBB dysfunction to AD pathogenesis^[103,104] and its neuroinflammatory pathology^[33,105]. However, the mechanisms underlying BBB dysfunction in AD are currently not well-elucidated. The BBB is part of the neurovascular unit (NVU) in the brain, which consists of endothelial cells (ECs), mural cells, including vascular smooth muscle cells and pericytes, basement membrane, glia cells including astrocytes and microglia, and neurons [Figure 2]. The ECs of the BBB are a distinct characteristic of the NVU due to their tight junctions and lack of fenestrae. This allows the ECs to regulate the selective transport and metabolism of substances from blood to brain and vice versa, thereby separating the microenvironment of the brain parenchyma from changes in circulating ion and metabolite concentrations in the systemic circulation^[105].

In CNS injury, there are several potential mechanisms by which tPA is able to mediate changes in the permeability of the BBB [Figure 3], which in turn further exacerbates CNS injury by promoting neuroinflammation. AD is associated with BBB dysfunction in humans and animal models. Amyloid deposition activates gliosis that can alter the morphology of astrocytic endfeet, which are integral to the integrity of the neurovascular unit. As described previously with CAA, amyloid deposition can also injure the brain endothelium, which can additionally impair BBB integrity^[106]. Finally, A β oligomers stimulate fibrin production that complexes with amyloid plaques, and fibrin has been shown to be increased in the parenchyma and vasculature of AD brains^[107,108]. This fibrin-A β complex promotes further neuroinflammation and neurodegeneration. tPA is conformationally activated by fibrin deposition, but its enzymatic activity is inhibited by the elevated levels of PAI-1 found in AD parenchyma. However, as summarized in Figure 3, activated tPA has multiple plasmin-independent mechanisms by which it can compromise BBB integrity.

tPA in the CNS directly alters BBB integrity

tPA has long been known to play a significant role in the NVU, mostly in the context of stroke^[109-111]. tPA has been reported to directly alter the BBB integrity by triggering activation of LRP-1 on the surface of astrocytes^[12]. LRP-1 is a multifunctional signaling receptor that functions in receptor-mediated endocytosis and cellular signaling. LRP-1 binds many ligands, including tPA and amyloid-beta^[112], which thereby facilitates A β endocytosis across endothelial cells of the BBB^[113]. A β oligomers may compromise BBB integrity via activation of matrix metalloproteinases (MMPs)^[113]. Alternatively, tPA may cleave LRP-1 at its substrate binding ectodomain, activating NF- κ B, which promotes the synthesis of matrix metalloproteinases MMP-3 and MMP-9, leading to matrix protein degradation and BBB leakage^[12]. tPA-induced activation of LRP-1 shedding from astrocytic endfeet also promotes detachment of endfeet projections from tight junctions of the endothelial cells of the neurovascular unit, further compromising the BBB^[12]. Additionally, tPA can directly alter BBB integrity via platelet-derived growth factor PDGF-CC^[114]. Upregulated neuronal expression of tPA expression induced by CNS disease or injury results in the release of tPA into the extracellular matrix of the brain, where it cleaves complement subcomponents C1r/C1s, urchin EGF-like protein, and bone-morphogenic protein-1 (CUB) from PDGF-CC forming an active ligand that binds to PDGF receptor- α (PDGFR- α). PDGFR- α promotes BBB leakage that worsens cerebral edema, neuroinflammation and neuronal death^[114]. One study found this tPA-mediated activation of PDGF-CC to be inefficient in an *in vitro* stroke model^[115]. However, *in vivo*, the Mac-1 integrin expressed on microglia works cooperatively with the endocytic receptor LRP-1 to promote tPA-mediated activation of PDGF-




Figure 2. Cross-section of the neurovascular unit (NVU) in a normal brain vs. an Alzheimer's disease (AD) brain. The blood-brain barrier (BBB) consists of endothelial cells joined by tight junctions, basement membrane, mural cells (i.e., pericytes and vascular smooth muscle cells), enclosed by astrocytic endfeet. Neurons and microglia closely associate with the BBB. In the AD brain, the NVU undergoes morphological and structural changes due to AD pathology. Amyloid-beta plaques complexed to fibrin result in neuroinflammation and BBB disruption, including activated microglia, swollen astrocytic endfeet, and compromised tight junctions. Created with BioRender.com.

CC^[115]. Multiple studies have also implicated tPA in binding amyloid-beta, thereby facilitating A β endocytosis across endothelial cells of the BBB^[113].

Peripheral tPA alters BBB

In addition to its endogenous effects within the CNS, peripheral tPA can cross the intact BBB^[116], phosphorylate claudin-5 and occludin, thereby weakening endothelial tight junctions and increasing BBB permeability by plasmin-independent mechanisms^[117,118]. Chronic release of plasma tPA can induce a hyperfibrinolytic state that also directly increases vascular permeability of the BBB. Resultant plasmin activation by tPA also triggers bradykinin (BK) production^[119,120]. BK is a peptide mediator generated from its circulating precursor, high molecular weight kininogen (HMWK), and is known for its ability to induce vascular permeability and cause vasodilation of arteries and veins^[119]. It is a pro-inflammatory mediator, and its role as a neuromediator was identified in clinical conditions including AD^[119]. While it is still debated as to how the PAS triggers BK generation, two primary pathways have been proposed [Figure 3]. A direct mechanism identified using an *in vitro* model involves tPA-mediated conversion of plasminogen to plasmin, which then cleaves HMWK into BK. BK acts through the bradykinin 2 receptor (B2R) on endothelial cells, triggering a signaling cascade that promotes intracellular calcium release and downregulation of claudin-5, a critical protein involved in maintaining EC tight junctions^[120]. B2R activation can additionally induce tPA release from endothelial cells, further amplifying additional BK generation^[121]. The PAS alternatively can indirectly trigger BK formation through a plasmin-dependent pathway where plasmin activated by tPA then converts Factor XII (FXII) into Factor XIIa (FXIIa), which then converts plasma pre-kallikrein into plasma kallikrein (PKal)^[121]. PKal then serves to cleave HMWK, leading to BK formation and B2R signaling activation [Figure 3]. This indirect mechanism was demonstrated *ex vivo* and *in vivo* with the former using human plasma incubated with tPA, which resulted in the formation of active PKal; the latter demonstrating that intravenous injection of tPA in mice increased PKal activity^[121,122].




Figure 3. Mechanisms by which tPA may disrupt the blood-brain barrier. (1) tissue-type plasminogen activator (tPA) released from neurons cleaves lipoprotein receptor-related protein-1 (LRP-1) to activate an NF- κ B signaling cascade resulting in the production of MMP-9. tPA and LRP-1 can bind amyloid beta, which facilitates A β endocytosis across the blood-brain barrier (BBB). (2) Neuronal tPA degrades platelet-derived growth factor-CC (PDGF-CC) to release the active ligand for PDGF receptor- α (PDGFR- α) on astrocytic endfeet, causing them to retract from endothelial cells. (3) Plasma tPA activates plasmin to directly produce bradykinin that activates bradykinin 2 receptor (B2R) receptor on endothelial cells. (4) Plasma tPA cleaves plasminogen to generate plasmin that indirectly upregulates bradykinin expression through plasma kallikrein (PKal). Created with BioRender.com.

AD has been shown to produce BBB dysfunction in humans and animal models. Amyloid deposition activates gliosis that can alter the morphology of astrocytic endfeet, which are integral to the integrity of the neurovascular unit. As described previously with CAA, amyloid deposition can injure the brain endothelium, which can additionally impair BBB integrity^[106]. Finally, A β oligomers stimulate fibrin production that complexes with amyloid plaques and has been shown to be increased in the parenchyma and vasculature of AD brains^[107]. This fibrin-A β complex promotes further neuroinflammation and neurodegeneration. tPA is conformationally activated by fibrin deposition, but its enzymatic activity is inhibited by the elevated levels of PAI-1 found in AD parenchyma. However, as summarized in Figure 3, activated tPA has multiple plasmin-independent mechanisms by which it can compromise BBB integrity.

CONCLUSION

Over the past two decades following initial reports of histologic evidence of A β deposition in the brains of children chronically exposed to severe air pollution^[123], it has become clear that chronic peripheral inflammatory conditions, including those that involve lung, gut, liver, and metabolic syndrome, exacerbate or initiate neuroinflammatory disorders. This has been supported by epidemiologic findings of a positive association between chronic peripheral inflammatory conditions and increased incidence of dementia, including AD. More recently, there has been increased interest in the contribution of the peripheral PAS to the neuroinflammatory component of AD. Recently, it has become recognized that the risk of blood clots,

increased mortality, and persistent neuroinflammatory complications of COVID 19 viral infections are also associated with pre-existing systemic inflammatory disorders shown to chronically activate components of the PAS^[124]. With respect to AD, the available evidence suggests that the peripheral PAS may modulate the neuroinflammatory response via multiple mechanisms^[12,51]. Besides fostering the transcytosis of inflammatory cells across the BBB, components of the PAS have been shown to decrease BBB integrity and increase BBB permeability, consequences that have been independently linked to early cognitive dysfunction^[125] including progressive stages of AD^[126] perhaps in association with concomitant vascular disease^[127]. Overall, the means by which the PAS modulates BBB integrity by tPA and plasmin-dependent mechanisms is complex and requires further validation and investigation. tPA in the CNS has been shown to alter BBB permeability by LRP-1 and PDGF-CC-dependent mechanisms, while tPA produced from peripheral inflammation can cross the BBB where it may work in tandem with the kinin system to directly generate BK via plasmin, or indirectly by increased PKal. It is likely that tPA works multifactorially and that these mechanisms are not mutually exclusive [Figure 2]^[118]. Based on what is currently known, further studies investigating the role of the PAS in AD and other dementias are certainly warranted.

DECLARATIONS

Acknowledgments

The authors gratefully acknowledge Dr. Suzette Smiley-Jewell for assistance with manuscript preparation.

Authors' contributions

Made substantial contributions to conception and outline of the review: Tang MY, Gorin FA, Lein PJ

Performed literature search to identify relevant publications: Tang MY, Gorin FA

Wrote the initial draft of the manuscript and created figures: Tang MY

Significantly edited early versions of the manuscript: Gorin FA

Made final edits to the manuscript: Lein PJ

Obtained funding to support the work: Gorin FA, Lein PJ

Availability of data and materials

Not applicable.

Financial support and sponsorship

This work was supported by the National Institute of Aging (Grant No. P30 AG010129; No. R21 AG065908). The National Institute of Aging (NIA) was not involved in the design, layout, writing or decision to submit this review for publication. The contents of this work do not necessarily represent the official views of the NIA, and the NIA does not endorse the purchase of any commercial products or services mentioned in this publication.

Conflicts of interest

All authors declared that there are no conflicts of interest.

Ethical approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Copyright

© The Author(s) 2022.

REFERENCES

1. Morris GP, Clark IA, Vissel B. Inconsistencies and controversies surrounding the amyloid hypothesis of Alzheimer's disease. *Acta Neuropathol Commun* 2014;2:135. DOI PubMed PMC
2. Herrup K. Reimagining Alzheimer's disease--an age-based hypothesis. *J Neurosci* 2010;30:16755-62. DOI PubMed PMC
3. Makin S. The amyloid hypothesis on trial. *Nature* 2018;559:S4-7. DOI PubMed
4. Janelidze S, Mattsson N, Palmqvist S, et al. Plasma P-tau181 in Alzheimer's disease: relationship to other biomarkers, differential diagnosis, neuropathology and longitudinal progression to Alzheimer's dementia. *Nat Med* 2020;26:379-86. DOI PubMed
5. Honig LS, Vellas B, Woodward M, et al. Trial of Solanezumab for mild dementia due to Alzheimer's disease. *N Engl J Med* 2018;378:321-30. DOI PubMed
6. Decourt B, Boumelhem F, Pope ED 3rd, Shi J, Mari Z, Sabbagh MN. Critical appraisal of amyloid lowering agents in AD. *Curr Neurol Neurosci Rep* 2021;21:39. DOI PubMed PMC
7. Egan MF, Kost J, Tariot PN, et al. Randomized trial of verubecestat for mild-to-moderate Alzheimer's disease. *N Engl J Med* 2018;378:1691-703. DOI PubMed PMC
8. Gauthier S, Feldman HH, Schneider LS, et al. Efficacy and safety of tau-aggregation inhibitor therapy in patients with mild or moderate Alzheimer's disease: a randomised, controlled, double-blind, parallel-arm, phase 3 trial. *Lancet* 2016;388:2873-84. DOI PubMed PMC
9. Strooper B, Karran E. The cellular phase of Alzheimer's disease. *Cell* 2016;164:603-15. DOI PubMed
10. Huang Y, Mucke L. Alzheimer mechanisms and therapeutic strategies. *Cell* 2012;148:1204-22. DOI PubMed PMC
11. Nelson PT, Alafuzoff I, Bigio EH, et al. Correlation of Alzheimer disease neuropathologic changes with cognitive status: a review of the literature. *J Neuropathol Exp Neurol* 2012;71:362-81. DOI PubMed PMC
12. Mehra A, Ali C, Parcq J, Vivien D, Docagne F. The plasminogen activation system in neuroinflammation. *Biochim Biophys Acta* 2016;1862:395-402. DOI PubMed
13. Akiyama H, Barger S, Barnum S, et al. Inflammation and Alzheimer's disease. *Neurobiol Aging* 2000;21:383-421. DOI PubMed PMC
14. Jevtic S, Sengar AS, Salter MW, McLaurin J. The role of the immune system in Alzheimer disease: etiology and treatment. *Ageing Res Rev* 2017;40:84-94. DOI PubMed
15. Kinney JW, Bemiller SM, Murtishaw AS, Leisgang AM, Salazar AM, Lamb BT. Inflammation as a central mechanism in Alzheimer's disease. *Alzheimers Dement (N Y)* 2018;4:575-90. DOI PubMed PMC
16. Heneka MT, Carson MJ, Khoury JE, et al. Neuroinflammation in Alzheimer's disease. *Lancet Neurol* 2015;14:388-405. DOI PubMed PMC
17. Ni J, Wu Z. Inflammation spreading: negative spiral linking systemic inflammatory disorders and Alzheimer's disease. *Front Cell Neurosci* 2021;15:638686. DOI PubMed PMC
18. Oh S, Suh N, Kim I, Lee JY. Impacts of aging and amyloid- β deposition on plasminogen activators and plasminogen activator inhibitor-1 in the Tg2576 mouse model of Alzheimer's disease. *Brain Res* 2015;1597:159-67. DOI PubMed
19. Lee JY, Kweon HS, Cho E, et al. Upregulation of tPA/plasminogen proteolytic system in the periphery of amyloid deposits in the Tg2576 mouse model of Alzheimer's disease. *Neurosci Lett* 2007;423:82-7. DOI PubMed
20. Ries M, Sastre M. Mechanisms of A β clearance and degradation by glial cells. *Front Aging Neurosci* 2016;8:160. DOI PubMed PMC
21. Tarasoff-Conway JM, Carare RO, Osorio RS, et al. Clearance systems in the brain-implications for Alzheimer disease. *Nat Rev Neurol* 2015;11:457-70. DOI PubMed PMC
22. Gerrits E, Brouwer N, Kooistra SM, et al. Distinct amyloid- β and tau-associated microglia profiles in Alzheimer's disease. *Acta Neuropathol* 2021;141:681-96. DOI PubMed PMC
23. Hong S, Dissing-Olesen L, Stevens B. New insights on the role of microglia in synaptic pruning in health and disease. *Curr Opin Neurobiol* 2016;36:128-34. DOI PubMed PMC
24. Rajendran L, Paolicelli RC. Microglia-mediated synapse loss in Alzheimer's disease. *J Neurosci* 2018;38:2911-9. DOI PubMed PMC
25. Stephan AH, Barres BA, Stevens B. The complement system: an unexpected role in synaptic pruning during development and disease. *Annu Rev Neurosci* 2012;35:369-89. DOI PubMed
26. Liddelow SA, Guttenplan KA, Clarke LE, et al. Neurotoxic reactive astrocytes are induced by activated microglia. *Nature* 2017;541:481-7. DOI PubMed PMC
27. Hansen DV, Hanson JE, Sheng M. Microglia in Alzheimer's disease. *J Cell Biol* 2018;217:459-72. DOI PubMed PMC
28. Olmos-Alonso A, Schettters ST, Sri S, et al. Pharmacological targeting of CSF1R inhibits microglial proliferation and prevents the progression of Alzheimer's-like pathology. *Brain* 2016;139:891-907. DOI PubMed PMC
29. Hong S, Beja-Glasser VF, Nfonoyim BM, et al. Complement and microglia mediate early synapse loss in Alzheimer mouse models. *Science* 2016;352:712-6. DOI PubMed PMC
30. Cisbani G, Rivest S. Targeting innate immunity to protect and cure Alzheimer's disease: opportunities and pitfalls. *Mol Psychiatry* 2021;26:5504-15. DOI PubMed

31. Heissig B, Salama Y, Takahashi S, Osada T, Hattori K. The multifaceted role of plasminogen in inflammation. *Cell Signal* 2020;75:109761. DOI PubMed PMC
32. Newcombe EA, Camats-Perna J, Silva ML, Valmas N, Huat TJ, Medeiros R. Inflammation: the link between comorbidities, genetics, and Alzheimer's disease. *J Neuroinflammation* 2018;15:276. DOI PubMed PMC
33. Le Page A, Dupuis G, Frost EH, et al. Role of the peripheral innate immune system in the development of Alzheimer's disease. *Exp Gerontol* 2018;107:59-66. DOI PubMed
34. Svenningsen P, Hinrichs GR, Zachar R, Ydegaard R, Jensen BL. Physiology and pathophysiology of the plasminogen system in the kidney. *Pflugers Arch* 2017;469:1415-23. DOI PubMed
35. Calderón-Garcidueñas L, Franco-Lira M, Mora-Tiscareño A, Medina-Cortina H, Torres-Jardón R, Kavanaugh M. Early Alzheimer's and Parkinson's disease pathology in urban children: Friend versus foe responses--it is time to face the evidence. *Biomed Res Int* 2013;2013:161687. DOI PubMed PMC
36. Calderón-Garcidueñas L, Kavanaugh M, Block M, et al. Neuroinflammation, hyperphosphorylated tau, diffuse amyloid plaques, and down-regulation of the cellular prion protein in air pollution exposed children and young adults. *J Alzheimers Dis* 2012;28:93-107. DOI PubMed
37. Jialal I, Devaraj S. Subcutaneous adipose tissue biology in metabolic syndrome. *Horm Mol Biol Clin Investig* 2018;33. DOI PubMed
38. Zhang B, Wang HE, Bai YM, et al. Inflammatory bowel disease is associated with higher dementia risk: a nationwide longitudinal study. *Gut* 2021;70:85-91. DOI PubMed
39. Chana-Muñoz A, Jendroszek A, Sønnichsen M, et al. Origin and diversification of the plasminogen activation system among chordates. *BMC Evol Biol* 2019;19:27. DOI PubMed PMC
40. Cesari M, Pahor M, Incalzi RA. Plasminogen activator inhibitor-1 (PAI-1): a key factor linking fibrinolysis and age-related subclinical and clinical conditions. *Cardiovasc Ther* 2010;28:e72-91. DOI PubMed PMC
41. Kruihof EK. Regulation of plasminogen activator inhibitor type 1 gene expression by inflammatory mediators and statins. *Thromb Haemost* 2008;100:969-75. PubMed
42. Lee TW, Tsang VW, Birch NP. Physiological and pathological roles of tissue plasminogen activator and its inhibitor neuroserpin in the nervous system. *Front Cell Neurosci* 2015;9:396. DOI PubMed PMC
43. Ricagno S, Caccia S, Sorrentino G, Antonini G, Bolognesi M. Human neuroserpin: structure and time-dependent inhibition. *J Mol Biol* 2009;388:109-21. DOI PubMed
44. Chandler WL, Alessi MC, Aillaud MF, Henderson P, Vague P, Juhan-Vague I. Clearance of tissue plasminogen activator (TPA) and TPA/plasminogen activator inhibitor type 1 (PAI-1) complex: relationship to elevated TPA antigen in patients with high PAI-1 activity levels. *Circulation* 1997;96:761-8. DOI PubMed
45. Castellino FJ, Ploplis VA. Structure and function of the plasminogen/plasmin system. *Thromb Haemost* 2005;93:647-54. DOI PubMed
46. Andreasen PA, Egelund R, Petersen HH. The plasminogen activation system in tumor growth, invasion, and metastasis. *Cell Mol Life Sci* 2000;57:25-40. DOI PubMed
47. Zhang L, Seiffert D, Fowler BJ, Jenkins GR, Thinnies TC, et al. Plasminogen has a broad extrahepatic distribution. *Thromb Haemost* 2002;87:493-501. PubMed
48. Samad F, Ruf W. Inflammation, obesity, and thrombosis. *Blood* 2013;122:3415-22. DOI PubMed PMC
49. Tjärnlund-Wolf A, Brogren H, Lo EH, Wang X. Plasminogen activator inhibitor-1 and thrombotic cerebrovascular diseases. *Stroke* 2012;43:2833-9. DOI PubMed PMC
50. Atsev S, Tomov N. Using antifibrinolytics to tackle neuroinflammation. *Neural Regen Res* 2020;15:2203-6. DOI PubMed PMC
51. Baker SK, Strickland S. A critical role for plasminogen in inflammation. *J Exp Med* 2020;217:e20191865. DOI PubMed PMC
52. Diaz A, Merino P, Guo JD, et al. Urokinase-type plasminogen activator protects cerebral cortical neurons from soluble A β -induced synaptic damage. *J Neurosci* 2020;40:4251-63. DOI PubMed PMC
53. Diaz A, Merino P, Manrique LG, et al. A cross talk between neuronal urokinase-type plasminogen activator (uPA) and astrocytic uPA receptor (uPAR) promotes astrocytic activation and synaptic recovery in the ischemic brain. *J Neurosci* 2017;37:10310-22. DOI PubMed PMC
54. Jeanneret V, Ospina JP, Diaz A, et al. Tissue-type plasminogen activator protects the postsynaptic density in the ischemic brain. *J Cereb Blood Flow Metab* 2018;38:1896-910. DOI PubMed PMC
55. Yepes M. The plasminogen activating system in the pathogenesis of Alzheimer's disease. *Neural Regen Res* 2021;16:1973-7. DOI PubMed PMC
56. Chevillet A, Lesept F, Lenoir S, Ali C, Parcq J, Vivien D. Impacts of tissue-type plasminogen activator (tPA) on neuronal survival. *Front Cell Neurosci* 2015;9:415. DOI PubMed PMC
57. Lenoir S, Varangot A, Lebouvier L, Galli T, Hommet Y, Vivien D. Post-synaptic release of the neuronal tissue-type plasminogen activator (tPA). *Front Cell Neurosci* 2019;13:164. DOI PubMed PMC
58. Sappino AP, Madani R, Huarte J, et al. Extracellular proteolysis in the adult murine brain. *J Clin Invest* 1993;92:679-85. DOI PubMed PMC
59. Vassalli J, Belin D. Amiloride selectively inhibits the urokinase-type plasminogen activator. *FEBS Letters* 1987;214:187-91. DOI PubMed
60. Yepes M, Lawrence DA. Neuroserpin: a selective inhibitor of tissue-type plasminogen activator in the central nervous system. *Thromb Haemost* 2004;91:457-64. DOI PubMed
61. Lee TW, Tsang VW, Loeff EJ, Birch NP. Physiological and pathological functions of neuroserpin: regulation of cellular responses

- through multiple mechanisms. *Semin Cell Dev Biol* 2017;62:152-9. DOI PubMed
62. Davis RL, Shrimpton AE, Holohan PD, et al. Familial dementia caused by polymerization of mutant neuroserpin. *Nature* 1999;401:376-9. DOI PubMed
 63. Salles FJ, Strickland S. Localization and regulation of the tissue plasminogen activator-plasmin system in the hippocampus. *J Neurosci* 2002;22:2125-34. PubMed PMC
 64. Stevenson TK, Lawrence DA. Characterization of tissue plasminogen activator expression and trafficking in the adult murine brain. *eNeuro* 2018;5:ENEURO. DOI PubMed PMC
 65. Zhu J, Wan Y, Xu H, Wu Y, Hu B, Jin H. The role of endogenous tissue-type plasminogen activator in neuronal survival after ischemic stroke: friend or foe? *Cell Mol Life Sci* 2019;76:1489-506. DOI PubMed
 66. Melchor JP, Strickland S. Tissue plasminogen activator in central nervous system physiology and pathology. *Thromb Haemost* 2005;93:655-60. DOI PubMed PMC
 67. Seeds NW, Basham ME, Ferguson JE. Absence of tissue plasminogen activator gene or activity impairs mouse cerebellar motor learning. *J Neurosci* 2003;23:7368-75. PubMed PMC
 68. Yepes M. The plasminogen activation system promotes neurorepair in the ischemic brain. *Curr Drug Targets* 2019;20:953-9. DOI PubMed PMC
 69. Yepes M. Tissue-type plasminogen activator is a neuroprotectant in the central nervous system. *Front Cell Neurosci* 2015;9:304. DOI PubMed PMC
 70. Yepes M, Sandkvist M, Moore EG, Bugge TH, Strickland DK, Lawrence DA. Tissue-type plasminogen activator induces opening of the blood-brain barrier via the LDL receptor-related protein. *J Clin Invest* 2003;112:1533-40. DOI PubMed PMC
 71. Dent MA, Sumi Y, Morris RJ, Seeley PJ. Urokinase-type plasminogen activator expression by neurons and oligodendrocytes during process outgrowth in developing rat brain. *Eur J Neurosci* 1993;5:633-47. DOI PubMed
 72. Merino P, Diaz A, Jeanneret V, et al. Urokinase-type plasminogen activator (uPA) binding to the uPA receptor (uPAR) promotes axonal regeneration in the central nervous system. *J Biol Chem* 2017;292:2741-53. DOI PubMed PMC
 73. Jacobsen JS, Comery TA, Martone RL, et al. Enhanced clearance of Abeta in brain by sustaining the plasmin proteolysis cascade. *Proc Natl Acad Sci U S A* 2008;105:8754-9. DOI PubMed PMC
 74. Kutz SM, Higgins CE, Higgins PJ. Novel combinatorial therapeutic targeting of PAI-1 (SERPINE1) gene expression in Alzheimer's disease. *Mol Med Ther* 2012;1:106. DOI PubMed PMC
 75. Ledesma MD, Da Silva JS, Crassaerts K, Delacourte A, De Strooper B, Dotti CG. Brain plasmin enhances APP alpha-cleavage and Abeta degradation and is reduced in Alzheimer's disease brains. *EMBO Rep* 2000;1:530-5. DOI PubMed PMC
 76. Liu RM, van Groen T, Katre A, et al. Knockout of plasminogen activator inhibitor 1 gene reduces amyloid beta peptide burden in a mouse model of Alzheimer's disease. *Neurobiol Aging* 2011;32:1079-89. DOI PubMed PMC
 77. Tucker HM, Kihiko M, Caldwell JN, et al. The plasmin system is induced by and degrades amyloid-beta aggregates. *J Neurosci* 2000;20:3937-46. PubMed PMC
 78. Oh SB, Byun CJ, Yun JH, et al. Tissue plasminogen activator arrests Alzheimer's disease pathogenesis. *Neurobiol Aging* 2014;35:511-9. DOI PubMed
 79. Melchor JP, Pawlak R, Strickland S. The tissue plasminogen activator-plasminogen proteolytic cascade accelerates amyloid-beta (Abeta) degradation and inhibits Abeta-induced neurodegeneration. *J Neurosci* 2003;23:8867-71. PubMed PMC
 80. Tucker HM, Kihiko-Ehmann M, Estus S. Urokinase-type plasminogen activator inhibits amyloid-beta neurotoxicity and fibrillogenesis via plasminogen. *J Neurosci Res* 2002;70:249-55. DOI PubMed
 81. Tucker HM, Kihiko-Ehmann M, Wright S, Rydel RE, Estus S. Tissue plasminogen activator requires plasminogen to modulate amyloid-beta neurotoxicity and deposition. *J Neurochem* 2000;75:2172-7. DOI PubMed
 82. Samarakoon R, Higgins SP, Higgins CE, Higgins PJ. The TGF- β 1/p53/PAI-1 signaling axis in vascular senescence: role of caveolin-1. *Biomolecules* 2019;9:341. DOI PubMed PMC
 83. Vaughan DE, Rai R, Khan SS, Eren M, Ghosh AK. Plasminogen activator inhibitor-1 is a marker and a mediator of senescence. *Arterioscler Thromb Vasc Biol* 2017;37:1446-52. DOI PubMed PMC
 84. Yamamoto K, Takeshita K, Kojima T, Takamatsu J, Saito H. Aging and plasminogen activator inhibitor-1 (PAI-1) regulation: implication in the pathogenesis of thrombotic disorders in the elderly. *Cardiovasc Res* 2005;66:276-85. DOI PubMed
 85. Oh J, Lee HJ, Song JH, Park SI, Kim H. Plasminogen activator inhibitor-1 as an early potential diagnostic marker for Alzheimer's disease. *Exp Gerontol* 2014;60:87-91. DOI PubMed
 86. Akhter H, Huang WT, van Groen T, Kuo HC, Miyata T, Liu RM. A small molecule inhibitor of plasminogen activator inhibitor-1 reduces brain amyloid- β load and improves memory in an animal model of Alzheimer's disease. *J Alzheimers Dis* 2018;64:447-57. DOI PubMed
 87. Hohensinner PJ, Baumgartner J, Kral-Pointner JB, et al. PAI-1 (plasminogen activator inhibitor-1) expression renders alternatively activated human macrophages proteolytically quiescent. *Arterioscler Thromb Vasc Biol* 2017;37:1913-22. DOI PubMed PMC
 88. Medina MG, Ledesma MD, Domínguez JE, et al. Tissue plasminogen activator mediates amyloid-induced neurotoxicity via Erk1/2 activation. *EMBO J* 2005;24:1706-16. DOI PubMed PMC
 89. Pineda D, Ampurdanés C, Medina MG, et al. Tissue plasminogen activator induces microglial inflammation via a noncatalytic molecular mechanism involving activation of mitogen-activated protein kinases and Akt signaling pathways and AnnexinA2 and Galectin-1 receptors. *Glia* 2012;60:526-40. DOI PubMed
 90. Hou B, Eren M, Painter CA, et al. Tumor necrosis factor alpha activates the human plasminogen activator inhibitor-1 gene through a distal nuclear factor kappaB site. *J Biol Chem* 2004;279:18127-36. DOI PubMed

91. Loures CMG, Duarte RCF, Silva MVF, et al. Hemostatic abnormalities in dementia: a systematic review and meta-analysis. *Semin Thromb Hemost* 2019;45:514-22. DOI PubMed
92. Sutton R, Keohane ME, VanderBerg SR, Gonias SL. Plasminogen activator inhibitor-1 in the cerebrospinal fluid as an index of neurological disease. *Blood Coagul Fibrinolysis* 1994;5:167-71. DOI PubMed
93. Barker R, Kehoe PG, Love S. Activators and inhibitors of the plasminogen system in Alzheimer's disease. *J Cell Mol Med* 2012;16:865-76. DOI PubMed PMC
94. Higgins PJ. The TGF-beta1/upstream stimulatory factor-regulated PAI-1 gene: potential involvement and a therapeutic target in Alzheimer's disease. *J Biomed Biotechnol* 2006;2006:15792. DOI PubMed PMC
95. Kawarada Y, Inoue Y, Kawasaki F, et al. TGF- β induces p53/Smads complex formation in the PAI-1 promoter to activate transcription. *Sci Rep* 2016;6:35483. DOI PubMed PMC
96. Jeon H, Kim JH, Kim JH, Lee WH, Lee MS, Suk K. Plasminogen activator inhibitor type 1 regulates microglial motility and phagocytic activity. *J Neuroinflammation* 2012;9:149. DOI PubMed PMC
97. Cao C, Lawrence DA, Li Y, et al. Endocytic receptor LRP together with tPA and PAI-1 coordinates Mac-1-dependent macrophage migration. *EMBO J* 2006;25:1860-70. DOI PubMed PMC
98. Hamilton JA, Whitty GA, Wojta J, Gallichio M, McGrath K, Ianches G. Regulation of plasminogen activator inhibitor-1 levels in human monocytes. *Cell Immunol* 1993;152:7-17. DOI PubMed
99. Fabbro S, Seeds NW. Plasminogen activator activity is inhibited while neuroserpin is up-regulated in the Alzheimer disease brain. *J Neurochem* 2009;109:303-15. DOI PubMed
100. Angelucci F, Čechová K, Průša R, Hort J. Amyloid beta soluble forms and plasminogen activation system in Alzheimer's disease: Consequences on extracellular maturation of brain-derived neurotrophic factor and therapeutic implications. *CNS Neurosci Ther* 2019;25:303-13. DOI PubMed PMC
101. Mutimer CA, Keragala CB, Markus HS, Werring DJ, Cloud GC, Medcalf RL. Cerebral amyloid angiopathy and the fibrinolytic system: is plasmin a therapeutic target? *Stroke* 2021;52:2707-14. DOI PubMed
102. Park L, Zhou J, Koizumi K, et al. tPA deficiency underlies neurovascular coupling dysfunction by amyloid- β . *J Neurosci* 2020;40:8160-73. DOI PubMed PMC
103. Deane R, Zlokovic BV. Role of the blood-brain barrier in the pathogenesis of Alzheimer's disease. *Curr Alzheimer Res* 2007;4:191-7. DOI PubMed
104. Zenaro E, Piacentino G, Constantin G. The blood-brain barrier in Alzheimer's disease. *Neurobiol Dis* 2017;107:41-56. DOI PubMed PMC
105. Huang X, Hussain B, Chang J. Peripheral inflammation and blood-brain barrier disruption: effects and mechanisms. *CNS Neurosci Ther* 2021;27:36-47. DOI PubMed PMC
106. Bardehle S, Rafalski VA, Akassoglou K. Breaking boundaries-coagulation and fibrinolysis at the neurovascular interface. *Front Cell Neurosci* 2015;9:354. DOI PubMed PMC
107. Cortes-Canteli M, Mattei L, Richards AT, Norris EH, Strickland S. Fibrin deposited in the Alzheimer's disease brain promotes neuronal degeneration. *Neurobiol Aging* 2015;36:608-17. DOI PubMed PMC
108. Ryu JK, Rafalski VA, Meyer-Franke A, et al. Fibrin-targeting immunotherapy protects against neuroinflammation and neurodegeneration. *Nat Immunol* 2018;19:1212-23. DOI PubMed PMC
109. Kim JS. tPA helpers in the treatment of acute ischemic stroke: are they ready for clinical use? *J Stroke* 2019;21:160-74. DOI PubMed PMC
110. Lopes Pinheiro MA, Kooij G, Mizze MR, et al. Immune cell trafficking across the barriers of the central nervous system in multiple sclerosis and stroke. *Biochim Biophys Acta* 2016;1862:461-71. DOI PubMed
111. Sarvari S, Moakedi F, Hone E, Simpkins JW, Ren X. Mechanisms in blood-brain barrier opening and metabolism-challenged cerebrovascular ischemia with emphasis on ischemic stroke. *Metab Brain Dis* 2020;35:851-68. DOI PubMed PMC
112. Spuch C, Orlolano S, Navarro C. LRP-1 and LRP-2 receptors function in the membrane neuron. Trafficking mechanisms and proteolytic processing in Alzheimer's disease. *Front Physiol* 2012;3:269. DOI PubMed PMC
113. Cai Z, Qiao PF, Wan CQ, Cai M, Zhou NK, Li Q. Role of blood-brain barrier in Alzheimer's disease. *J Alzheimers Dis* 2018;63:1223-34. DOI PubMed
114. Lewandowski SA, Fredriksson L, Lawrence DA, Eriksson U. Pharmacological targeting of the PDGF-CC signaling pathway for blood-brain barrier restoration in neurological disorders. *Pharmacol Ther* 2016;167:108-19. DOI PubMed PMC
115. Su EJ, Cao C, Fredriksson L, et al. Microglial-mediated PDGF-CC activation increases cerebrovascular permeability during ischemic stroke. *Acta Neuropathol* 2017;134:585-604. DOI PubMed PMC
116. Benchenane K, Berezowski V, Ali C, et al. Tissue-type plasminogen activator crosses the intact blood-brain barrier by low-density lipoprotein receptor-related protein-mediated transcytosis. *Circulation* 2005;111:2241-9. DOI PubMed
117. Abu Fanne R, Nassar T, Yarovoi S, et al. Blood-brain barrier permeability and tPA-mediated neurotoxicity. *Neuropharmacology* 2010;58:972-80. DOI PubMed PMC
118. Kaur J, Tuor UI, Zhao Z, Barber PA. Quantitative MRI reveals the elderly ischemic brain is susceptible to increased early blood-brain barrier permeability following tissue plasminogen activator related to claudin 5 and occludin disassembly. *J Cereb Blood Flow Metab* 2011;31:1874-85. DOI PubMed PMC
119. Golias C, Charalabopoulos A, Stagikas D, Charalabopoulos K, Batistatou A. The kinin system--bradykinin: biological effects and clinical implications. Multiple role of the kinin system--bradykinin. *Hippokratia* 2007;11:124-8. PubMed PMC
120. Marcos-Contreras OA, Martinez de Lizarrondo S, Bardou I, et al. Hyperfibrinolysis increases blood-brain barrier permeability by a

- plasmin- and bradykinin-dependent mechanism. *Blood* 2016;128:2423-34. DOI PubMed
121. Gauberti M, Potzeka F, Vivien D, Martinez de Lizarrondo S. Impact of bradykinin generation during thrombolysis in ischemic stroke. *Front Med (Lausanne)* 2018;5:195. DOI PubMed PMC
122. Simão F, Feener EP. The effects of the contact activation system on hemorrhage. *Front Med (Lausanne)* 2017;4:121. DOI PubMed PMC
123. Calderón-Garcidueñas L, Reynoso-Robles R, Vargas-Martínez J, et al. Prefrontal white matter pathology in air pollution exposed Mexico City young urbanites and their potential impact on neurovascular unit dysfunction and the development of Alzheimer's disease. *Environ Res* 2016;146:404-17. DOI PubMed
124. Ji HL, Zhao R, Matalon S, Matthay MA. Elevated plasmin(ogen) as a common risk factor for COVID-19 susceptibility. *Physiol Rev* 2020;100:1065-75. DOI PubMed PMC
125. Nation DA, Sweeney MD, Montagne A, et al. Blood-brain barrier breakdown is an early biomarker of human cognitive dysfunction. *Nat Med* 2019;25:270-6. DOI PubMed PMC
126. Sweeney MD, Sagare AP, Zlokovic BV. Blood-brain barrier breakdown in Alzheimer disease and other neurodegenerative disorders. *Nat Rev Neurol* 2018;14:133-50. DOI PubMed PMC
127. Blennow K, Wallin A, Fredman P, Karlsson I, Gottfries CG, Svennerholm L. Blood-brain barrier disturbance in patients with Alzheimer's disease is related to vascular factors. *Acta Neurol Scand* 1990;81:323-6. DOI PubMed

Review

Open Access



Recent developments in understanding brain aging: sex differences, mechanisms, and implications in diseases

Jing Yang^{1,2,3,5,#}, Jing Qu^{2,3,4,5,#}, Huan Ma^{2,3,4,5}

¹Department of Basic Medicine, School of Medicine, Zhejiang University City College, Hangzhou 310015, Zhejiang, China.

²Department of Neurobiology, Affiliated Mental Health Center and Hangzhou Seventh People's Hospital, Zhejiang University School of Medicine, Hangzhou 310058, Zhejiang, China.

³NHC and CAMS Key Laboratory of Medical Neurobiology, MOE Frontier Science Center for Brain Research and Brain-Machine Integration, School of Brain Science and Brain Medicine, Zhejiang University, Hangzhou 310058, Zhejiang, China.

⁴Liangzhu Laboratory, The MOE Frontier Research Center of Brain & Brain-Machine Integration, Zhejiang University School of Brain Science and Brain Medicine, 1369 West Wenyi Road, Hangzhou 311121, Zhejiang, China.

⁵Research Units for Emotion and Emotion Disorders, Chinese Academy of Medical Sciences, Beijing 100730, China.

#Authors contributed equally.

Correspondence to: Prof. Huan Ma, Department of Neurobiology, Affiliated Mental Health Center and Hangzhou Seventh People's Hospital, Zhejiang University School of Medicine, 866 Yuhangtang Rd Hangzhou 310058, Zhejiang, China. E-mail: mah@zju.edu.cn; Dr. Jing Yang, Department of Basic Medicine, School of Medicine, Zhejiang University City College, 51 Huzhou Street Hangzhou 310015, Zhejiang, China. E-mail: yangjing@zucc.edu.cn

How to cite this article: Yang J, Qu J, Ma H. Recent developments in understanding brain aging: sex differences, mechanisms, and implications in diseases. *Ageing Neur Dis* 2022;2:3. <https://dx.doi.org/10.20517/and.2022.03>

Received: 14 Jan 2022 **First Decision:** 18 Feb 2022 **Revised:** 3 Mar 2022 **Accepted:** 16 Mar 2022 **Published:** 29 Mar 2022

Academic Editors: Weidong Le, Pingyi Xu **Copy Editor:** Xi-Jun Chen **Production Editor:** Xi-Jun Chen

Abstract

Exemplified by the disproportionate cases of Alzheimer's disease among women, many diseases show considerable sexual disparity in the aging process. Given that such a sex bias varies significantly in different neurological conditions, considering sex differences is necessary for the diagnosis as well as the treatment of neurological disorders. However, currently, relatively few studies have specifically focused on sex differences in brain aging or the general aging process, which has prevented the development of precision medicine for these sex-different diseases. Here, we summarize age-related disparities relating to cognitive function and dysfunction for males and females from human cross-sectional and longitudinal studies. By discussing potential anatomical and physiological bases underlying such differences, we highlight the importance of sex for aging studies in this review, which may hopefully shed light on understanding the precise causes of different brain diseases.



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, sharing, adaptation, distribution and reproduction in any medium or format, for any purpose, even commercially, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.



Keywords: Sex difference, brain aging, dementia, cognitive decline

INTRODUCTION

According to figures from the World Health Organization (WHO), the number and proportion of people aged 60 and older are expanding and estimated to increase to 1.4 billion by 2030 and 2.1 billion by 2050, up from 1 billion in 2020^[1]. The process of aging is associated with both normal and pathologic cognitive changes, which significantly affect older adults' daily life and society. The most recent data suggest that the prevalence of dementia will double in Europe and triple worldwide by 2050. Economic costs for dementia reached 957.56 billion dollars and are set to increase to 2.54 trillion dollars worldwide^[2]. Alzheimer's disease (AD) is a main cause of dementia, the related total costs of which will reach 507.49 billion dollars by 2030 in China and 1.89 trillion by 2050^[2]. Despite this, the clinical diagnosis of AD still faces many problems. There is a clear lack of precise gold standards for both diagnosis and treatment, and scientists have yet to develop multiple effective therapies for AD, especially for patients suffering in the later stages of the disease. Hence, it is clear that, for AD and other broader age-related conditions, research on aging and age-related diseases requires urgent attention.

During the human aging process, females show longer lifespans overall^[3,4] but often also display more frailty than males^[5]. For example, women aged 45-79 had a higher frailty index based on standards^[6] including 28 variables on function, cognition, co-morbidity, health attitudes and practices, and physical performance measures^[7]. This is known as the "male-female health-survival paradox"^[8], and the sex variable can make a difference for health risks in males and females. To date, sex factors have attracted wide attention in the studies of human aging. In the discoveries of brain aging, sex bias has been well-recognized in the prevalence of certain brain aging-related diseases. For example, females with AD or other dementias exhibit a two-fold incidence compared with males^[9]. Conversely, the prevalence of another progressive and age-related neurodegenerative disorder, Parkinson's disease (PD), is 1.5 times more common in men than women^[10]. In addition, sex and gender can affect the risk factors and disease progression of aging-related diseases such as AD^[11]. Thus, it is important to understand the sex difference in changes of normal brain aging, which should provide specific clues for understanding the sex-related mechanisms for age-related diseases and, in turn, may facilitate improved and personalized care during aging.

Here, we focus on reviewing the current literature reporting the sex difference in the functional changes (cognitive decline and vulnerability to neurodegenerative diseases), structural changes, and cellular hallmark changes of normal brain aging. To address this, we used the terms "sex difference", "brain aging", "cognitive aging", "brain structure", and keywords for cellular hallmarks of brain aging (mitochondria, oxidative stress, glia, ubiquitin-proteasome system, autophagy, DNA repair, stem cell exhaustion, and aberrant neuronal network activity) to search the literature in databases such as PubMed.

We first briefly review human data relating to the sex difference in cognitive decline and vulnerability to neurodegenerative diseases in the process of normal brain aging. Next, we discuss sex difference in potential anatomical changes underlying functional changes of brain aging with human evidence. Finally, we summarize the discoveries on sex difference in cellular hallmarks such as oxidative stress of brain aging from animal experiments and human data, which may offer clues for better therapeutics to cognitive decline in aging and neurodegenerative diseases.

SEXUAL DIFFERENCES IN VULNERABILITY OF MALES AND FEMALES TO COGNITIVE DECLINE IN AGING AND IN RELATED DISEASES

Sex difference of cognitive decline in normal brain aging

Cognitive function includes a variety of mental processes such as perception, attention, memory, decision making, and language comprehension. General sex differences in specific cognitive functions have been reported, with the most accepted findings being that men outperform women on spatial-based aspects, especially visual-spatial working memory tasks^[12], while females excel in verbal memory and location memory tasks^[13,14]. These differences seem to remain consistent from adolescents and young adults into older ages^[15,16]. In line with previous reports, several studies have demonstrated that certain cognitive functions decline along with the normal process of the aging of the human brain^[17,18]. Such studies monitor the trajectories of cognition change along with the process of aging in order to pick up any related changes of cognitive function that occur concurrently with stages of the aging process.

Longitudinal studies have been demonstrated as particularly useful and applicable in the study of both the difference of cognitive performance levels and the rates of cognitive change over time. In line with the results of the cross-sectional studies mentioned above, De Frias *et al.*^[19] found that women performed better on episodic memory tasks and men had higher visuospatial ability, and this sex difference was stable across age groups (35-80 years) over a 10-year period. When detecting the cognition decline rate between females and males with aging, although a review published in 2013 which screened 13 longitudinal studies concluded that no sex differences were found in the rate of overall cognitive decline between the ages of 60-80 years^[20], there are many other investigations that have shown sex differences existed in some specific cognitive tasks or in much older age (> 80) [Table 1]. Finkel *et al.*^[21], for example, found men had a faster linear decline than women on a card rotation test from middle age (50). Another study conducted by Casaletto *et al.*^[22] detected the age-related cognitive decline of 314 normal adults (average 69.3) and found that men tended to develop a declining episodic memory trajectory. Meanwhile, in recent longitudinal studies, McCarrey *et al.*^[23] administered a series of memory and other cognitive tests to participants from the Baltimore Longitudinal Study of Aging to detect the cognitive change of females and males with age. They found men showed steeper rates of decline on measures of mental status, perceptuomotor speed and integration, and visuospatial ability, but no significantly differing declines on other cognitive abilities tested compared to women^[23]. However, when analyzing increased numbers of people of the older age brackets, and after adjusting for age, education, and vascular factors, one study demonstrated that women showed a steeper decline of cognition than men after 80 years old^[24]. Finkel *et al.*^[21]'s study also demonstrated that women had a faster decline in information tests than men at ages beyond 65, with a much steeper decline after 80. However, McDowell *et al.*^[25] showed a trend for a steeper decline in men when compared with women after 80 years. Because of the complexity of human studies, it is still difficult to have a consistent conclusion, and further studies are still warranted. Nevertheless, these findings seem to indicate that women tend to show more cognitive decline at later old age than men (> 80), particularly for some for specific cognition functions, while men may have a faster cognitive decline in earlier old age (50-65).

Limitations and difficulties of human research techniques may contribute to these disparities. Firstly, the backgrounds of individuals may represent differences in aspects such as education^[24,26], lifestyle^[27], physical activities^[28], and weight^[29], and these factors not only affect the baseline of individual cognitive function but also affect the rate of cognitive decline. This makes the study on the effect of the single factor of sex on the rate of cognitive decline difficult to isolate. Moreover, in longitudinal studies, subjects are incorporated into such studies from many different age groups, in which case a limited number of subjects will be of the same age, despite a large sample of total subjects being involved. Different cohorts also show different aspects of cognitive aging^[30,31]. All these factors may cause discrepancies in results. To distinguish the true effect of sex

Table 1. Longitudinal studies for sex difference in normal cognitive decline

Ref.	Age (year)	Type of cognitive task	Decline rate
Finkel <i>et al.</i> [21]	> 65	Information test	F > M
	50-65	Card rotation test	M > F
Casaleto <i>et al.</i> [22]	47-99	The California verbal learning test	M > F
McCarrey <i>et al.</i> [23]	64.9-69.7	MMSE, fluent language production, digital symbol, card rotation	M > F
Proust-lima <i>et al.</i> [24]	> 80	Digit symbol substitution task	F > M
McDowell <i>et al.</i> [25]	> 65	Modified mini-mental state (3MS) cognitive Screening test	F > M (institution)
	> 80	Modified mini-mental state (3MS) cognitive Screening test	M > F

F: Female; M: male.

on cognitive decline with aging, larger sample sizes of confirmed similar backgrounds and the same ages should be involved, and the observation durations should be extended for these groups.

Sex differences in vulnerability to aging associated cognitive disorders

Brain aging is a natural process that results in a certain level of associated cognitive decline. However, as the brain ages, it is more susceptible to neurodegenerative diseases such as Alzheimer's disease (AD), Lewy body dementia (LBD), frontotemporal dementia (FTD), or Parkinson's disease (PD). These diseases usually occur in later life, worsening with subsequent aging. They often manifest with increasing age-related cognitive impairment, finally leading to dementia. Unlike the inconsistent results for the sex difference of cognition decline in normal brain aging, relatively consistent results have been demonstrated relating to females and males for some aspects of the vulnerabilities to these diseases [Table 2].

AD is the leading cause of dementia, which accounts for 60%-80% of dementia cases. Over the past 20 years, many studies have investigated sex differences in the risk, incidence, prevalence, or development of AD. For the analysis of risk, the Framingham Heart Study showed that, for a 65-year-old woman, the risk of AD over her remaining lifetime was 21.2%, while for a man, it was 11.6%. Correspondingly, the ratio of female to male risk for AD was noted as about 2:1 [32]. For overall numbers, many epidemiological studies of varied global locations also highlighted a higher number of women than men with AD [33,34]. The prevailing explanation for such a difference is that women live longer than men on average, and that the incidences of AD correspond with increased age [32]. The argument could therefore be formulated that any apparent sex-based difference for this disease is simply due to the increased average longevity for women. However, upon a more specific analysis of prevalence, results seem to conflict with the lifespan explanation. Plassman *et al.* [35]'s Aging, Demographics, and Memory Study (ADAMS), including subjects aged 71 years and older from all regions of the USA, showed a higher prevalence of AD in women than in men of corresponding ages. In the 2015 World Alzheimer Report, Prince *et al.* [36] showed that, in East Asia, South Asia, the Caribbean, Western Europe, and Latin America, the prevalence of dementia for men was lower than for women, although no significant difference was noted for other regions. In addition, their review in 2016, summarizing several similar studies from Europe, indicates the same higher prevalence of AD in women [37]. As for the incidence of AD, the conclusion that there is no sex difference has been noted in some studies from the United States [38,39], while others have shown that women tended to have a higher incidence of AD [40]. These disparities may stem from regional and socioeconomic factors and/or cultural effects (such as diet and lifestyle). The 10/66 Dementia Research Group study of dementia in low- and middle-income countries found a higher incidence in women [41], and most studies on European [42,43] and Asian populations [44,45] have also observed a higher incidence in women at an older age. With regards to any sex differences related to the rate of cognitive decline beyond the diagnosis of AD, many studies show that the

Table 2. Sex difference of neurodegenerative diseases

Neurodegenerative diseases	Factors	Sex difference	Ref.
Alzheimer's disease (AD) The leading cause of dementia, which accounts for 60%-80% of dementia cases	Number	F > M	Chêne et al. ^[32] Hebert et al. ^[33] Alzheimer's Association 2021 ^[34]
	Prevalence	F > M	Plassman et al. ^[35] Prince et al. ^[36] Prince et al. ^[37]
		No difference	Prince et al. ^[36] Rocca ^[152]
	Incidence	F > M	Fitzpatrick et al. ^[40] Prince et al. ^[41] Ruitenberg et al. ^[42] Rizzi et al. ^[43] Chen et al. ^[44] Yamada et al. ^[45]
		F = M (equal)	Tom et al. ^[38] Zahodne et al. ^[39]
		F > M	Hebert et al. ^[33] Holland et al. ^[46] Tifratene et al. ^[47] Lin et al. ^[48] Laws et al. ^[49] Gamberger et al. ^[50]
	Cognitive decline rate	F > M	
Lewy body dementia (LBD) The second most prevalent cause of neurodegenerative dementia	Number	M > F	Nelson et al. ^[54]
	Incidence	M > F	Savica et al. ^[55] Goodman et al. ^[56]
Frontotemporal dementia (FTD) The third most prevalent form of neurodegenerative dementias	Prevalence	M > F	Goodman et al. ^[56] Mercy et al. ^[58]
		F > M	Bernardi et al. ^[59] Ikeda et al. ^[60]
		No difference	Borroni et al. ^[61]
Parkinson's disease (PD) A movement disorder that can also lead to dementia	Prevalence	M > F	Pringsheim et al. ^[63] (worldwide) Hirsch et al. ^[64] Abbas et al. ^[65] GBD 2016 Neurology Collaborators ^[66]
		No difference	Pringsheim et al. ^[63] (Asia) Taylor et al. ^[67]
	Cognitive decline	M > F	Reekes et al. ^[71]

F: Female; M: male.

cognitive deterioration rate is faster in women than men in the progression of AD^[33,46-50]. However, despite this faster progression of the disease in women, several studies have shown that men have an overall shorter lifespan beyond diagnosis with AD, as summarized by a recent review^[51].

LBD is a neurodegenerative disease with abnormal α -synuclein accumulation (Lewy body proteins) in neurons, which can cause cognitive decline. LBD is the second most frequent form of neurodegenerative dementia. Reviews show the prevalence of LBD ranges 0.3%-24% in the general population and 3%-7% in the patients with dementia^[52,53]. In accessing the registry autopsy series from the University of Kentucky Alzheimer's Disease Center and the National Alzheimer's Coordinating Center, researchers found that the number of male patients dying with Lewy body-associated pathologies was three times that of females^[54]. Similarly, a study on the population of Olmsted County, Minnesota, showed that men had a higher incidence of LBD than women across the age spectrum^[55]. A study on the prevalence of dementia subtypes in United States Medicare showed the same result^[56]. FTD is considered the third most frequent form of

neurodegenerative dementia with more relatively young patients than other types of dementia (< 65)^[57]. While many studies have demonstrated sex differences in the prevalence of this disorder, no clear consensus has been reached. Studies in the population of Cambridgeshire, UK, and a United States Medicare study both showed greater FTD prevalence in men than women^[56,58]. However, many other studies failed to support these results^[59-61]. The discrepancy here may be due to difficulties in the exact diagnosis of FTD, which presents with similar clinical symptoms to late onset psychiatric disorders and amyotrophic lateral sclerosis (ALS). Recently, Curtis's meta-analysis focusing on the sex difference of the prevalence of genetic mutations in FTD and ALS indicated a higher prevalence of progranulin (GRN)-mutated FTD in female patients but no sex differences in chromosome 9 open reading frame 72 (C9orf72)- and microtubule-associated protein tau related FTD (MFTD), which should further help clarify the sex differences of prevalence of FTD^[62]. Another neurodegenerative disease is PD. PD is a movement disorder with bradykinesia, rigidity, tremor at rest, gait disturbance, and difficulty with speech. PD can also lead to dementia, and the proportion of patients with PD who are also diagnosed with Parkinson's disease dementia ranges from 10% to 15%. Studies are fairly consistent in demonstrating that there is a higher prevalence of PD presented among men than women from worldwide epidemiological data^[63-66], especially in Western and South American populations^[10,67-69]. However, there are reports showing the prevalence rates were almost equal between men and women in Asian populations^[63,67]. Thus, there appears to be a difference between Asian and Western populations, which may stem from sex different behaviors such as smoking, methodologies, genetics, and ethnicity^[65,70]. For the cognitive decline in PD, Reekes *et al.*^[71] indicated that males with PD have significantly greater executive and processing speed impairments compared to women.

As mentioned above, the aging brain undergoes cognitive functional change and becomes increasingly susceptible to a number of cognitive diseases. Although no results have consistently or conclusively shown differences in cognitive decline rate between females and males, apparent sex differences have been shown to be involved in the cognitive performance and disease susceptibility of the elderly. In addition, females have been demonstrated as more susceptible to AD, and males are more vulnerable to LBD and PD. The underlying mechanism for such sex-based differences in brain aging behaviors and related diseases is a key area for further study.

BRAIN STRUCTURE AS A BASIS FOR SEX DIFFERENCES IN BRAIN AGING

To explain how function changes with aging, the most widely investigated aspect is the structural changes of the aging brain. With the advantages of noninvasive imaging techniques, researchers were able to study the aging brain in healthy living individuals. As many healthy volunteers were incorporated into such studies, this provided the opportunity to analyze the sex differences of the human brain anatomy relating to aging. Using magnetic resonance imaging, many researchers found more profound age-related decline in cortical grey matter volume in males than females^[72-74]. However, there are many investigations that do not support the hypothesis that the effect of aging is accelerated in men and have failed to find age-by-sex interactions in adult and elderly populations^[75-77]. When considering the sex differences of subcortical gray matter structure in the aging brain, conclusions are no more consistent [Table 3]. The subcortical structures studied include the basal ganglia (caudate, nucleus accumbens, putamen, and pallidum), thalamus, hippocampus, and amygdala. Among these, the hippocampus is the most studied, and some hippocampus studies have reported that females have larger volumes in the aging brain^[78-80] while others have opposing results^[81]. In old age, for thalamus, some studies found the male had a larger volume^[72,80], while others opposed this^[82]. When correcting for brain volume, Li *et al.*^[78] found no significant sex difference in the relative volume of thalamus. Similarly, for the caudate, some found females had larger volume^[83,84], while others found males had larger volume^[78,80]. More consistently, the amygdala^[78,85], pallidum^[78,80], and putamen^[78,80] have been invariably found to be larger in males.

Table 3. Sex difference of subcortical regions in the aging brain of cross-sectional studies

Subcortical regions	Ref.	Age (year)	Sex difference of volume in older age (> 45)	Sex difference of decline rate in older age (> 45)
Hippocampus	Li <i>et al.</i> ^[78]	19-70	F > M (relative volume)	M > F (relative volume)
	Nemeth <i>et al.</i> ^[79]	21-58	F > M	M > F
	Wang <i>et al.</i> ^[80]	19-86	F > M (> 70)	M > F
	Goto <i>et al.</i> ^[81]	41-77	M > F (absolute volume)	F > M (absolute volume)
Thalamus	Sullivan <i>et al.</i> ^[72]	20-85	M > F	Similar
	Li <i>et al.</i> ^[78]	19-70	Similar (relative volume)	Similar (relative volume)
	Wang <i>et al.</i> ^[80]	19-86	M > F	M > F
	Takahashi <i>et al.</i> ^[82]	20 to ≥ 80	F > M	M > F
Caudate nuclei	Li <i>et al.</i> ^[78]	19-70	M > F (relative volume)	F > M (relative volume)
	Wang <i>et al.</i> ^[80]	19-86	M > F	Similar
	Good <i>et al.</i> ^[83]	18-79	F > M	No test
	Luders <i>et al.</i> ^[84]	18-82	F > M	No test
Putamen	Li <i>et al.</i> ^[78]	19-70	M > F (relative volume)	F > M (relative volume)
	Wang <i>et al.</i> ^[80]	19-86	M > F	M > F (right)
Pallium	Li <i>et al.</i> ^[78]	19-70	M > F (relative volume)	F > M (relative volume)
	Wang <i>et al.</i> ^[80]	19-86	M > F	M > F (right)
Accumbens	Wang <i>et al.</i> ^[80]	19-86	Similar	Similar
Amygdala	Li <i>et al.</i> ^[78]	19-70	M > F (relative volume)	F > M (relative volume)
	Cheng <i>et al.</i> ^[85]	20-50	M > F	No test

F: Female; M: male.

Differences in these findings may be due to differences in the age range of subjects evaluated and methods used for analysis. The previous findings of brain structure changes in the aged brain of females and males are mainly based on cross-sectional studies, which only show the status at one specific time or with different status at different specific times. Longitudinal studies may be better suited to address the conflicts in cross-sectional studies. Over the last 15 years, increasing numbers of longitudinal studies have been performed to investigate the rate of brain change with aging. Among them, some studies have paid attention to the sex difference in aging^[86-90]. Taki *et al.*^[86]'s studies showed the annual percentage change in the grey matter ratio (APC_{GMR}) in the older female group was substantially lower than in the older male group, using such a longitudinal design running over a period of over six years in 381 healthy community-dwelling individuals. Jiang *et al.*^[88] chose individuals aged 70-90 years as subjects. After a two-year follow-up, they found that women had thicker cortical regions but greater rates of cortical atrophy^[88]. For the structural changes of cortex subregions, Pfefferbaum *et al.*^[87] focused on the change of regional brain volume with aging in longitudinal studies. They found a more rapid increase of lateral ventricle volume and Sylvian fissures and more rapid decline of the centrum semiovale, anterior cingulate, parietal and precentral cortices, and thalamus in older men than older women, especially in those beyond 60 years of age^[87]. Narvacan *et al.*^[89] scanned a cohort of 55 subjects approximately three years apart. While finding that overall males had larger volumes than females for all subcortical structures, no sex differences in trajectories of change were detected^[89]. Such differences may stem from longitudinal studies which can be limited by the age range of participants, sex distribution of the samples, or scanning intervals. In a recent study, Vinke *et al.*^[90] used the Rotterdam study^[91] to understand the different aspects in brain aging of middle- and old-aged males and females based on a large prospective population-based cohort study. Their analysis showed that an earlier acceleration of decrease for normal-appearing white matter volume, gray matter volume, total brain volume, hippocampus volume, and pallidum volume and increased cerebrospinal fluid volume had occurred in men compared with women. Meanwhile, men tended to have a higher prevalence of focal

lesions (microbleeds, lacunes, and cortical infarcts) compared with women. Although shorter time intervals and less time for scanning posed some limitations for the reliable representation of the longitudinal effect of those of older ages, this study, with the largest sample used for aging-related research, provides good background information for understanding the different changes in the female and male brain due to aging.

As for other studies dealing with functional age-related change in the human brain, limitations exist in both cross-sectional and longitudinal studies for investigating how the brain's structure changes with aging. Although no consistent results have been reached for such sex differences in brain structural changes, most studies have indicated that males have accelerated atrophy in the grey matter of the cortex. This may support some findings for the faster cognitive decline in males mentioned above. The noted differences in changes for different subcortical regions may help us to understand why females outperformed males in some specific tasks while males outperformed females in others. In their review, Nemeth *et al.*^[79] demonstrated the possible functional consequences of sex difference of the subcortical grey matter, noting that their findings were relevant for dementia occurrence. However, the direct link among brain structures, cognition, and behavior is not currently clear and requires further investigation.

CELLULAR BASIS FOR SEX DIFFERENCES IN BRAIN AGING

Mattson and Arumugam^[92] 2018 paper summarizes these findings well and organizes the main aspects of brain aging into nine hallmarks: (1) mitochondrial dysfunction; (2) oxidative damage; (3) impaired cellular “waste disposal” mechanisms (autophagy-lysosome and proteasome functionality); (4) impaired adaptive stress response signaling; (5) impaired DNA repair; (6) aberrant neuronal network activity; (7) dysregulated neuronal Ca²⁺ handling; (8) stem cell exhaustion; and (9) glia cell activation and inflammation. To date, several studies (mainly from animal experiments) have shown sex differences for these hallmarks, indicating possible cellular and molecular mechanisms for sex disparity of brain aging [Figure 1].

Sex difference in mitochondrial dysfunction and oxidative stress with brain aging

The mitochondrion is an important organelle in the cell, which plays crucial roles in ATP production, storage calcium ions, and the regulation of cellular proliferation^[93]. Sex difference has been found in many aspects of brain mitochondrial function including morphology, pathways of biogenesis, autophagy, cell death, calcium, and redox homeostasis^[94]. However, such results have been mainly based on investigations of the adult brain or injured brain. Investigations on sex differences of brain-based mitochondrial dysfunction related to the normal aging process are relatively sparse. However, sex factors of redox homeostasis have been directly studied relating to brain aging, where the mechanism of the balance of free radical production and antioxidants is required to maintain redox homeostasis.

Upon aging, brain neurons tend to suffer from oxidative damage by excessive generation of free radicals and reduced antioxidant defense. Animal studies showed that, in young adult rats, females exhibit lower release of cytochrome c and lower levels of mitochondrial hydrogen peroxide than males^[95]. Moreover, in rats of similar ages, female brain mitochondria generated half the amount of peroxides and imposed dramatically less oxidative damage to mitochondrial DNA than those of males^[96]. In contrast, Guevara *et al.*^[97,98]'s studies on rat brains of different ages (6, 12, 18, and 24 months old) showed no significant sex-based differences for H₂O₂ production in any age class, despite H₂O₂ production being increased with age in both sexes.

For the antioxidant system, studies on rodents showed that young female brains have higher expression or activities of the antioxidant enzymes SOD and CAT^[99-101]. However, upon brain aging, the factors of sex difference relating to oxidative stress become complicated. Although higher antioxidant defense occurred in young female rats, after ovariectomy, mitochondrial peroxide and glutathione (GSH) levels in females




Figure 1. Sex differences implicated in cellular hallmarks of the aging brain. Shown in colored sections and dashed boxes, sex differences may exist in some hallmarks of brain aging.

became similar to those of males^[96], indicating decreased antioxidant ability in menopausal female rats. Accordingly, other non-human studies also showed higher CAT activity in the aging (18-20 months) male mouse brain and higher SOD1 protein levels in the brain of aged (28-month-old) males^[102]. Similar results were found in Human studies. Viña and Borrás^[103] proposed that young females have a better ability to fight against oxidative stress with higher antioxidant levels prior to menopause. Mandal *et al.*^[104] also found that young females (\pm 26 years old) have higher GSH levels than young men in the frontal and parietal cortex. Interestingly, the GSH levels decrease in the brains of older women (56 years old) compared with the younger women^[104], indicating that the antioxidant ability changes with aging in women. In line with this, Rekkas *et al.*^[105] found that oxidative stress in the brain increased rapidly in perimenopausal women. Bilateral oophorectomy-induced menopause was associated with increases in the GSH/GSSG ratio (increase in GSH, decrease in GSSG) and reduction in SOD and glutathione peroxidase (Gpx) mRNA expression^[106], suggesting that women had decreased antioxidant ability after menopause.

When detecting oxidative damage in the brain, several studies in rodents have demonstrated males may exhibit greater oxidative damage, expressed as higher DNA oxidation^[107] or higher levels of lipid peroxidation^[107-109], than age-matched females over relatively young ages. This may be caused by lower free radical production and higher antioxidant defense in young females. However, other studies show no such sex differences^[110,111] or higher oxidative damage in females^[100,112]. These differences may stem from the different brain regions under examination or the different times of detection. Guevara *et al.*^[97,98]'s studies on rats of different ages (6, 12, 18, and 24 months old) showed lesser oxidative damaged proteins and lipids in female brains, even in old age, an observation which they attributed in part to higher Gpx activity.

Uzun *et al.*^[113]'s study also found that protein damages were greater in male brains of 24-month-old rats. However, in the brain of 6-month-old ovariectomy mice, females showed increased levels of lipid peroxidation compared to the sham group^[114], indicating increased oxidative damage in postmenopausal females.

Overall, these findings on the sex difference in oxidative stress in the aging brain suggest that young females have a better ability to defend against oxidative stress, but these advantages may be lost upon aging, especially beyond menopause. As reviewed in Grimm *et al.*^[115]'s paper, the change of redox state with aging is associated with variation in sex steroid levels. This finding may aid in the understanding of sex differences in vulnerability to neurodegenerative disease^[115].

Sex differences in glia cell activation upon brain aging

Glial cells are distinguished from neurons in the brain and play important roles in brain function via facilitating crosstalk with neurons, maintaining the normal function of neurons, and defending against harmful stimuli. There are different types of glial cells in the brain, including astrocytes, microglia, and oligodendrocytes, each with distinct functions. Oligodendrocytes insulate axons and provide them with trophic support^[116]; microglia are regarded as macrophages participating in local immunity^[117], and astrocytes provide biochemical support of neuronal activities by facilitating the appropriate glial surroundings in conjunction with microglia^[117,118]. Various wide-ranging studies have demonstrated sex differences related to glial cells in physiological conditions or in response to pathological insults^[119-121]. However, sex differences in aged glial cells remain relatively under-studied, and the discoveries mainly stem from animal experiments.

Sex differences in microglia are the most studied sex-related aspect in the process of brain aging, with previous studies quantifying changes of microglia in the aging brain^[122,123]. Mouton *et al.*^[123]'s study showed that, in both young and old mice, females had more astrocytes and microglia in DG and CA1 of the hippocampus than did age-matched male mice. In addition, with the development of sequencing technology, recent studies have paid increasing attention to the gene expression relationships for microglia in aging^[124,125]. Mangold *et al.*^[126] also detected the sex differences for microglial gene expression in the mouse hippocampus and cortex. They found that inflammatory genes were more highly expressed in microglia of older females than in corresponding older males^[126]. Importantly, using single-cell RNA-sequencing analysis, Sala Frigerio *et al.*^[127] found that aging or progressive amyloid- β accumulation accelerated the two main activated microglia states and that female mice progressed faster in this than males, which also converged with the pathway of sex differences relating to aging and AD. In addition, Kang *et al.*^[128] mentioned in their article that tauopathy, amyloidosis, and aging had been shown to share a common APOE-driven transcriptional signature in microglia, which indicates that the increased expression of many of these transcripts of microglia in older mouse brains may be related to increased susceptibility to Alzheimer's disease in females. As regards to the functions of microglia in aging, one recent study on phagocytosis showed that aged female microglia had a greater ability for phagocytosis of neuronal debris, but they had lost their ability to adapt their phagocytic activity to inflammatory conditions^[129]. Another study on microglial function, analyzing microglial Ca²⁺ signaling and process motility, suggested "faster aging" for microglia in female mice^[130]. Taken together, the more active/faster aging microglia in older females may render them more vulnerable to some age-related neurodegenerative disease such as AD.

Relatively few studies have focused on astrocytes and oligodendrocytes in aging. Research on astrocytes has mainly been conducted in vitro via the detection of differences in the changes between the sexes under specific stimuli or in response to various pathological insults^[131]. No specific analysis of the sex differences

relating to the aging process seems to have been conducted relating to astrocytes. For oligodendrocytes, Cergnet *et al.*^[132,133] examined the sex difference of oligodendrocytes in the rat and mouse brain and found that the density of oligodendrocytes in the corpus callosum, fornix, and myelin proteins and myelin gene expression were all greater in males, and shorter life of oligodendrocytes was noted in females, a finding which did not simply represent younger mice and rats, but also held true for old mice. These differences for oligodendrocytes and myelin may be associated with the sex difference in the white matter volume in the adult and aging brain, which is discussed in “BRAIN STRUCTURE AS A BASIS FOR SEX DIFFERENCES IN BRAIN AGING”. Oligodendrocyte precursor cells can generate new mature oligodendrocytes to defend against myelin impairment in the adult brain. Transcriptomic analyses have identified sex differences in oligodendrocyte precursor cells in the expression of genes encoding for proteins involved in the cell cycle, proliferation, maturation, and myelination, among other functions^[134,135]. This difference renders older (12-month-old) female rats with greater abilities of remyelination than males after demyelination lesions^[136].

Sex differences in proteasome degradation and autophagy upon brain aging

In most organisms, a balance in the protein system, which forms the basis for gene expression and protein synthesis and degradation, is important for the normal function of cells. Aging often shifts this balance with the subsequently altered gene expressions and protein synthesis and disrupted protein degradation, resulting in some notable pathological features^[137]. In recent studies, many molecules have been demonstrated to be associated with brain aging including SFRS11^[138], CD22^[139], REST^[140], and BAZ-2 and SET-6^[141], the expressions of which change along with the aging process. However, no particular sex differences have been noted for these. Whether this is due to there not being any notable sex differences, or that this is an aspect yet to be properly examined, remains to be clarified.

Another important part of the protein system is protein degradation, which has been verified to be disrupted with brain aging^[142]. In cells, two main protein degradation systems exist, namely the ubiquitin-proteasome system (UPS) and autophagy^[143,144]. A few non-human data demonstrate that sex differences exist in these protein degradation systems of the aged brain. Ding and Keller^[145]'s study showed that proteasome inhibition occurs with aging in the central nervous system, while Zeng *et al.*^[146]'s study demonstrated that, in older (15-month-old) female mice and rats, catalytic activities of the proteasome are decreased in the cortex, striatum, cerebellum, globus pallidus, and substantia nigra with aging. In Jenkin *et al.*^[147]'s study, investigators compared the activity of the proteasome in nine tissues of young (3-5-month-old) and middle-aged (10-15-month-old) female and male mice. They found young females showed no significant differences in their proteasome activity in the brain as compared to young males. However, from middle age onwards, males showed significant decreases in proteasome activity with subsequent aging, while no such change was noted for females^[147]. One study on older fruit flies showed that basal activities of 20S proteasome had decreased in both female and male fruit-fly heads. However, in young female flies, the proteolytic capacity of the 20S proteasome could be increased via the induction of H₂O₂, but with this effect diminishing for older female flies. However, male flies showed no such age-related adaptation of the 20S proteasome^[148]. This indicated the potential for an age-related sex difference for proteasome activity in its primed adaption from external stimuli. In another recent study, old (22-month-old) male rats showed impaired fear memory with impaired UPS activity (reduced phosphorylation of the Rpt6 proteasome subunit and accumulated K48 polyubiquitinated proteins) in the basolateral amygdala (BLA), while no behavioral change was noted for old females. In this study, such changes in the activity-driven markers of UPS activity occurred within the medial prefrontal cortex but not in the BLA of old females^[149]. Such an observation of the sex differences of UPS activity in different brain regions of the aged brain aid in the understanding of sex differences related to cognitive decline with aging.

As for autophagy, while no direct articles have reported sex differences in the normal aging brain, multiple studies have demonstrated its role in AD^[150]. Many investigations on other tissues (not brain tissue) have shown that females appear to have overall lower levels of autophagy, and ovariectomized animals show increased basal levels of autophagy in several cell types^[151]. This highlights the changes in the nature and extent of autophagy after menopause in women. Although more research is needed, considering the results of Jenkin *et al.*^[147]'s study, younger females show an overall higher proteostasis capacity, where there may be a particularly well-established balance in the two types of protein degradation systems, but these could then become disrupted upon aging, rendering elderly women more vulnerable to a number of associated diseases.

For the many other hallmarks of brain aging, few studies have paid attention to any potential sex differences. Although many of the above-noted differences in the cellular changes between the female and male aging brains are not fully understood, these discoveries will likely attract increasing numbers of researchers to consider the sex factor and to further illuminate the related cellular mechanisms. These will help link the hormonal effects with those that relate to cognition and behavior, in the process of aging.

CONCLUSION

Sex differences in the brain, as they function in developmental and adult stages, have been widely investigated. However, studies on the sex factors related to the aging brain are lagging behind and deserve increasing attention, particularly under the current situation of a rapidly aging global population. Although there are no consistent general conclusions related to sex differences on cognitive decline with aging, the differences of some aspects of cognitive performance in older adults and the increased vulnerability of females and males to various aspects of dementia are becoming fairly well established, especially for AD and PD where sex is regarded as a primary risk factor for these neurodegenerative diseases. Specifically, many studies have demonstrated that women have a higher prevalence of AD than age-matched men and exhibit faster cognitive decline beyond AD diagnosis. Conversely, almost all recent studies have demonstrated that men have a higher prevalence of PD. However, for treatment, the current interventions for dementia often fail to consider the sex factor. One possible reason may be that the underlying mechanisms of sex difference for the functional changes that occur during the process of brain aging are not yet fully clear and that present techniques and other socioeconomic factors render such factors difficult to examine in detail. However, it is clear that age-related cohorts need to be established and traced to provide information about how human aging-related phenotypes and molecular changes upon aging relate to sex, in order to guide future health improvements. Beyond human-based studies, to unpack the possible structural and cellular mechanisms for sex differences of the aging brain, the use of animal models should be increasingly established, with experiments designed that can incorporate the benefits of the model animal's simple genetic background. With a better understanding of the biological mechanism for the sex differences in brain aging, we can better understand the overall functional changes in the brain, elucidate how sex creates differences in disease risk, lay a stronger foundation for dealing with the newly emerging aspects of neurodegenerative disease, explore more directly the biomarkers for brain aging, and further promote personalized medicine that incorporates the factor of sex for improved and more individualized disease treatment.

DECLARATIONS

Authors' contributions

Made contributions to conception of this review article: Yang J, Ma H

Screened and gathered articles and wrote the abstract and introduction: Qu J

Wrote the other sections and made tables: Yang J

Availability of data and materials

Not applicable.

Financial support and sponsorship

This study was supported by the National Key R&D Program of China to Ma H (grant number 2019YFA0508603); Science and Technology Innovation 2030-Major Project to Ma H (2021ZD0203501); the National Natural Science Foundation of China (grant numbers 81930030, 31771109, and 31722023 to Ma H; 81901154 to Yang J); Project for Hangzhou Medical Disciplines of Excellence; Key Project for Hangzhou Medical Disciplines.

Conflicts of interest

All authors declared that there are no conflicts of interest.

Ethical approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Copyright

© The Author(s) 2022.

REFERENCES

1. Aging and health. Available from: <https://www.who.int/news-room/fact-sheets/detail/ageing-and-health> [Last accessed on 25 Mar 2022].
2. Jia J, Wei C, Chen S, et al. The cost of Alzheimer's disease in China and re-estimation of costs worldwide. *Alzheimers Dement* 2018;14:483-91. DOI PubMed
3. Sampathkumar NK, Bravo JI, Chen Y, et al. Widespread sex dimorphism in aging and age-related diseases. *Hum Genet* 2020;139:333-56. DOI PubMed PMC
4. Bronikowski AM, Altmann J, Brockman DK, et al. Aging in the natural world: comparative data reveal similar mortality patterns across primates. *Science* 2011;331:1325-8. DOI PubMed PMC
5. Fan J, Yu C, Guo Y, et al. Frailty index and all-cause and cause-specific mortality in Chinese adults: a prospective cohort study. *Lancet Public Health* 2020;5:e650-60. DOI PubMed PMC
6. Searle SD, Mitnitski A, Gahbauer EA, Gill TM, Rockwood K. A standard procedure for creating a frailty index. *BMC Geriatr* 2008;8:24. DOI PubMed PMC
7. Gordon EH, Peel NM, Samanta M, Theou O, Howlett SE, Hubbard RE. Sex differences in frailty: a systematic review and meta-analysis. *Exp Gerontol* 2017;89:30-40. DOI PubMed
8. Oksuzyan A, Petersen I, Stovring H, Bingley P, Vaupel JW, Christensen K. The male-female health-survival paradox: a survey and register study of the impact of sex-specific selection and information bias. *Ann Epidemiol* 2009;19:504-11. DOI PubMed PMC
9. Rajan KB, Weuve J, Barnes LL, McAninch EA, Wilson RS, Evans DA. Population estimate of people with clinical Alzheimer's disease and mild cognitive impairment in the United States (2020-2060). *Alzheimers Dement* 2021;17:1966-75. DOI PubMed
10. Moisan F, Kab S, Mohamed F, et al. Parkinson disease male-to-female ratios increase with age: French nationwide study and meta-analysis. *J Neurol Neurosurg Psychiatry* 2016;87:952-7. DOI PubMed PMC
11. Nebel RA, Aggarwal NT, Barnes LL, et al. Understanding the impact of sex and gender in Alzheimer's disease: a call to action. *Alzheimers Dement* 2018;14:1171-83. DOI PubMed PMC
12. Voyer D, Voyer SD, Saint-Aubin J. Sex differences in visual-spatial working memory: a meta-analysis. *Psychon Bull Rev* 2017;24:307-34. DOI PubMed
13. Voyer D, Postma A, Brake B, Imperato-McGinley J. Gender differences in object location memory: a meta-analysis. *Psychon Bull Rev* 2007;14:23-38. DOI PubMed
14. Asperholm M, Högman N, Rafi J, Herlitz A. What did you do yesterday? *Psychol Bull* 2019;145:785-821. DOI PubMed
15. Jansen P, Kaltner S. Object-based and egocentric mental rotation performance in older adults: the importance of gender differences and motor ability. *Neuropsychol Dev Cogn B Aging Neuropsychol Cogn* 2014;21:296-316. DOI PubMed
16. Munro CA, Winicki JM, Schretlen DJ, et al. Sex differences in cognition in healthy elderly individuals. *Neuropsychol Dev Cogn B*

- Aging Neuropsychol Cogn* 2012;19:759-68. DOI PubMed PMC
17. Harada CN, Natelson Love MC, Triebel KL. Normal cognitive aging. *Clin Geriatr Med* 2013;29:737-52. DOI PubMed PMC
 18. Salthouse TA. Trajectories of normal cognitive aging. *Psychol Aging* 2019;34:17-24. DOI PubMed PMC
 19. de Frias CM, Nilsson LG, Nilsson A. Sex differences in cognition are stable over a 10-year period in adulthood and old age. *Neuropsychol Dev Cogn B Aging Neuropsychol Cogn* 2006;13:574-87. DOI
 20. Ferreira L, Ferreira Santos-Galduróz R, Ferri CP, Fernandes Galduróz JC. Rate of cognitive decline in relation to sex after 60 years-of-age: a systematic review. *Geriatr Gerontol Int* 2014;14:23-31. DOI PubMed
 21. Finkel D, Reynolds CA, Berg S, Pedersen NL. Surprising lack of sex differences in normal cognitive aging in twins. *Int J Aging Hum Dev* 2006;62:335-57. DOI PubMed
 22. Casaletto KB, Elahi FM, Staffaroni AM, et al. Cognitive aging is not created equally: differentiating unique cognitive phenotypes in “normal” adults. *Neurobiol Aging* 2019;77:13-9. DOI PubMed PMC
 23. McCarrey AC, An Y, Kitner-Triolo MH, Ferrucci L, Resnick SM. Sex differences in cognitive trajectories in clinically normal older adults. *Psychol Aging* 2016;31:166-75. DOI PubMed PMC
 24. Proust-Lima C, Amieva H, Letenneur L, Orgogozo JM, Jacqmin-Gadda H, Dartigues JF. Gender and education impact on brain aging: a general cognitive factor approach. *Psychol Aging* 2008;23:608-20. DOI PubMed
 25. McDowell I, Xi G, Lindsay J, Tuokko H. Canadian study of health and aging: study description and patterns of early cognitive decline. *Aging, Neuropsychology, and Cognition* 2004;11:149-68. DOI
 26. Wilson RS, Hebert LE, Scherr PA, Barnes LL, Mendes de Leon CF, Evans DA. Educational attainment and cognitive decline in old age. *Neurology* 2009;72:460-5. DOI PubMed PMC
 27. Yu F, Ryan LH, Schaie KW, Willis SL, Kolanowski A. Factors associated with cognition in adults: the Seattle Longitudinal Study. *Res Nurs Health* 2009;32:540-50. DOI PubMed PMC
 28. Finkel D, Andel R, Gatz M, Pedersen NL. The role of occupational complexity in trajectories of cognitive aging before and after retirement. *Psychol Aging* 2009;24:563-73. DOI PubMed PMC
 29. Noh HM, Han J, Kim YJ, Jung JH, Roh YK, Song HJ. Sex differences in the relationship between cognitive impairment and overweight or obesity in late life: a 3-year prospective study. *Medicine (Baltimore)* 2019;98:e14736. DOI PubMed PMC
 30. Gerstorf D, Ram N, Hoppmann C, Willis SL, Schaie KW. Cohort differences in cognitive aging and terminal decline in the Seattle Longitudinal Study. *Dev Psychol* 2011;47:1026-41. DOI PubMed PMC
 31. Karlsson P, Thorvaldsson V, Skoog I, Gudmundsson P, Johansson B. Birth cohort differences in fluid cognition in old age: comparisons of trends in levels and change trajectories over 30 years in three population-based samples. *Psychol Aging* 2015;30:83-94. DOI PubMed
 32. Chêne G, Beiser A, Au G, et al. Gender and incidence of dementia in the Framingham Heart Study from mid-adult life. *Alzheimers Dement* 2015;11:310-20. DOI PubMed PMC
 33. Hebert LE, Weuve J, Scherr PA, Evans DA. Alzheimer disease in the United States (2010-2050) estimated using the 2010 census. *Neurology* 2013;80:1778-83. DOI PubMed PMC
 34. 2021 Alzheimer's disease facts and figures. *Alzheimers Dement* 2021;17:327-406. DOI PubMed
 35. Plassman BL, Langa KM, Fisher GG, et al. Prevalence of dementia in the United States: the aging, demographics, and memory study. *Neuroepidemiology* 2007;29:125-32. DOI PubMed PMC
 36. Wimo A, Ali GC, Guerchet M, Prince M, Prina M, Wu YT; Alzheimer's Disease International. World Alzheimer Report 2015 - The Global Impact of Dementia: An analysis of prevalence, incidence, cost and trends. Available from: <https://www.alzint.org/resource/world-alzheimer-report-2015/> [Last accessed on 25 Mar 2022].
 37. Prince M, Ali GC, Guerchet M, Prina AM, Albanese E, Wu YT. Recent global trends in the prevalence and incidence of dementia, and survival with dementia. *Alzheimers Res Ther* 2016;8:23. DOI PubMed PMC
 38. Tom SE, Hubbard RA, Crane PK, et al. Characterization of dementia and Alzheimer's disease in an older population: updated incidence and life expectancy with and without dementia. *Am J Public Health* 2015;105:408-13. DOI PubMed PMC
 39. Zahodne LB, Schofield PW, Farrell MT, Stern Y, Manly JJ. Bilingualism does not alter cognitive decline or dementia risk among Spanish-speaking immigrants. *Neuropsychology* 2014;28:238-46. DOI PubMed PMC
 40. Fitzpatrick AL, Kuller LH, Ives DG, et al. Incidence and prevalence of dementia in the Cardiovascular Health Study. *J Am Geriatr Soc* 2004;52:195-204. DOI PubMed
 41. Prince M, Acosta D, Ferri CP, et al. Dementia incidence and mortality in middle-income countries, and associations with indicators of cognitive reserve: a 10/66 Dementia Research Group population-based cohort study. *Lancet* 2012;380:50-8. DOI PubMed PMC
 42. Ruitenberg A, Ott A, van Swieten JC, Hofman A, Breteler MM. Incidence of dementia: does gender make a difference? *Neurobiol Aging* 2001;22:575-80. DOI PubMed
 43. Rizzi L, Rosset I, Roriz-Cruz M. Global epidemiology of dementia: Alzheimer's and vascular types. *Biomed Res Int* 2014;2014:908915. DOI PubMed PMC
 44. Chen R, Hu Z, Wei L, Ma Y, Liu Z, Copeland JR. Incident dementia in a defined older Chinese population. *PLoS One* 2011;6:e24817. DOI PubMed PMC
 45. Yamada M, Mimori Y, Kasagi F, et al. Incidence of dementia, Alzheimer disease, and vascular dementia in a Japanese population: Radiation Effects Research Foundation adult health study. *Neuroepidemiology* 2008;30:152-60. DOI PubMed
 46. Holland D, Desikan RS, Dale AM, McEvoy LK; Alzheimer's Disease Neuroimaging Initiative. Higher rates of decline for women and apolipoprotein E epsilon4 carriers. *AJNR Am J Neuroradiol* 2013;34:2287-93. DOI PubMed PMC
 47. Tifratene K, Robert P, Metelkina A, Pradier C, Dartigues JF. Progression of mild cognitive impairment to dementia due to AD in

- clinical settings. *Neurology* 2015;85:331-8. DOI PubMed
48. Lin FC, Chuang YS, Hsieh HM, et al. Early statin use and the progression of Alzheimer disease: a total population-based case-control study. *Medicine (Baltimore)* 2015;94:e2143. DOI PubMed PMC
49. Laws KR, Irvine K, Gale TM. Sex differences in cognitive impairment in Alzheimer's disease. *World J Psychiatry* 2016;6:54-65. DOI PubMed PMC
50. Gamberger D, Lavrač N, Srivatsa S, Tanzi RE, Doraiswamy PM. Identification of clusters of rapid and slow decliners among subjects at risk for Alzheimer's disease. *Sci Rep* 2017;7:6763. DOI PubMed PMC
51. Todd S, Barr S, Roberts M, Passmore AP. Survival in dementia and predictors of mortality: a review. *Int J Geriatr Psychiatry* 2013;28:1109-24. DOI PubMed
52. Vann Jones SA, O'Brien JT. The prevalence and incidence of dementia with Lewy bodies: a systematic review of population and clinical studies. *Psychol Med* 2014;44:673-83. DOI PubMed
53. Hogan DB, Fiest KM, Roberts JI, et al. The prevalence and incidence of dementia with Lewy bodies: a systematic review. *Can J Neurol Sci* 2016;43 Suppl 1:S83-95. DOI PubMed
54. Nelson PT, Schmitt FA, Jicha GA, et al. Association between male gender and cortical Lewy body pathology in large autopsy series. *J Neurol* 2010;257:1875-81. DOI PubMed PMC
55. Savica R, Grossardt BR, Bower JH, Boeve BF, Ahlskog JE, Rocca WA. Incidence of dementia with Lewy bodies and Parkinson disease dementia. *JAMA Neurol* 2013;70:1396-402. DOI PubMed PMC
56. Goodman RA, Lochner KA, Thambisetty M, Wingo TS, Posner SF, Ling SM. Prevalence of dementia subtypes in United States Medicare fee-for-service beneficiaries, 2011-2013. *Alzheimers Dement* 2017;13:28-37. DOI PubMed PMC
57. Bang J, Spina S, Miller BL. Frontotemporal dementia. *Lancet* 2015;386:1672-82. DOI PubMed PMC
58. Mercy L, Hodges JR, Dawson K, Barker RA, Brayne C. Incidence of early-onset dementias in Cambridgeshire, United Kingdom. *Neurology* 2008;71:1496-9. DOI PubMed
59. Bernardi L, Frangipane F, Smirne N, et al. Epidemiology and genetics of frontotemporal dementia: a door-to-door survey in southern Italy. *Neurobiol Aging* 2012;33:2948.e1-2948.e10. DOI PubMed PMC
60. Ikeda M, Ishikawa T, Tanabe H. Epidemiology of frontotemporal lobar degeneration. *Dement Geriatr Cogn Disord* 2004;17:265-8. DOI PubMed
61. Borroni B, Alberici A, Grassi M, et al. Is frontotemporal lobar degeneration a rare disorder? *J Alzheimers Dis* 2010;19:111-6. DOI PubMed
62. Curtis AF, Masellis M, Hsiung GR, et al. Sex differences in the prevalence of genetic mutations in FTD and ALS: a meta-analysis. *Neurology* 2017;89:1633-42. DOI PubMed PMC
63. Pringsheim T, Jette N, Frolkis A, Steeves TD. The prevalence of Parkinson's disease: a systematic review and meta-analysis. *Mov Disord* 2014;29:1583-90. DOI PubMed
64. Hirsch L, Jette N, Frolkis A, Steeves T, Pringsheim T. The incidence of Parkinson's disease: a systematic review and meta-analysis. *Neuroepidemiology* 2016;46:292-300. DOI PubMed
65. Abbas MM, Xu Z, Tan LCS. Epidemiology of Parkinson's disease-East versus West. *Mov Disord Clin Pract* 2018;5:14-28. DOI PubMed PMC
66. GBD 2016 Neurology Collaborators. Global, regional, and national burden of neurological disorders, 1990-2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet Neurol* 2019;18:459-80. DOI PubMed PMC
67. Taylor KS, Cook JA, Counsell CE. Heterogeneity in male to female risk for Parkinson's disease. *J Neurol Neurosurg Psychiatry* 2007;78:905-6. DOI PubMed PMC
68. de Lau LM, Verbaan D, Marinus J, van Hilten JJ. Survival in Parkinson's disease. Relation with motor and non-motor features. *Parkinsonism Relat Disord* 2014;20:613-6. DOI PubMed
69. Pinter B, Diem-Zangerl A, Wenning GK, et al. Mortality in Parkinson's disease: a 38-year follow-up study. *Mov Disord* 2015;30:266-9. DOI PubMed
70. Ascherio A, Schwarzschild MA. The epidemiology of Parkinson's disease: risk factors and prevention. *Lancet Neurol* 2016;15:1257-72. DOI PubMed
71. Reekes TH, Higginson CI, Ledbetter CR, Sathivadivel N, Zweig RM, Disbrow EA. Sex specific cognitive differences in Parkinson disease. *NPJ Parkinsons Dis* 2020;6:7. DOI PubMed PMC
72. Sullivan EV, Rosenbloom M, Serventi KL, Pfefferbaum A. Effects of age and sex on volumes of the thalamus, pons, and cortex. *Neurobiol Aging* 2004;25:185-92. DOI PubMed
73. DeCarli C, Massaro J, Harvey D, et al. Measures of brain morphology and infarction in the framingham heart study: establishing what is normal. *Neurobiol Aging* 2005;26:491-510. DOI PubMed
74. Ikram MA, Vrooman HA, Vernooij MW, et al. Brain tissue volumes in the general elderly population. The Rotterdam Scan Study. *Neurobiol Aging* 2008;29:882-90. DOI PubMed
75. Salat DH, Tuch DS, Greve DN, et al. Age-related alterations in white matter microstructure measured by diffusion tensor imaging. *Neurobiol Aging* 2005;26:1215-27. DOI PubMed
76. Lemaître H, Crivello F, Gratiot B, Alperovitch A, Tzourio C, Mazoyer B. Age- and sex-related effects on the neuroanatomy of healthy elderly. *Neuroimage* 2005;26:900-11. DOI PubMed
77. Jäncke L, Mérillat S, Liem F, Hänggi J. Brain size, sex, and the aging brain. *Hum Brain Mapp* 2015;36:150-69. DOI PubMed PMC
78. Li W, van Tol MJ, Li M, et al. Regional specificity of sex effects on subcortical volumes across the lifespan in healthy aging. *Hum Brain Mapp* 2014;35:238-47. DOI PubMed PMC

79. Nemeth VL, Must A, Horvath S, Király A, Kincses ZT, Vécsei L. Gender-specific degeneration of dementia-related subcortical structures throughout the lifespan. *J Alzheimers Dis* 2017;55:865-80. DOI PubMed
80. Wang Y, Xu Q, Luo J, Hu M, Zuo C. Effects of age and sex on subcortical volumes. *Front Aging Neurosci* 2019;11:259. DOI PubMed PMC
81. Goto M, Abe O, Miyati T, et al. Accelerated hippocampal volume reduction in post-menopausal women: an additional study with Atlas-based method. *Radiol Phys Technol* 2011;4:185-8. DOI PubMed
82. Takahashi R, Ishii K, Kakigi T, Yokoyama K. Gender and age differences in normal adult human brain: voxel-based morphometric study. *Hum Brain Mapp* 2011;32:1050-8. DOI PubMed PMC
83. Good CD, Johnsrude I, Ashburner J, Henson RN, Friston KJ, Frackowiak RS. Cerebral asymmetry and the effects of sex and handedness on brain structure: a voxel-based morphometric analysis of 465 normal adult human brains. *Neuroimage* 2001;14:685-700. DOI PubMed
84. Luders E, Gaser C, Narr KL, Toga AW. Why sex matters: brain size independent differences in gray matter distributions between men and women. *J Neurosci* 2009;29:14265-70. DOI PubMed PMC
85. Cheng Y, Chou KH, Decety J, et al. Sex differences in the neuroanatomy of human mirror-neuron system: a voxel-based morphometric investigation. *Neuroscience* 2009;158:713-20. DOI PubMed
86. Taki Y, Kinomura S, Sato K, Kawashima R, Fukuda H. A longitudinal study of gray matter volume decline with age and modifying factors. *Neurobiol Aging* 2011;32:907-15. DOI PubMed
87. Pfefferbaum A, Rohlfing T, Rosenbloom MJ, Chu W, Colrain IM, Sullivan EV. Variation in longitudinal trajectories of regional brain volumes of healthy men and women (ages 10 to 85 years) measured with atlas-based parcellation of MRI. *Neuroimage* 2013;65:176-93. DOI PubMed PMC
88. Jiang J, Sachdev P, Lipnicki DM, et al. A longitudinal study of brain atrophy over two years in community-dwelling older individuals. *Neuroimage* 2014;86:203-11. DOI PubMed
89. Narvacan K, Treit S, Camicioli R, Martin W, Beaulieu C. Evolution of deep gray matter volume across the human lifespan. *Hum Brain Mapp* 2017;38:3771-90. DOI PubMed PMC
90. Vinke EJ, de Groot M, Venkatraghavan V, et al. Trajectories of imaging markers in brain aging: the Rotterdam Study. *Neurobiol Aging* 2018;71:32-40. DOI PubMed
91. Ikram MA, Brusselle GGO, Murad SD, et al. The Rotterdam Study: 2018 update on objectives, design and main results. *Eur J Epidemiol* 2017;32:807-50. DOI PubMed PMC
92. Mattson MP, Arumugam TV. Hallmarks of brain aging: adaptive and Pathological modification by metabolic states. *Cell Metab* 2018;27:1176-99. DOI PubMed PMC
93. Rizzuto R, De Stefani D, Raffaello A, Mammucari C. Mitochondria as sensors and regulators of calcium signalling. *Nat Rev Mol Cell Biol* 2012;13:566-78. DOI PubMed
94. Demarest TG, McCarthy MM. Sex differences in mitochondrial (dys)function: Implications for neuroprotection. *J Bioenerg Biomembr* 2015;47:173-88. DOI PubMed PMC
95. Lloret A, Badía MC, Mora NJ, et al. Gender and age-dependent differences in the mitochondrial apoptogenic pathway in Alzheimer's disease. *Free Radic Biol Med* 2008;44:2019-25. DOI PubMed
96. Borrás C, Sastre J, Garcia-sala D, Lloret A, Pallardó FV, Viña J. Mitochondria from females exhibit higher antioxidant gene expression and lower oxidative damage than males. *Free Radic Biol Med* 2003;34:546-52. DOI PubMed
97. Guevara R, Santandreu FM, Valle A, Gianotti M, Oliver J, Roca P. Sex-dependent differences in aged rat brain mitochondrial function and oxidative stress. *Free Radic Biol Med* 2009;46:169-75. DOI PubMed
98. Guevara R, Gianotti M, Oliver J, Roca P. Age and sex-related changes in rat brain mitochondrial oxidative status. *Exp Gerontol* 2011;46:923-8. DOI PubMed
99. Gagnard P, Savouroux S, Liere P, et al. Effect of sex differences on brain mitochondrial function and its suppression by ovariectomy and in aged mice. *Endocrinology* 2015;156:2893-904. DOI PubMed
100. Dkhil MA, Al-Shaebi EM, Lubbad MY, Al-Quraishy S. Impact of sex differences in brain response to infection with Plasmodium berghei. *Parasitol Res* 2016;115:415-22. DOI PubMed
101. Khalifa AR, Abdel-Rahman EA, Mahmoud AM, et al. Sex-specific differences in mitochondria biogenesis, morphology, respiratory function, and ROS homeostasis in young mouse heart and brain. *Physiol Rep* 2017;5:e13125. DOI PubMed PMC
102. Ali SS, Xiong C, Lucero J, Behrens MM, Dugan LL, Quick KL. Gender differences in free radical homeostasis during aging: shorter-lived female C57BL6 mice have increased oxidative stress. *Aging Cell* 2006;5:565-74. DOI PubMed
103. Viña J, Borrás C. Women live longer than men: understanding molecular mechanisms offers opportunities to intervene by using estrogenic compounds. *Antioxid Redox Signal* 2010;13:269-78. DOI PubMed
104. Mandal PK, Tripathi M, Sugunan S. Brain oxidative stress: detection and mapping of anti-oxidant marker 'Glutathione' in different brain regions of healthy male/female, MCI and Alzheimer patients using non-invasive magnetic resonance spectroscopy. *Biochem Biophys Res Commun* 2012;417:43-8. DOI PubMed
105. Rekkas PV, Wilson AA, Lee VW, et al. Greater monoamine oxidase a binding in perimenopausal age as measured with carbon 11-labeled harmine positron emission tomography. *JAMA Psychiatry* 2014;71:873-9. DOI PubMed PMC
106. Bellanti F, Matteo M, Rollo T, et al. Sex hormones modulate circulating antioxidant enzymes: impact of estrogen therapy. *Redox Biol* 2013;1:340-6. DOI PubMed PMC
107. Candeias E, Duarte AI, Sebastião I, et al. Middle-aged diabetic females and males present distinct susceptibility to Alzheimer disease-like pathology. *Mol Neurobiol* 2017;54:6471-89. DOI PubMed

108. Sobocanec S, Balog T, Kusić B, Sverko V, Sarić A, Marotti T. Differential response to lipid peroxidation in male and female mice with age: correlation of antioxidant enzymes matters. *Biogerontology* 2008;9:335-43. DOI PubMed
109. Silva TLA, Braz GRF, Silva SCA, et al. Serotonin transporter inhibition during neonatal period induces sex-dependent effects on mitochondrial bioenergetics in the rat brainstem. *Eur J Neurosci* 2018;1620-34. DOI PubMed
110. Cole TB, Coburn J, Dao K, et al. Sex and genetic differences in the effects of acute diesel exhaust exposure on inflammation and oxidative stress in mouse brain. *Toxicology* 2016;374:1-9. DOI PubMed PMC
111. Monte AS, Mello BSF, Borella VCM, et al. Two-hit model of schizophrenia induced by neonatal immune activation and peripubertal stress in rats: study of sex differences and brain oxidative alterations. *Behav Brain Res* 2017;331:30-7. DOI PubMed
112. Al-Gubory KH, Garrel C. Sex-specific divergence of antioxidant pathways in fetal brain, liver, and skeletal muscles. *Free Radic Res* 2016;50:366-73. DOI PubMed
113. Uzun H, Kayali R, Cakatay U. The chance of gender dependency of oxidation of brain proteins in aged rats. *Arch Gerontol Geriatr* 2010;50:16-9. DOI PubMed
114. Yao J, Irwin R, Chen S, Hamilton R, Cadenas E, Brinton RD. Ovarian hormone loss induces bioenergetic deficits and mitochondrial β -amyloid. *Neurobiol Aging* 2012;33:1507-21. DOI PubMed PMC
115. Grimm A, Mensah-Nyagan AG, Eckert A. Alzheimer, mitochondria and gender. *Neurosci Biobehav Rev* 2016;67:89-101. DOI PubMed
116. Serrano-Regal MP, Bayón-Cordero L, Ordaz RP, et al. Expression and function of GABA receptors in myelinating cells. *Front Cell Neurosci* 2020;14:256. DOI PubMed PMC
117. Bernaus A, Blanco S, Sevilla A. Glia crosstalk in neuroinflammatory diseases. *Front Cell Neurosci* 2020;14:209. DOI PubMed PMC
118. Augusto-Oliveira M, Arrifano GP, Takeda PY, et al. Astroglia-specific contributions to the regulation of synapses, cognition and behaviour. *Neurosci Biobehav Rev* 2020;118:331-57. DOI PubMed
119. Thion MS, Low D, Silvín A, et al. Microbiome influences prenatal and adult microglia in a sex-specific manner. *Cell* 2018;172:500-516.e16. DOI PubMed PMC
120. Villa A, Gelosa P, Castiglioni L, et al. Sex-specific features of microglia from adult mice. *Cell Rep* 2018;23:3501-11. DOI PubMed PMC
121. Hanamsagar R, Alter MD, Block CS, Sullivan H, Bolton JL, Bilbo SD. Generation of a microglial developmental index in mice and in humans reveals a sex difference in maturation and immune reactivity. *Glia* 2017;65:1504-20. DOI PubMed PMC
122. Hayakawa N, Kato H, Araki T. Age-related changes of astrocytes, oligodendrocytes and microglia in the mouse hippocampal CA1 sector. *Mech Ageing Dev* 2007;128:311-6. DOI PubMed
123. Mouton PR, Long JM, Lei D, et al. Age and gender effects on microglia and astrocyte numbers in brains of mice. *Brain Res* 2002;956:30-5. DOI PubMed
124. Sierra A, Gottfried-Blackmore AC, McEwen BS, Bulloch K. Microglia derived from aging mice exhibit an altered inflammatory profile. *Glia* 2007;55:412-24. DOI PubMed
125. Bonham LW, Sirkis DW, Yokoyama JS. The transcriptional landscape of microglial genes in aging and neurodegenerative disease. *Front Immunol* 2019;10:1170. DOI PubMed PMC
126. Mangold CA, Wronowski B, Du M, et al. Sexually divergent induction of microglial-associated neuroinflammation with hippocampal aging. *J Neuroinflammation* 2017;14:141. DOI PubMed PMC
127. Sala Frigerio C, Wolfs L, Fattorelli N, et al. The major risk factors for Alzheimer's disease: age, sex, and genes modulate the microglia response to A β plaques. *Cell Rep* 2019;27:1293-1306.e6. DOI PubMed PMC
128. Kang SS, Ebbert MTW, Baker KE, et al. Microglial translational profiling reveals a convergent APOE pathway from aging, amyloid, and tau. *J Exp Med* 2018;215:2235-45. DOI PubMed PMC
129. Yanguas-Casás N, Crespo-Castrillo A, Arevalo MA, Garcia-Segura LM. Aging and sex: impact on microglia phagocytosis. *Aging Cell* 2020;19:e13182. DOI PubMed PMC
130. Del Moral M, Fröhlich N, Figarella K, Mojtahedi N, Garaschuk O. Effect of caloric restriction on the in vivo functional properties of aging microglia. *Front Immunol* 2020;11:750. DOI PubMed PMC
131. Chowen JA, Garcia-Segura LM. Role of glial cells in the generation of sex differences in neurodegenerative diseases and brain aging. *Mech Ageing Dev* 2021;196:111473. DOI PubMed
132. Cerghet M, Skoff RP, Bessert D, Zhang Z, Mullins C, Ghandour MS. Proliferation and death of oligodendrocytes and myelin proteins are differentially regulated in male and female rodents. *J Neurosci* 2006;26:1439-47. DOI PubMed PMC
133. Cerghet M, Skoff RP, Swamydas M, Bessert D. Sexual dimorphism in the white matter of rodents. *J Neurol Sci* 2009;286:76-80. DOI PubMed PMC
134. Young KM, Psachoulia K, Tripathi RB, et al. Oligodendrocyte dynamics in the healthy adult CNS: evidence for myelin remodeling. *Neuron* 2013;77:873-85. DOI PubMed PMC
135. Yasuda K, Maki T, Kinoshita H, et al. Sex-specific differences in transcriptomic profiles and cellular characteristics of oligodendrocyte precursor cells. *Stem Cell Res* 2020;46:101866. DOI PubMed
136. Li WW, Penderis J, Zhao C, Schumacher M, Franklin RJ. Females remyelinate more efficiently than males following demyelination in the aged but not young adult CNS. *Exp Neurol* 2006;202:250-4. DOI
137. Anisimova AS, Alexandrov AI, Makarova NE, Gladyshev VN, Dmitriev SE. Protein synthesis and quality control in aging. *Aging (Albany NY)* 2018;10:4269-88. DOI PubMed PMC
138. Raihan O, Brishti A, Li Q, et al. SFRS11 loss leads to aging-associated cognitive decline by modulating LRP8 and ApoE. *Cell Rep*

- 2019;28:78-90.e6. [DOI](#) [PubMed](#)
139. Pluvinage JV, Haney MS, Smith BAH, et al. CD22 blockade restores homeostatic microglial phagocytosis in ageing brains. *Nature* 2019;568:187-92. [DOI](#) [PubMed](#) [PMC](#)
140. Zullo JM, Drake D, Aron L, et al. Regulation of lifespan by neural excitation and REST. *Nature* 2019;574:359-64. [DOI](#) [PubMed](#) [PMC](#)
141. Yuan J, Chang SY, Yin SG, et al. Two conserved epigenetic regulators prevent healthy ageing. *Nature* 2020;579:118-22. [DOI](#) [PubMed](#)
142. Tsuji T, Shimohama S. Protein degradation in Alzheimer's disease and aging of the brain. *Prog Mol Subcell Biol* 2002;29:43-60. [DOI](#) [PubMed](#)
143. Galluzzi L, Baehrecke EH, Ballabio A, et al. Molecular definitions of autophagy and related processes. *EMBO J* 2017;36:1811-36. [DOI](#) [PubMed](#) [PMC](#)
144. VerPlank JJS, Goldberg AL. Regulating protein breakdown through proteasome phosphorylation. *Biochem J* 2017;474:3355-71. [DOI](#) [PubMed](#) [PMC](#)
145. Ding Q, Keller JN. Proteasomes and proteasome inhibition in the central nervous system. *Free Radic Biol Med* 2001;31:574-84. [DOI](#) [PubMed](#)
146. Zeng BY, Medhurst AD, Jackson M, Rose S, Jenner P. Proteasomal activity in brain differs between species and brain regions and changes with age. *Mech Ageing Dev* 2005;126:760-6. [DOI](#) [PubMed](#)
147. Jenkins EC, Shah N, Gomez M, et al. Proteasome mapping reveals sexual dimorphism in tissue-specific sensitivity to protein aggregations. *EMBO Rep* 2020;21:e48978. [DOI](#) [PubMed](#) [PMC](#)
148. Pomatto LCD, Wong S, Carney C, Shen B, Tower J, Davies KJA. The age- and sex-specific decline of the 20s proteasome and the Nrf2/CncC signal transduction pathway in adaption and resistance to oxidative stress in *Drosophila melanogaster*. *Aging (Albany NY)* 2017;9:1153-85. [DOI](#) [PubMed](#) [PMC](#)
149. Dulka BN, Trask S, Helmstetter FJ. Age-related memory impairment and sex-specific alterations in phosphorylation of the Rpt6 proteasome subunit and polyubiquitination in the basolateral amygdala and medial prefrontal cortex. *Front Aging Neurosci* 2021;13:656944. [DOI](#) [PubMed](#) [PMC](#)
150. Li Q, Liu Y, Sun M. Autophagy and Alzheimer's disease. *Cell Mol Neurobiol* 2017;37:377-88. [DOI](#) [PubMed](#)
151. Congdon EE. Sex differences in autophagy contribute to female vulnerability in Alzheimer's disease. *Front Neurosci* 2018;12:372. [DOI](#) [PubMed](#) [PMC](#)
152. Rocca WA. Time, sex, gender, history, and dementia. *Alzheimer Dis Assoc Disord* 2017;31:76-9. [DOI](#) [PubMed](#) [PMC](#)

Review

Open Access



Influence of sleep disruption on protein accumulation in neurodegenerative diseases

Xiying Wang^{1,2}, Rui Wang³, Jiada Li^{1,4}

¹Center for Medical Genetics, School of Life Sciences, Central South University, Changsha 410078, Hunan, China.

²Xiangya School of Medicine, Central South University, Changsha 410078, Hunan, China.

³School of Life Science and Technology, Key Laboratory of Developmental Genes and Human Disease, Southeast University, Nanjing 210096, Jiangsu, China.

⁴Hunan Key Laboratory of Animal Models for Human Diseases, and Hunan Key Laboratory of Medical Genetics, Changsha 410078, Hunan, China.

Correspondence to: Prof. Jiada Li, Center for Medical Genetics, School of Life Sciences, Central South University, 172 Tong-Zi-Po Road, Changsha 410078, Hunan, China. E-mail: lijada@sklmg.edu.cn

How to cite this article: Wang X, Wang R, Li J. Influence of sleep disruption on protein accumulation in neurodegenerative diseases. *Ageing Neur Dis* 2022;2:4. <https://dx.doi.org/10.20517/and.2021.10>

Received: 29 Dec 2021 **First Decision:** 1 Mar 2022 **Revised:** 10 Mar 2022 **Accepted:** 23 Mar 2022 **Published:** 31 Mar 2022

Academic Editor: Weidong Le **Copy Editor:** Xi-Jun Chen **Production Editor:** Xi-Jun Chen

Abstract

Abnormal accumulation of disease proteins in the central nervous system is a neuropathological feature in neurodegenerative disorders. Recently, a growing body of evidence has supported a role of disruption of the sleep-wake cycle in disease development, pathological changes and abnormal protein accumulation in neurodegenerative diseases, especially in Alzheimer's disease and Parkinson's disease. Sleep deprivation promotes abnormal accumulation of disease proteins. Interestingly, amyloid- β ($A\beta$) has daily oscillations in human cerebral spinal fluid (CSF) and is cleared more in sleep. Both circadian genes and circadian hormones are associated with disease protein deposition. Recently, the glymphatic pathway and meningeal lymphatics have been shown to play a critical role in $A\beta$ clearance, which is mediated by the aquaporin (AQP-4) water channel on astrocytes. The rate of the clearance of $A\beta$ by the glymphatic pathway is different during the sleep/wake cycle. Most importantly, circadian rhythms facilitate glymphatic clearance of solutes and $A\beta$ in the CSF and interstitial fluid in an AQP-4-dependent manner, which further provides evidence for the involvement of circadian rhythms in disease protein clearance.



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, sharing, adaptation, distribution and reproduction in any medium or format, for any purpose, even commercially, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.



Keywords: Neurodegenerative diseases, protein accumulation, glymphatic clearance, Alzheimer's disease, Parkinson's disease, circadian disruption

INTRODUCTION

The sleep-wake cycle is controlled by circadian rhythms. Circadian rhythms are biological processes that follow a daily cycle and respond to light and darkness in an organism's environment. Circadian rhythms are endogenously synchronized to the day/night cycle to form oscillations of 24 h^[1]. The central pacemaker in mammals is the suprachiasmatic nucleus (SCN), which receives light signals from the retina through the retinohypothalamic pathway^[2]. Ablation of the SCN in animals, aneurysms near the SCN, or pituitary tumors in patients disrupts daily rhythms^[3-5]. Moreover, intrahypothalamic grafts of neonatal SCN tissue to SCN-ablated animals restore rest-activity rhythms^[3]. Thus, the SCN plays a principal role in circadian rhythms.

The SCN clock entrains the peripheral clock, nearly all of the cells in the body, through autonomic innervation, body temperature, humoral signals, and feeding-related cues^[6]. The rhythms of the cells from different organs are affected not only by the oscillation of hormones, the nervous system and temperature but also by local circadian oscillators^[1]. Increasing lines of evidence demonstrate that dysfunction of circadian rhythms is associated with diseases in many systems and organs, including neurodegenerative diseases^[7,8], cancer^[9], hypertension^[10], diabetes^[11], and autoimmune diseases^[12]. The circadian rhythms in mammals are generated internally but synchronized to the external environment (light and dark cycles by the Earth's 24 h rotation^[13]). Sleep/wake cycles have a typical 24 h pattern, which is linked to circadian rhythms^[13]. Sleep loss is a representative symptom of circadian rhythm disruption.

Aberrant accumulation of abnormal disease proteins is a common pathological feature of neurodegenerative diseases. The aggregation of proteins associated with neurodegenerative disease is present in patient brains in both familial and sporadic cases, such as the formation of extracellular plaques by β -amyloid ($A\beta$) and intracellular neurofibrillary tangles by tau in Alzheimer's disease (AD), intracellular Lewy bodies by α -synuclein (α -syn) in Parkinson's disease (PD), TDP-43 inclusions in amyotrophic lateral sclerosis and nuclear inclusions in Huntington disease^[14-16]. Circadian rhythm abnormalities are frequently present in patients with neurodegenerative diseases, in which patients obviously show abnormalities in the sleep-wake cycle. Patients, especially with AD or PD, often decrease activities during the daytime and increase activities at night, showing changes in rest-activity patterns^[7,8]. Increasing evidence suggests that circadian rhythms are associated with protein homeostasis, which functions in the clearance of abnormal proteins^[17,18]. Disruption of circadian rhythms increases the accumulation of disease proteins, which promotes pathological changes in neurodegenerative diseases.

In this review, we discuss the associations of circadian rhythm disruption and protein homeostasis dysfunction in neurodegenerative diseases. We focused on the influence of protein aggregation and pathological changes by circadian rhythm abnormalities. We summarized the studies on circadian rhythm disruption in AD and PD and the protein homeostasis regulated by circadian rhythms. We also described the pathological significance of circadian rhythm disruption and the underlying mechanisms in neurodegenerative diseases.

CLINICAL ASSOCIATIONS BETWEEN SLEEP DISRUPTION AND NEURODEGENERATIVE DISEASES

Alterations in circadian rhythms can be presented by behavioral and physiological changes, including sleep-wake rhythms, body temperature, blood pressure and hormone levels. The sleep-wake cycle is often disrupted in patients with neurodegenerative diseases. The degeneration of neurons in neurodegenerative diseases is associated with or leads to circadian dysfunction and sleep disturbance, and it was previously believed that the disruption of circadian rhythms is a consequence of neurodegeneration. However, increasing evidence suggests that circadian dysfunction contributes to the formation of pathological changes, such as the accumulation of abnormal proteins and the progression of neurodegenerative diseases [Figure 1]. Importantly, sleep disorders are presented in many neurodegenerative diseases, including early AD, PD, dementia with Lewy bodies, frontotemporal dementia (FTD) and multiple system atrophy (MSA) patients^[8,19-23], further suggesting that they play a role in disease development.

Alzheimer's disease

Sleep disturbance can occur at all stages in AD patients. AD patients lose a normal resting-activity pattern, showing an increase in daytime sleepiness. It has been reported that sleep fragmentation can increase the risk of AD and the rate of cognitive decline^[24]. In a Swedish cohort, 214 Swedish adults aged 75 and over participated in a longitudinal study with 9 years of follow-up. All participants were dementia-free at baseline and the first follow-up (3 years later). After a total of 9 years of follow-up, participants with moderate or severe sleep problems showed an increased risk for developing dementia, particularly AD^[25]. After adjusting for age, education and gender, moderate/severe sleep disruption causes a 2.5 times greater risk of all-cause dementia and a more than 3 times greater risk of AD^[25]. In a prospective study, women without dementia were subjected to overnight polysomnography measurements to determine sleep status. After a 5-year follow-up, their cognitive status was evaluated. Women with sleep-disordered breathing have a high incidence of developing mild cognitive impairment or dementia, suggesting that sleep problems are associated with the development of dementia^[26]. Shift workers are subjected to sleep deprivation, which causes circadian disruption. By analysis of a Danish nurse cohort (28,731 female nurses), shift work increased AD risk^[27]. Moreover, sleep disorder is believed to occur at the preclinical stage of AD^[28]. Using a tailored light treatment that maximally regulates the circadian system, AD patients show significant improvements in sleep, mood and behavior, further suggesting that there is an association between circadian system disruption and the symptoms in AD patients and that an entrainment of circadian rhythms benefits AD patients^[29].

Parkinson's disease

It is estimated that nearly half of dopamine neurons in the substantia nigra are lost when patients start motor symptoms. Before motor symptoms, PD patients often demonstrate nonmotor symptoms, including mood disorders, pain, gastrointestinal dysfunction and sleep disorders. Sleep disturbance, including insomnia, excessive daytime somnolence (EDS), fragmented sleep and rapid eye movement sleep behavior disorder (RBD), is the most common nonmotor symptom, which appears in up to 90% of PD patients^[30-32]. The changes in temporal sleep patterns, including insomnia, EDS and fragmented sleep, that are controlled by circadian rhythms suggest a disruption of the circadian system in PD. Characterized by polysomnography signals, sleep is divided into two states: rapid eye movement (REM) and nonrapid eye movement. Patients with PD at an early stage show increased sleep latency and decreased REM sleep as well as sleep efficiency^[7]. In a study involving 3078 men free of PD and dementia, the risk of developing PD within 10 years was threefold higher in men with EDS than in those without EDS^[33], suggesting that EDS increases the risk of PD and is a prodromal stage of PD. In a population-based prevalence study, RBD was identified as a symptom in the early phase of PD, which occurs in nearly 30% of newly diagnosed PD patients^[34]. The prevalence of RBD in PD patients is estimated to be near 40%, showing that the risk for the



Figure 1. Interactions between sleep disturbance and neurodegenerative diseases. Sleep disturbance plays roles in neurodegenerative disease. It promotes the pathogenesis of Alzheimer's disease (AD) and Parkinson's disease (PD). Sleep disturbance is tightly associated with A β and tau deposition in AD and α -syn accumulation in PD. It also promotes the transmission of disease proteins, such as tau in AD and α -syn in PD. Furthermore, sleep disturbance is positively related to dementia in AD and excessive daytime somnolence (EDS) and rapid eye movement sleep behavior disorder (RBD) in PD. However, there are bidirectional relationships between sleep disturbance and neurodegenerative diseases. Neurodegeneration and pathogenesis in neurodegenerative diseases induce sleep disturbance, further leading to deterioration of the disease. A β : Amyloid- β ; α -syn: α -synuclein.

conversion from idiopathic RBD to PD is extremely high. RBD is associated not only with the development of PD but also with a risk for dementia^[20]. Importantly, among the sleep disorders in PD, RBD is mostly associated with the development of PD pathology. In an observational cohort study, patients with idiopathic RBD were followed up for 7 years and examined with dopamine transport imaging, transcranial sonography and olfactory testing^[35]. With this cohort, 82% of patients with idiopathic RBD show neurodegenerative syndrome with an increased risk for developing PD and Lewy body dementia. In three patients with antemortem diagnosis of PD and Lewy body dementia, there were widespread Lewy bodies in the brains and α -syn aggregates in the peripheral autonomic nervous system, suggesting an association between RBD and synucleinopathy^[35]. Interestingly, these three patients show Lewy pathology as well as a loss of neurons in the brainstem nuclei that regulate REM sleep atonia^[35], which may also reflect a connection between synucleinopathy and REM sleep without atonia in PD.

Frontotemporal dementia

FTD includes a group of heterogeneous dementias with the frontal and temporal lobe atrophy in patients. Behavioral variant FTD (bvFTD) is a common form of FTD syndromes. In comparison to AD patients, FTD patients have more severe daytime somnolence and sleep disturbance^[36]. The sleep disturbance appears earlier in FTD than in AD patients. Moreover, FTD patients have longer sleep onset and less total sleep time than AD patients^[37]. In the prodromal symptoms, sleep disturbance occurs more frequent in bvFTD patients (40%) than in AD patients (12%)^[38].

The hexanucleotide (G4C2) repeat expansion in the chromosome 9 open reading frame 72 (*C9orf72*) gene causes both FTD and amyotrophic lateral sclerosis (ALS)^[39,40]. In a screen for *C9orf72* repeat expansion in 344 RBD patients, two of them have G4C2 expansion, suggesting a possible linkage between RBD and *C9orf72* mutation^[41]. It is well known that dipeptide repeat proteins (DPRs) that are encoded by the expansion of *C9orf72* by non-ATG translation form inclusions in c9FTD/ALS. Interestingly, the abundant DPR inclusions are presented in pineal gland, as well as the supraoptic nucleus and paraventricular nucleus (PVN) that are related to the SCN, implying an association of sleep disruption and c9FTD/ALS^[42].

Multiple system atrophy

The sleep disturbance occurs highly in MSA patients. RBD is a very common symptom in MSA patients. In a meta-analysis, the prevalence of RBD in MSA ranks from 25% to 100%^[43]. In a cross-sectional study in which 165 MSA patients are engaged, sleep disorders are observed in most patients^[44]. RBD occurs in 49.7% of patients, and the frequency of EDS is 27.3%^[44]. Importantly, there is a positive correlation between sleep

disturbances and the severity of MSA^[44]. Another study also shows that there is an association of RBD and MSA, with a frequency of RBD as high as 70.4% in MSA patients^[45]. A cross-sectional study shows that the MSA patients with EDS have a higher score of Non-Motor Symptoms Scale and a higher apnea-hypopnea index as compared the MSA patients without EDS, suggesting that EDS in MSA patients is more associated with sleep-related breathing disorder and other the non-motor symptoms^[46].

PATHOLOGICAL ACCUMULATION OF DISEASE PROTEINS IN ASSOCIATION WITH SLEEP DISRUPTION

The pathological hallmark is the presence of abnormal protein deposition in diseased brains in patients with neurodegenerative diseases. The formation of amyloid plaques by A β and intracellular neurofibrillary tangles by tau in AD or the formation of Lewy bodies by α -syn is a typical pathological process in AD or PD brains^[47]. Soluble A β , tau or α -syn can form oligomers, protofibrils and fibrils, which accumulate and deposit in diseased brains (Soto and Pritzkow^[48], 2018) [Figure 1].

A β

The homeostasis of A β , either accumulation or clearance, is important for AD pathology. A β is a small peptide that is released from amyloid precursor protein (APP) after APP is cleaved by β -secretase and subsequently cleaved by γ -secretase^[49,50]. Mutations in APP or presenilin (catalytic subunit of γ -secretase) that cause early-onset familial AD increase the processing of APP, which promotes the generation of A β peptides^[51,52]. The deposition of A β in AD brains and in APP transgenic mice is a typical pathological feature.

It has been reported that A β has daily oscillations^[53]. The levels of soluble A β in brain interstitial fluid (ISF) in the hippocampus in wild-type or human APP transgenic mice have a pronounced diurnal rhythm^[53]. Moreover, the fluctuation of A β also occurs in the cerebral spinal fluid (CSF) in humans^[53]. In sleep-deprived mice, A β is increased in the ISF in the hippocampus in both wild-type and APP transgenic mice; however, A β is decreased in the ISF in the hippocampus in animals with more sleep^[53]. In *APP^{Swe}/PS1 δ E9* mice, diurnal fluctuation of A β in the ISF and the hippocampus but not in the striatum was attenuated at 6 months of age. Diurnal fluctuation of A β in ISF in the striatum is decreased at 9 months of age when A β plaques appear in the striatum and more often in the hippocampus^[54]. Animals also have significant disturbances in the sleep-wake cycle at 6 months of age^[54]. Thus, the decrease in diurnal fluctuation of A β in the ISF occurs earlier in the hippocampus than in the striatum, and A β plaques appear earlier in the hippocampus than in the striatum, suggesting a link between A β pathological progression and daily oscillations. In addition, active immunization with A β , which decreases A β deposition in *APP^{Swe}/PS1 δ E9* mice, restores the diurnal fluctuation of A β in the ISF and normal sleep-wake cycle^[54], suggesting that A β deposition disrupts A β oscillations and the sleep-wake cycle.

In humans, the daily oscillation of A β in the CSF shows that A β increases during the day, reaches a peak in the evening, and then decreases overnight^[53]. Loss of diurnal fluctuation of the CSF A β occurs in patients with presenilin mutation when A β deposition is detected by amyloid imaging with Pittsburgh Compound B^[54]. An attenuation of diurnal fluctuation of the CSF A β also occurs in patients with presenilin mutation^[54]. Most interestingly, using a radiotracer 18F-florbetaben that binds to A β , images from healthy individuals scanned by positron emission tomography show that even one night of sleep deprivation can increase A β in the hippocampus and thalamus, the regions vulnerable to damage in AD^[55]. This study demonstrates the first evidence, in living humans, that sleep disturbance is directly associated with A β accumulation^[55].

In animal models, *APP^{swe}/PS1^{ΔE9}* transgenic mice show changed sleep architecture compared with control mice. Moreover, sleep disturbance occurs at 4 months of age, but plaque deposition and tau phosphorylation typically occur at 6 months of age^[56], suggesting that sleep changes occur earlier than AD pathology. APP transgenic mice also develop more Aβ plaques in the hippocampus after exposure to sleep deprivation^[53,57]. Chronic sleep deprivation of 8 h per day for 2 months or chronic sleep deprivation of 20 h per day for 21 days increases Aβ plaques in the hippocampus in APP transgenic mice^[53,57]. Sleep deprivation increases Aβ levels in the CSF in humans^[53] and amyloid plaques in the hippocampus in APP transgenic mice^[53,57], suggesting an impact of sleep disorders on Aβ accumulation. However, Aβ has a role in circadian rhythms in multiple transgenic mouse models with Aβ pathology. In *5×FAD* mice that model AD, the circadian rhythms are changed, evidenced by alterations in home cage activity and body temperature^[58]. Thus, there is a bidirectional relationship between Aβ accumulation and sleep disorders^[59].

The association of the accumulation of Aβ with sleep disturbance has been further verified in patients who accept sleep intervention or in animal models that are treated with drugs to improve sleep. In patients with obstructive sleep apnea, the Aβ levels in blood^[60] and the Aβ deposition in brain^[61] are increased. In AD patients with sleep-disordered breathing, a 6-month sleep intervention with a constitutive positive airway pressure ventilation decreases blood Aβ_{42/40} ratio in patients who are completely recovered from sleep disturbance^[62]. Interestingly, there are no changes in blood Aβ_{42/40} ratio in those patients who have no improvement in sleep^[62], suggesting an involvement of sleep in the regulation of Aβ levels. In animal, an administration of nobiletin, a natural compound that is able to enhance the amplitude of clock gene oscillation^[63], improves clock gene expressions and decreases Aβ deposition in APP/PS1 mouse brains^[64].

Tau

Tau is another important protein in association with AD. In the AD brain, tau is hyperphosphorylated and forms intracellular neurofibrillary tangles, a pathological hallmark of Aβ plaques. Tau pathology occurs in the early AD. Aggregates of tau first appear in the brainstem locus coeruleus (LC) and then spread to the transentorhinal and entorhinal regions and subsequently to the hippocampus^[65]. The LC is strongly linked to wakefulness and sends norepinephrine-containing projections to the cortex, amygdala, hippocampus, cerebellum and spinal cord. In tau *P301S* transgenic mice expressing the human P301S tau mutation that is associated with parkinsonism linked to chromosome 17, chronic short sleep at the age of 2-3 months induces deterioration in behaviors and increases in tau oligomers with sustained increases in phosphorylated tau and neuronal loss in the LC and the amygdala^[66]. The brain ISF tau in mice and the CSF tau in humans are also regulated by the sleep-wake cycle, although tau is known as an intracellular protein^[18]. Tau in brain ISF in the hippocampus has diurnal oscillations similar to Aβ. In sleep-deprived mice, tau levels are 2-fold increased in brain ISF^[18]. Interestingly, the increase in tau in brain ISF induced by sleep deprivation can be blocked by terodotoxin, a toxin that blocks voltage-dependent sodium channels, suggesting that neuronal activity is related to tau levels in brain ISF. In addition, sleep deprivation in humans increases CSF Aβ by 30%, while it increases CSF tau by over 50% in the same participants^[18]. With chronic sleep deprivation, the administration of recombinant P301S human tau fibrils does not change tau seeding in the hippocampus; however, it increases tau spreading to the LC, a nucleus associated with wakefulness^[18], suggesting that sleep disorders may induce tau spreading to other brain regions that are associated with the wake-sleep cycle, producing feedback effects on sleep.

α-syn

It has been reported that the severity of hypothalamic Lewy pathology is correlated with Braak stages^[67]. Moreover, Lewy pathology is presented in the SCN with mild or moderate severity in most cases of PD, much fewer in pineal gland^[67]. In multiple system atrophy, no Lewy pathology is present in the SCN or pineal gland^[68], suggesting a tight association between circadian dysfunction and PD pathology. According

to the Braak hypothesis, α -syn pathology can spread from the peripheral autonomic nerve endings of the gastrointestinal tract to the brain^[69]. In a study involving 602 patients with clinical assessment for RBD and neuropathological examination for Lewy-type α -synucleinopathy, Lewy-type α -synucleinopathy occurred in 79.2% of patients with RBD but only 39.5% of those without RBD^[70]. In addition to the brain, α -syn pathology occurs in many organs, including the vagus nerve, gastrointestinal tract, adrenal gland and heart, in PD patients^[71]. The presence of pathological α -syn in the gastrointestinal tract is identified in prodromal PD patients up to 20 years prior to diagnosis^[72]. Moreover, the transmission of α -syn inclusions has been well identified in recent studies that show the development of α -syn pathology in the brain after injecting α -syn preformed fibrils into the duodenum^[73,74]. Thus, peripheral tissues with α -syn pathology may reflect, at least partially, central pathology. In a study using biopsy samples of labial minor salivary glands, α -syn pathology occurred in labial minor salivary glands in 50% of patients with idiopathic RBD and 54% of patients with PD but only 3% of controls^[75]. Moreover, the deposits of α -syn in the parotid gland in idiopathic RBD patients occur at the prodromal stage of PD^[76]. Thus, studies suggest an association of α -syn deposition and RBD.

TDP-43

TDP-43 that is encoded by *TARDBP* is a major pathogenic protein in FTD and ALS^[77]. Although the mutations in *TARDBP* only cover about 5% familial ALS cases and even less in FTD patients^[78]. The TDP-43 pathology is presented in about 97% of ALS patients and 45 % FTD patients^[79]. In ALS patients, the volume of the hypothalamus is decreased^[80]. Moreover, the volume of the PVN in the hypothalamus is also decreased. Furthermore, TDP-43-positive inclusions are observed in the PVN, lateral hypothalamus and fornix^[80]. TDP-43 inclusions are also presented in the reticular formation in the brainstem, which has an important role in the rhythmical cycle of sleep and wakefulness^[81]. The presence of TDP-43 pathology in sleep-related brain regions and nuclei suggests a linkage between the abnormal accumulation of TDP-43 and the symptoms of sleep disorders in ALS and FTD patients.

FACTORS INVOLVED IN CIRCADIAN RHYTHM DISRUPTION IN NEURODEGENERATIVE DISEASES

Clock gene

The molecular basis of circadian rhythms is the oscillation of 24 h clock gene expression in the SCN^[1]. The core clock gene products circadian locomotor output cycles kaput (CLOCK) and brain and muscle arnt-like 1 (BMAL1) form heterodimers that bind to E-boxes and drive the expression of period (PERs) and cryptochrome (CRYs). The expression of PERs and CRYs in turn represses CLOCK-BMAL1 activity, thus inhibiting their own expression, which forms a feedback loop that takes 24 h^[1]. Thus, the molecules involved in the molecular clock are important for the maintenance of circadian rhythms.

It has been reported that the alteration of DNA methylation of BMAL1 in early AD patients causes changes in BMAL1 expression patterns in both amplitude and phase^[82]. Knockout of *Bmal1* disrupts A β oscillation in brain ISF in the hippocampus. Moreover, in tamoxifen-inducible global *Bmal1* knockout mice with an APP transgenic background, *Bmal1* knockout significantly increases hippocampal A β plaques 4 months after tamoxifen induction^[17]. In AD patients, there is a disruption of circadian rhythms, typically showing sleep problems. Sleep deprivation also affects clock gene expression and the DNA binding patterns of BMAL1 and CLOCK heterodimers, which disturbs clock function. Three different polymorphisms of the CLOCK gene are associated with AD in the Chinese population^[83,84]. Moreover, changes in expression patterns and decreases in expression levels of *Bmal1* and *Clock* in senescent cells and aged rodent brains suggest a role of circadian genes in aging^[84]. In 5 \times FAD mice, the amplitude of BMAL1 and PER2 in a 24 h oscillation is greatly decreased^[58]. Moreover, A β is able to induce BMAL1 degradation *in vitro*^[58]. In

APP/PS1 mice, there is also a modest alteration of PER2 in the SCN, which is consistent with the findings in *5×FAD* mice^[58,85]. In addition, at circadian time 4 (CT4, 4 h after the onset of activity of diurnal organisms, based on the free-running period of a rhythm), when microglia express higher levels of BMAL1 than at CT12, the clearance ability for fibrillary A β by microglia at CT4 is also higher than that at CT12 in *5×FAD* mice^[86]. Pharmacological inhibition of REV-ERBs that is transactivated by BMAL1 and negative feedback on BMAL1 activity promote a microglial M2-like phenotype. Moreover, a constitutive deletion of *Rev-erb α* in *5×FAD* mice can repress amyloid plaque formation^[86], further suggesting a role of the clock gene in protein homeostasis.

It has been reported that BMAL1 oscillation amplitude is decreased in PD patients and that BMAL1 mRNA levels are decreased in leukocytes in PD patients, suggesting that dysfunction of the clock gene *Bmal1* may be associated with PD.

Melatonin

In the brain, circadian information in the SCN can be sent to different areas. Through multisynaptic projections, the SCN circadian information is integrated and sent to the superior cervical ganglia (SCG). The noradrenergic neurons in the SCG send projections to the pineal gland^[87]. The pineal gland produces melatonin, the hormone that synchronizes and stabilizes circadian rhythms, which is important for the maintenance of the biological clock of the brain in the SCN. Melatonin is synthesized in the dark and shows oscillation at its level, which is controlled by the SCN [Figure 2]. The secretion of melatonin starts at early night and usually reaches a peak level at 3:00 to 4:00 am. With aging or neurodegenerative diseases, the circadian amplitude is decreased with desynchronization of physiological rhythms, leading to a decrease in melatonin levels^[88]. Melatonin binds to melatonin receptor 1 (MT1) and MT2, which are G-protein coupled receptors and are highly expressed in neurons in the SCN, hippocampus, thalamus, vestibular nuclei and cerebral and cerebellar cortex^[89]. In addition to the maintenance of circadian rhythms, melatonin has multiple functions, including antioxidative stress and regulating metabolism^[88].

AD patients have lower melatonin levels than normal controls. The melatonin level is decreased in the CSF in early AD patients before clinical symptoms^[90]. Meanwhile, the loss of neurons in the SCN further shows a correlation between AD and circadian rhythm dysfunction^[91]. In patients with AD at the early stage, the initial evening secretion of melatonin is delayed and mildly decreased, suggesting that a circadian phase shift occurs in early AD patients^[92]. Moreover, the risk variant rs12506228, which is located downstream of *MTNR1A* (MT1A-encoding gene), is associated with AD^[93]. The rs12506228 variant leads to a decrease in MT1A expression and is associated with both clinical and pathological changes in AD patients^[93], further suggesting a role of melatonin in AD.

It has been reported that melatonin has effects on A β deposits in AD. In *APP/PS1* mice, long-term melatonin treatment at ages starting from 2-2.5 months decreases A β deposits in the hippocampus and the entorhinal cortex when the animals are examined at the age of 7.5 months^[94]. Meanwhile, there was a decrease in inflammatory cytokines in the hippocampus in *APP/PS1* mice treated with melatonin, suggesting a role of melatonin in anti-inflammation and decreasing A β accumulation^[94]. It has also been reported that melatonin can induce lymphatic clearance of A β in *Tg2576* mice^[95].

Using real-time PCR analysis, it has been identified that MT1 and MT2 are decreased in both the substantia nigra and the amygdala in PD patients compared to controls^[96]. In PD patients, circulating melatonin is decreased in early PD patients compared with controls^[7]. Moreover, plasma melatonin, both the amplitude of daily oscillation and the levels, is lower in PD patients than in controls^[97]. In PD patients who accept




Figure 2. Impact of circadian genes and hormones on neurodegeneration and the glymphatic system. The secretion of the circadian hormone melatonin is controlled by circadian rhythm. In addition to functioning in circadian rhythm, melatonin has protective effects on neurons. In neurodegenerative diseases, the levels of melatonin are decreased.

melatonin treatment at a dose of 5 mg or a high dose of 50 mg 30 min before bedtime over 14 days, the nocturnal sleep that is evaluated with actigraphy is significantly improved in PD patients during melatonin treatment at both doses^[98]. Interestingly, the treatment of PD patients with levodopa increases melatonin levels^[99].

In a rotenone PD animal model^[100] or in a 6-hydroxydopamine with or without pinealectomy animal model^[101], melatonin protects DA neurons. Melatonin also improves rotenone-induced defects in behaviors, including grip strength and performance on rotarods^[102]. In *MPTP* mice, melatonin protects DA neurons against MPTP-induced neurotoxicity, which may be mediated by antioxidant effects and neuroprotective effects of melatonin^[103-105]. In *MPTP* mice that receive both melatonin and levodopa treatments, melatonin increases the therapeutic effects of levodopa in the improvement of MPTP-induced akinesia and catalepsy^[105]. In a lentivirus-infected animal model that expresses A30P pathogenic α -syn, melatonin administration protects DA neurons against A30P α -syn-induced TH neuronal loss^[106]. In amphetamine-treated 4-day-old postnatal rats, melatonin decreases amphetamine-induced α -syn accumulation in the substantia nigra, dorsal striatum, nucleus accumbens, and prefrontal cortex^[107]. Thus, melatonin has multiple effects on circadian rhythm regulation, neuroprotection and protein clearance, which can be associated with the pathogenesis of neurodegenerative diseases [Figure 2]

GLYMPHATIC PATHWAY

Although the central nervous system is thought to anatomically lack lymphatic vessels for the removal of interstitial metabolic waste products, it has been recently discovered that glymphatic (glial-lymphatic)^[108] and meningeal lymphatic vessels^[109,110] execute functions to transport and drain brain wastes, such as the peripheral lymphatic system. The glymphatic system is a glial-dependent perivascular network that has lymphatic functions^[111]. The CSF produced by the choroid plexuses flows into the brain along periaxillary spaces surrounding cerebral arteries and arterioles, running in the same direction as blood flow, which is driven by arterial pulsation^[112]. The CSF enters the brain parenchyma and mixes with ISF, which is facilitated by the aquaporin (AQP-4) water channel on the perivascular end-foot processes of astrocytes^[113]. Mixtures of CSF and ISF with interstitial solutes outflow along the perivenous space and drain out of the brain via meningeal lymphatic vessels or along cranial and spinal nerves^[113] [Figure 3]. The meningeal




Figure 3. The glymphatic system in association with the clearance of neurodegenerative disease proteins. The glymphatic system is a perivascular network that has lymphatic functions. Driven by arterial pulsation, the CSF flows into the brain along periarterial spaces surrounding cerebral arteries and arterioles, running in the same direction as blood flow. The CSF enters the brain parenchyma mediated by the aquaporin (AQP-4) water channel on the perivascular end-foot processes of astrocytes. Mixtures of CSF and ISF with interstitial solutes outflow along the perivenous space and drain out of the brain, which can clear Aβ in the ISF. The glymphatic system depends on AQP-4 on astrocytes. Sleep enhances glymphatic system function to promote the clearance of Aβ and other disease proteins in the brain. CSF: Cerebral spinal fluid; ISF: interstitial fluid; Aβ: amyloid-β.

lymphatic vessels are located in the dura matter^[109,110]. The metabolites in the CSF are cleared from the brain by fluid flow and finally drained to extracranial deep cervical lymph nodes^[114]. Dysfunction of the glymphatic system may be a final common pathway to dementia^[115]. Using gadobutrol as the CSF tracer with intrathecal administration, the glymphatic system assessed with magnetic resonance imaging demonstrates delayed glymphatic clearance of gadobutrol from the subarachnoid space, with an enhancement of the signal in the brain parenchyma, in idiopathic normal hydrocephalus, which is the first study in humans showing glymphatic clearance of a CSF tracer (solute)^[116].

The glymphatic system is tightly associated with the clearance of neurodegenerative disease proteins, such as Aβ^[108,113]. In PD patients, the glymphatic system is impaired, and in animals, blockage of meningeal lymphatic vessels increases α-syn preformed fibril-induced α-syn pathology^[117]. Interestingly, the activity of the brain glymphatic system is controlled by circadian rhythms^[118], which are related to the clearance of neurodegenerative disease proteins^[119]. Glymphatic clearance is facilitated by AQP-4, which influences CSF influx into the brain parenchyma where it mixes with ISF^[108]. In *Aqp4* (AQP-4 encoding gene) knockout mice, CSF tracer influx into the brain parenchyma is markedly reduced; however, the periarterial movement of the tracer is not significantly decreased^[108]. Moreover, in *Aqp4* knockout mice, a 55% reduction in the clearance of Aβ occurs compared with wild-type mice when ¹²⁵I-Aβ₄₂ is infused into the striatum^[108]. The interstitial space in sleeping mice is larger than that in awake mice^[119], which increases the convective ISF

bulk flow in the brain parenchyma to increase glymphatic clearance of A β ^[119] [Figure 3]. In *APP/PS1* mice, A β also shows an impact on the glymphatic pathway, leading to dysfunction of the glymphatic pathway^[120]. The inflow of A β 40 into the brain and the clearance of A β 40 by the glymphatic pathway in the brain are decreased in *APP/PS1* mice (Peng *et al.*^[120], 2016). Disruption of meningeal lymphatic vessels in 5 \times *FAD* mice also aggravates A β deposition in the meninges and A β accumulation in the brain parenchyma^[121]. Most recently, it has been reported that there is a circadian rhythm in glymphatic influx, and loss of AQP-4 eliminates circadian CSF distribution, further suggesting a linkage between circadian rhythms and glymphatic clearance^[118]. Interestingly, the localization of AQP-4 to the endfeet of astrocytes surrounding the vasculature has diurnal variation, showing an increase in AQP-4 polarization surrounding the vasculature during the day^[118]. Correspondingly, glymphatic influx and the clearance of solutes are increased during the day. In addition, the influx that is indicated by a CSF tracer shows an increase during the rest phase compared with the active phase of animal behavior, further suggesting a role of circadian rhythms in glymphatic influx^[118]. There are also some controversial studies. In *Aqp4* knockout mice and rats, no significant difference in tracer (Alexa 647-labeled ovalbumin) penetration into the striatal parenchyma from the paravascular space was observed^[122]. The mechanisms for the clearance of waste by the glymphatic pathway are still largely unknown. Furthermore, the linkage between circadian rhythms and the glymphatic pathway is still being identified.

CONCLUSION

It is clear that there is a link between circadian disruption and neurodegenerative diseases. Sleep problems often start at the early stage in patients with neurodegenerative diseases. Clinical studies and animal models have revealed an association between the pathological progression of neurodegenerative diseases and circadian disruption. It has been well documented that sleep deprivation aggravates the deposition of neurodegenerative disease proteins in animal models. It is also known that sleep increases the glymphatic clearance of A β and tau in the CSF. However, our understanding of the role of circadian rhythms in protein homeostasis and disease development is very limited. There are still some key questions that need to be addressed: (1) whether circadian rhythms influence protein quality control systems, such as the ubiquitin-proteasomal pathway or autophagic pathway, facilitating disease protein degradation; (2) why RBD, within different sleep disturbances, is the highest risk factor associated with synucleinopathy, PD and dementia; (3) how the phenotypes of AQP-4 deficiency can be linked to the phenotypes of neurodegenerative diseases in animal models; and (4) whether there are different or the same mechanisms accounting for the interactions between abnormal accumulation of disease proteins and the dysfunction of circadian rhythms in patients with different neurodegenerative diseases. Further studies should address how circadian rhythm affects the clearance or accumulation of disease proteins and by which mechanism the abnormal proteins lead to dysfunction of circadian, which may guide the development of novel strategies for clinical treatment and drug targets for neurodegenerative diseases.

DECLARATIONS

Authors' contributions

Conceived the idea of this review article: Li J

Drafted the manuscript: Wang X

Revised the manuscript: Li J

Designed pictures: Wang R, Wang X

Completed the pictures: Wang R

Available of data and materials

Not applicable.

Financial support and sponsorship

This work was supported by the National Natural Science Foundation of China (31972913), Key Research and Development Programs from Hunan Province (2018DK2010 and 2018DK2013), Guangdong Key Project in Development of new tools for diagnosis and treatment of Autism (2018B030335001) and National Undergraduate Training Program for Innovation (No. 201910533108).

Conflicts of interest

All authors declared that there are no conflicts of interest.

Ethical approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Copyright

© The Author(s) 2022.

REFERENCES

1. Patke A, Young MW, Axelrod S. Molecular mechanisms and physiological importance of circadian rhythms. *Nat Rev Mol Cell Biol* 2020;21:67-84. DOI PubMed
2. Hastings MH, Maywood ES, Brancaccio M. Generation of circadian rhythms in the suprachiasmatic nucleus. *Nat Rev Neurosci* 2018;19:453-69. DOI PubMed
3. Ralph MR, Foster RG, Davis FC, Menaker M. Transplanted suprachiasmatic nucleus determines circadian period. *Science* 1990;247:975-8. DOI PubMed
4. Bloch KE, Brack T, Wirz-Justice A. Transient short free running circadian rhythm in a case of aneurysm near the suprachiasmatic nuclei. *J Neurol Neurosurg Psychiatry* 2005;76:1178-80. DOI PubMed PMC
5. Borgers AJ, Romeijn N, van Someren E, Fliers E, Alkemade A, Bisschop PH. Compression of the optic chiasm is associated with permanent shorter sleep duration in patients with pituitary insufficiency. *Clin Endocrinol (Oxf)* 2011;75:347-53. DOI PubMed
6. Mohawk JA, Green CB, Takahashi JS. Central and peripheral circadian clocks in mammals. *Annu Rev Neurosci* 2012;35:445-62. DOI PubMed PMC
7. Breen DP, Vuono R, Nawarathna U, et al. Sleep and circadian rhythm regulation in early Parkinson disease. *JAMA Neurol* 2014;71:589-95. DOI PubMed PMC
8. Musiek ES, Bhimasani M, Zangrilli MA, Morris JC, Holtzman DM, Ju YS. Circadian rest-activity pattern changes in aging and preclinical Alzheimer disease. *JAMA Neurol* 2018;75:582-90. DOI PubMed PMC
9. Masri S, Sassone-Corsi P. The emerging link between cancer, metabolism, and circadian rhythms. *Nat Med* 2018;24:1795-803. DOI PubMed PMC
10. Douma LG, Gumz ML. Circadian clock-mediated regulation of blood pressure. *Free Radic Biol Med* 2018;119:108-14. DOI PubMed PMC
11. Mason IC, Qian J, Adler GK, Scheer FAJL. Impact of circadian disruption on glucose metabolism: implications for type 2 diabetes. *Diabetologia* 2020;63:462-72. DOI PubMed PMC
12. Man K, Loudon A, Chawla A. Immunity around the clock. *Science* 2016;354:999-1003. DOI PubMed PMC
13. Foster RG. Sleep, circadian rhythms and health. *Interface Focus* 2020;10:20190098. DOI PubMed PMC
14. Taylor JP, Hardy J, Fischbeck KH. Toxic proteins in neurodegenerative disease. *Science* 2002;296:1991-5. DOI PubMed
15. Zhang S, Wang R, Wang G. Impact of dopamine oxidation on dopaminergic neurodegeneration. *ACS Chem Neurosci* 2019;10:945-53. DOI PubMed
16. Wang R, Sun H, Ren H, Wang G. α -Synuclein aggregation and transmission in Parkinson's disease: a link to mitochondria and lysosome. *Sci China Life Sci* 2020;63:1850-9. DOI PubMed
17. Kress GJ, Liao F, Dimitry J, et al. Regulation of amyloid- β dynamics and pathology by the circadian clock. *J Exp Med* 2018;215:1059-68. DOI PubMed PMC
18. Holth JK, Fritsch SK, Wang C, et al. The sleep-wake cycle regulates brain interstitial fluid tau in mice and CSF tau in humans.

- Science* 2019;363:880-4. DOI PubMed PMC
19. Sinforiani E, Terzaghi M, Pasotti C, Zucchella C, Zambrelli E, Manni R. Hallucinations and sleep-wake cycle in Alzheimer's disease: a questionnaire-based study in 218 patients. *Neurol Sci* 2007;28:96-9. DOI PubMed
 20. Li S, Wang Y, Wang F, Hu LF, Liu CF. A new perspective for Parkinson's disease: circadian rhythm. *Neurosci Bull* 2017;33:62-72. DOI PubMed PMC
 21. He C, Hu Z, Jiang C. Sleep disturbance: an early sign of Alzheimer's disease. *Neurosci Bull* 2020;36:449-51. DOI PubMed PMC
 22. McCarter SJ, St Louis EK, Boeve BF. Sleep disturbances in frontotemporal dementia. *Curr Neurol Neurosci Rep* 2016;16:85. DOI PubMed
 23. Fife K, De Boer T. The circadian system in Parkinson's disease, multiple system atrophy, and progressive supranuclear palsy. *Handb Clin Neurol* 2021;179:301-13. DOI PubMed
 24. Lim AS, Kowgier M, Yu L, Buchman AS, Bennett DA. Sleep fragmentation and the risk of incident Alzheimer's disease and cognitive decline in older persons. *Sleep* 2013;36:1027-32. DOI PubMed PMC
 25. Hahn EA, Wang HX, Andel R, Fratiglioni L. A change in sleep pattern may predict Alzheimer disease. *Am J Geriatr Psychiatry* 2014;22:1262-71. DOI PubMed
 26. Yaffe K, Laffan AM, Harrison SL, et al. Sleep-disordered breathing, hypoxia, and risk of mild cognitive impairment and dementia in older women. *JAMA* 2011;306:613-9. DOI PubMed PMC
 27. Jørgensen JT, Karlsen S, Stayner L, Hansen J, Andersen ZJ. Shift work and overall and cause-specific mortality in the Danish nurse cohort. *Scand J Work Environ Health* 2017;43:117-26. DOI PubMed
 28. Liguori C, Mercuri NB, Izzi F, et al. Obstructive sleep apnea is associated with early but possibly modifiable Alzheimer's disease biomarkers changes. *Sleep* 2017. DOI PubMed
 29. Figueiro MG, Plitnick B, Roohan C, Sahin L, Kalsher M, Rea MS. Effects of a tailored lighting intervention on sleep quality, rest-activity, mood, and behavior in older adults with Alzheimer disease and related dementias: a randomized clinical trial. *J Clin Sleep Med* 2019;15:1757-67. DOI PubMed PMC
 30. Shen Y, Huang JY, Li J, Liu CF. Excessive daytime sleepiness in Parkinson's disease: clinical implications and management. *Chin Med J (Engl)* 2018;131:974-81. DOI PubMed PMC
 31. Li T, Le W. Biomarkers for Parkinson's disease: how good are they? *Neurosci Bull* 2020;36:183-94. DOI PubMed PMC
 32. Zhang F, Niu L, Liu X, et al. Rapid eye movement sleep behavior disorder and neurodegenerative diseases: an update. *Aging Dis* 2020;11:315-26. DOI PubMed PMC
 33. Abbott RD, Ross GW, White LR, et al. Excessive daytime sleepiness and subsequent development of Parkinson disease. *Neurology* 2005;65:1442-6. DOI PubMed
 34. Gjerstad MD, Boeve B, Wentzel-Larsen T, Aarsland D, Larsen JP. Occurrence and clinical correlates of REM sleep behaviour disorder in patients with Parkinson's disease over time. *J Neurol Neurosurg Psychiatry* 2008;79:387-91. DOI PubMed
 35. Iranzo A, Tolosa E, Gelpi E, et al. Neurodegenerative disease status and post-mortem pathology in idiopathic rapid-eye-movement sleep behaviour disorder: an observational cohort study. *Lancet Neurol* 2013;12:443-53. DOI PubMed
 36. Sani TP, Bond RL, Marshall CR, et al. Sleep symptoms in syndromes of frontotemporal dementia and Alzheimer's disease: a proof-of-principle behavioural study. *eNeurologicalSci* 2019;17:100212. DOI PubMed PMC
 37. Bonakis A, Economou NT, Paparrigopoulos T, et al. Sleep in frontotemporal dementia is equally or possibly more disrupted, and at an earlier stage, when compared to sleep in Alzheimer's disease. *J Alzheimers Dis* 2014;38:85-91. DOI PubMed
 38. Korhonen T, Katisko K, Cajanus A, et al. Comparison of prodromal symptoms of patients with behavioral variant frontotemporal dementia and Alzheimer disease. *Dement Geriatr Cogn Disord* 2020;49:98-106. DOI PubMed
 39. DeJesus-Hernandez M, Mackenzie IR, Boeve BF, et al. Expanded GGGGCC hexanucleotide repeat in noncoding region of C9ORF72 causes chromosome 9p-linked FTD and ALS. *Neuron* 2011;72:245-56. DOI PubMed PMC
 40. Renton AE, Majounie E, Waite A, et al; ITALSGEN Consortium. A hexanucleotide repeat expansion in C9ORF72 is the cause of chromosome 9p21-linked ALS-FTD. *Neuron* 2011;72:257-68. DOI PubMed PMC
 41. Daoud H, Postuma RB, Bourassa CV, et al. C9orf72 repeat expansions in rapid eye movement sleep behaviour disorder. *Can J Neurol Sci* 2014;41:759-62. DOI PubMed
 42. Dedeene L, Van Schoor E, Vandenberghe R, Van Damme P, Poesen K, Thal DR. Circadian sleep/wake-associated cells show dipeptide repeat protein aggregates in C9orf72-related ALS and FTLD cases. *Acta Neuropathol Commun* 2019;7:189. DOI PubMed PMC
 43. Palma JA, Fernandez-Cordon C, Coon EA, et al. Prevalence of REM sleep behavior disorder in multiple system atrophy: a multicenter study and meta-analysis. *Clin Auton Res* 2015;25:69-75. DOI PubMed PMC
 44. Lin JY, Zhang LY, Cao B, et al. Sleep-related symptoms in multiple system atrophy: determinants and impact on disease severity. *Chin Med J (Engl)* 2020;134:690-8. DOI PubMed PMC
 45. Wu DD, Su W, Li SH, et al. A questionnaire-based study on clinical REM sleep behavior disorder and subtypes in multiple system atrophy. *Eur Neurol* 2021;84:368-74. DOI PubMed
 46. Wang H, Tang X, Zhou J, Xu Y. Excessive daytime sleepiness is associated with non-motor symptoms of multiple system atrophy: a cross-sectional study in China. *Front Neurol* 2021;12:798771. DOI PubMed PMC
 47. Soto C. Unfolding the role of protein misfolding in neurodegenerative diseases. *Nat Rev Neurosci* 2003;4:49-60. DOI PubMed
 48. Soto C, Pritzke S. Protein misfolding, aggregation, and conformational strains in neurodegenerative diseases. *Nat Neurosci*

- 2018;21:1332-40. DOI PubMed PMC
49. Zhao J, Liu X, Xia W, Zhang Y, Wang C. Targeting amyloidogenic processing of APP in Alzheimer's disease. *Front Mol Neurosci* 2020;13:137. DOI PubMed PMC
 50. Wang X. A bridge between the innate immunity system and amyloid- β production in Alzheimer's disease. *Neurosci Bull* 2021;37:898-901. DOI PubMed PMC
 51. De Jonghe C, Esselens C, Kumar-Singh S, et al. Pathogenic APP mutations near the gamma-secretase cleavage site differentially affect Abeta secretion and APP C-terminal fragment stability. *Hum Mol Genet* 2001;10:1665-71. DOI PubMed
 52. Selkoe DJ, Wolfe MS. Presenilin: running with scissors in the membrane. *Cell* 2007;131:215-21. DOI PubMed
 53. Kang JE, Lim MM, Bateman RJ, et al. Amyloid-beta dynamics are regulated by orexin and the sleep-wake cycle. *Science* 2009;326:1005-7. DOI PubMed PMC
 54. Roh JH, Huang Y, Bero AW, et al. Disruption of the sleep-wake cycle and diurnal fluctuation of β -amyloid in mice with Alzheimer's disease pathology. *Sci Transl Med* 2012;4:150ra122. DOI PubMed PMC
 55. Shokri-Kojori E, Wang GJ, Wiers CE, et al. β -Amyloid accumulation in the human brain after one night of sleep deprivation. *Proc Natl Acad Sci U S A* 2018;115:4483-8. DOI PubMed PMC
 56. Zhang F, Zhong R, Li S, et al. Alteration in sleep architecture and electroencephalogram as an early sign of Alzheimer's disease preceding the disease pathology and cognitive decline. *Alzheimers Dement* 2019;15:590-7. DOI PubMed
 57. Qiu H, Zhong R, Liu H, Zhang F, Li S, Le W. Chronic sleep deprivation exacerbates learning-memory disability and Alzheimer's disease-like pathologies in A β PP(swe)/PS1(Δ E9) mice. *J Alzheimers Dis* 2016;50:669-85. DOI PubMed
 58. Song H, Moon M, Choe HK, et al. A β -induced degradation of BMAL1 and CBP leads to circadian rhythm disruption in Alzheimer's disease. *Mol Neurodegener* 2015;10:13. DOI PubMed PMC
 59. Dufort-Gervais J, Mongrain V, Brouillette J. Bidirectional relationships between sleep and amyloid-beta in the hippocampus. *Neurobiol Learn Mem* 2019;160:108-17. DOI PubMed
 60. Bhuniya S, Goyal M, Chowdhury N, Mishra P. Intermittent hypoxia and sleep disruption in obstructive sleep apnea increase serum tau and amyloid-beta levels. *J Sleep Res* 2022. DOI PubMed
 61. Yun CH, Lee HY, Lee SK, et al. Amyloid burden in obstructive sleep apnea. *J Alzheimers Dis* 2017;59:21-9. DOI PubMed
 62. Huang H, Li M, Zhang M, et al. Sleep quality improvement enhances neuropsychological recovery and reduces blood A $\beta_{42/40}$ ratio in patients with mild-moderate cognitive impairment. *Medicina (Kaunas)* 2021;57:1366. DOI PubMed
 63. He B, Nohara K, Park N, et al. The small molecule nobiletin targets the molecular oscillator to enhance circadian rhythms and protect against metabolic syndrome. *Cell Metab* 2016;23:610-21. DOI PubMed PMC
 64. Kim E, Nohara K, Wirianto M, et al. Effects of the clock modulator nobiletin on circadian rhythms and pathophysiology in female mice of an Alzheimer's disease model. *Biomolecules* 2021;11:1004. DOI PubMed PMC
 65. Brettschneider J, Del Tredici K, Lee VM, Trojanowski JQ. Spreading of pathology in neurodegenerative diseases: a focus on human studies. *Nat Rev Neurosci* 2015;16:109-20. DOI PubMed PMC
 66. Zhu Y, Zhan G, Fenik P, et al. Chronic sleep disruption advances the temporal progression of tauopathy in P301S mutant mice. *J Neurosci* 2018;38:10255-70. DOI PubMed PMC
 67. Pablo-Fernandez E, Courtney R, Holton JL, Warner TT. Hypothalamic α -synuclein and its relation to weight loss and autonomic symptoms in Parkinson's disease. *Mov Disord* 2017;32:296-8. DOI PubMed
 68. Pablo-Fernández E, Courtney R, Warner TT, Holton JL. A histologic study of the circadian system in Parkinson disease, multiple system atrophy, and progressive supranuclear palsy. *JAMA Neurol* 2018;75:1008-12. DOI PubMed PMC
 69. Braak H, Rüb U, Gai WP, Del Tredici K. Idiopathic Parkinson's disease: possible routes by which vulnerable neuronal types may be subject to neuroinvasion by an unknown pathogen. *J Neural Transm (Vienna)* 2003;110:517-36. DOI PubMed
 70. Shprecher DR, Adler CH, Zhang N, et al. Predicting alpha-synuclein pathology by REM sleep behavior disorder diagnosis. *Parkinsonism Relat Disord* 2018;55:92-6. DOI PubMed PMC
 71. Gelpi E, Navarro-Otano J, Tolosa E, et al. Multiple organ involvement by alpha-synuclein pathology in Lewy body disorders. *Mov Disord* 2014;29:1010-8. DOI PubMed
 72. Stokholm MG, Danielsen EH, Hamilton-Dutoit SJ, Borghammer P. Pathological α -synuclein in gastrointestinal tissues from prodromal Parkinson disease patients. *Ann Neurol* 2016;79:940-9. DOI PubMed
 73. Kim S, Kwon SH, Kam TI, et al. Transneuronal propagation of pathologic α -synuclein from the gut to the brain models Parkinson's disease. *Neuron* 2019;103:627-41.e7. DOI PubMed PMC
 74. Van Den Berge N, Ferreira N, Gram H, et al. Evidence for bidirectional and trans-synaptic parasympathetic and sympathetic propagation of alpha-synuclein in rats. *Acta Neuropathol* 2019;138:535-50. DOI PubMed PMC
 75. Iranzo A, Borrego S, Vilaseca I, et al. α -Synuclein aggregates in labial salivary glands of idiopathic rapid eye movement sleep behavior disorder. *Sleep* 2018. DOI PubMed
 76. Fernández-Arcos A, Vilaseca I, Aldecoa I, et al. Alpha-synuclein aggregates in the parotid gland of idiopathic REM sleep behavior disorder. *Sleep Med* 2018;52:14-7. DOI PubMed
 77. Forman MS, Trojanowski JQ, Lee VM. TDP-43: a novel neurodegenerative proteinopathy. *Curr Opin Neurobiol* 2007;17:548-55. DOI PubMed PMC
 78. Kabashi E, Valdmanis PN, Dion P, et al. TARDBP mutations in individuals with sporadic and familial amyotrophic lateral sclerosis. *Nat Genet* 2008;40:572-4. DOI PubMed

79. Ling SC, Polymenidou M, Cleveland DW. Converging mechanisms in ALS and FTD: disrupted RNA and protein homeostasis. *Neuron* 2013;79:416-38. DOI PubMed PMC
80. Gabery S, Ahmed RM, Caga J, Kiernan MC, Halliday GM, Petersén Å. Loss of the metabolism and sleep regulating neuronal populations expressing orexin and oxytocin in the hypothalamus in amyotrophic lateral sclerosis. *Neuropathol Appl Neurobiol* 2021;47:979-89. DOI PubMed
81. Calderón-Garcidueñas L, Rajkumar RP, Stommel EW, et al. Brainstem quadruple aberrant hyperphosphorylated tau, beta-amyloid, alpha-synuclein and TDP-43 pathology, stress and sleep behavior disorders. *Int J Environ Res Public Health* 2021;18:6689. DOI PubMed PMC
82. Cronin P, McCarthy MJ, Lim ASP, et al. Circadian alterations during early stages of Alzheimer's disease are associated with aberrant cycles of DNA methylation in BMAL1. *Alzheimers Dement* 2017;13:689-700. DOI PubMed PMC
83. Chen HF, Huang CQ, You C, Wang ZR, Si-qing H. Polymorphism of CLOCK gene rs 4580704 C > G is associated with susceptibility of Alzheimer's disease in a Chinese population. *Arch Med Res* 2013;44:203-7. DOI PubMed
84. Musiek ES, Xiong DD, Holtzman DM. Sleep, circadian rhythms, and the pathogenesis of Alzheimer disease. *Exp Mol Med* 2015;47:e148. DOI PubMed PMC
85. Duncan MJ, Smith JT, Franklin KM, et al. Effects of aging and genotype on circadian rhythms, sleep, and clock gene expression in APPxPS1 knock-in mice, a model for Alzheimer's disease. *Exp Neurol* 2012;236:249-58. DOI PubMed
86. Lee J, Kim DE, Griffin P, et al. Inhibition of REV-ERBs stimulates microglial amyloid-beta clearance and reduces amyloid plaque deposition in the 5XFAD mouse model of Alzheimer's disease. *Aging Cell* 2020;19:e13078. DOI PubMed PMC
87. Bodinat C, Guardiola-Lemaitre B, Mocaër E, Renard P, Muñoz C, Millan MJ. Agomelatine, the first melatonergic antidepressant: discovery, characterization and development. *Nat Rev Drug Discov* 2010;9:628-42. DOI PubMed
88. Gunata M, Parlakpinar H, Acet HA. Melatonin: a review of its potential functions and effects on neurological diseases. *Rev Neurol (Paris)* 2020;176:148-65. DOI PubMed
89. Cardinali DP. Melatonin: clinical perspectives in neurodegeneration. *Front Endocrinol (Lausanne)* 2019;10:480. DOI PubMed PMC
90. Zhou JN, Liu RY, Kamphorst W, Hofman MA, Swaab DF. Early neuropathological Alzheimer's changes in aged individuals are accompanied by decreased cerebrospinal fluid melatonin levels. *J Pineal Res* 2003;35:125-30. DOI PubMed
91. Stopa EG, Volicer L, Kuo-Leblanc V, et al. Pathologic evaluation of the human suprachiasmatic nucleus in severe dementia. *J Neuropathol Exp Neurol* 1999;58:29-39. DOI PubMed
92. Manni R, Cremascoli R, Perretti C, et al. Evening melatonin timing secretion in real life conditions in patients with Alzheimer disease of mild to moderate severity. *Sleep Med* 2019;63:122-6. DOI PubMed
93. Sulkava S, Muggalla P, Sulkava R, et al. Melatonin receptor type 1A gene linked to Alzheimer's disease in old age. *Sleep* 2018;41:zsy103. DOI PubMed PMC
94. Olcese JM, Cao C, Mori T, et al. Protection against cognitive deficits and markers of neurodegeneration by long-term oral administration of melatonin in a transgenic model of Alzheimer disease. *J Pineal Res* 2009;47:82-96. DOI PubMed
95. Pappolla MA, Matsubara E, Vidal R, et al. Melatonin treatment enhances Aβ lymphatic clearance in a transgenic mouse model of amyloidosis. *Curr Alzheimer Res* 2018;15:637-42. DOI PubMed PMC
96. Adi N, Mash DC, Ali Y, Singer C, Shehadeh L, Papapetropoulos S. Melatonin MT1 and MT2 receptor expression in Parkinson's disease. *Med Sci Monit* 2010;16:BR61-7. PubMed
97. Videnovic A, Noble C, Reid KJ, et al. Circadian melatonin rhythm and excessive daytime sleepiness in Parkinson disease. *JAMA Neurol* 2014;71:463-9. DOI PubMed PMC
98. Dowling GA, Mastick J, Colling E, Carter JH, Singer CM, Aminoff MJ. Melatonin for sleep disturbances in Parkinson's disease. *Sleep Med* 2005;6:459-66. DOI PubMed
99. Bordet R, Devos D, Brique S, et al. Study of circadian melatonin secretion pattern at different stages of Parkinson's disease. *Clin Neuropharmacol* 2003;26:65-72. DOI PubMed
100. Carriere CH, Kang NH, Niles LP. Chronic low-dose melatonin treatment maintains nigrostriatal integrity in an intrastriatal rotenone model of Parkinson's disease. *Brain Res* 2016;1633:115-25. DOI PubMed
101. Li Y, Wang SM, Guo L, et al. Effects of melatonin levels on neurotoxicity of the medial prefrontal cortex in a rat model of Parkinson's disease. *Chin Med J (Engl)* 2017;130:2726-31. DOI PubMed PMC
102. Rasheed MZ, Andrabi SS, Salman M, et al. Melatonin improves behavioral and biochemical outcomes in a rotenone-induced rat model of Parkinson's disease. *J Environ Pathol Toxicol Oncol* 2018;37:139-50. DOI PubMed
103. Acuña-Castroviejo D, Coto-Montes A, Monti M, Ortiz GG, Reiter RJ. Melatonin is protective against MPTP-induced striatal and hippocampal lesions. *Life Sci* 1996;60:PL23-9. DOI PubMed
104. Jin BK, Shin DY, Jeong MY, et al. Melatonin protects nigral dopaminergic neurons from 1-methyl-4-phenylpyridinium (MPP+) neurotoxicity in rats. *Neurosci Lett* 1998;245:61-4. DOI PubMed
105. Naskar A, Prabhakar V, Singh R, Dutta D, Mohanakumar KP. Melatonin enhances L-DOPA therapeutic effects, helps to reduce its dose, and protects dopaminergic neurons in 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine-induced Parkinsonism in mice. *J Pineal Res* 2015;58:262-74. DOI PubMed
106. Brito-Armas JM, Baekelandt V, Castro-Hernández JR, González-Hernández T, Rodríguez M, Castro R. Melatonin prevents dopaminergic cell loss induced by lentiviral vectors expressing A30P mutant alpha-synuclein. *Histol Histopathol* 2013;28:999-1006.

[DOI PubMed](#)

107. Sae-Ung K, Uéda K, Govitrapong P, Phansuwan-Pujito P. Melatonin reduces the expression of alpha-synuclein in the dopamine containing neuronal regions of amphetamine-treated postnatal rats. *J Pineal Res* 2012;52:128-37. [DOI PubMed](#)
108. Iliff JJ, Wang M, Liao Y, et al. A paravascular pathway facilitates CSF flow through the brain parenchyma and the clearance of interstitial solutes, including amyloid β . *Sci Transl Med* 2012;4:147ra111. [DOI PubMed PMC](#)
109. Aspelund A, Antila S, Proulx ST, et al. A dural lymphatic vascular system that drains brain interstitial fluid and macromolecules. *J Exp Med* 2015;212:991-9. [DOI PubMed PMC](#)
110. Louveau A, Smirnov I, Keyes TJ, et al. Structural and functional features of central nervous system lymphatic vessels. *Nature* 2015;523:337-41. [DOI PubMed PMC](#)
111. Plog BA, Nedergaard M. The glymphatic system in central nervous system health and disease: past, present, and future. *Annu Rev Pathol* 2018;13:379-94. [DOI PubMed PMC](#)
112. Iliff JJ, Wang M, Zeppenfeld DM, et al. Cerebral arterial pulsation drives paravascular CSF-interstitial fluid exchange in the murine brain. *J Neurosci* 2013;33:18190-9. [DOI PubMed PMC](#)
113. Louveau A, Plog BA, Antila S, Alitalo K, Nedergaard M, Kipnis J. Understanding the functions and relationships of the glymphatic system and meningeal lymphatics. *J Clin Invest* 2017;127:3210-9. [DOI PubMed PMC](#)
114. Rasmussen MK, Mestre H, Nedergaard M. The glymphatic pathway in neurological disorders. *Lancet Neurol* 2018;17:1016-24. [DOI PubMed PMC](#)
115. Nedergaard M, Goldman SA. Glymphatic failure as a final common pathway to dementia. *Science* 2020;370:50-6. [DOI PubMed PMC](#)
116. Ringstad G, Vatnehol SAS, Eide PK. Glymphatic MRI in idiopathic normal pressure hydrocephalus. *Brain* 2017;140:2691-705. [DOI PubMed PMC](#)
117. Ding XB, Wang XX, Xia DH, et al. Impaired meningeal lymphatic drainage in patients with idiopathic Parkinson's disease. *Nat Med* 2021;27:411-8. [DOI PubMed](#)
118. Hablitz LM, Plá V, Giannetto M, et al. Circadian control of brain glymphatic and lymphatic fluid flow. *Nat Commun* 2020;11:4411. [DOI PubMed PMC](#)
119. Xie L, Kang H, Xu Q, et al. Sleep drives metabolite clearance from the adult brain. *Science* 2013;342:373-7. [DOI PubMed PMC](#)
120. Peng W, Achariyar TM, Li B, et al. Suppression of glymphatic fluid transport in a mouse model of Alzheimer's disease. *Neurobiol Dis* 2016;93:215-25. [DOI PubMed PMC](#)
121. Da Mesquita S, Louveau A, Vaccari A, et al. Functional aspects of meningeal lymphatics in ageing and Alzheimer's disease. *Nature* 2018;560:185-91. [DOI PubMed PMC](#)
122. Smith AJ, Yao X, Dix JA, Jin BJ, Verkman AS. Test of the 'glymphatic' hypothesis demonstrates diffusive and aquaporin-4-independent solute transport in rodent brain parenchyma. *Elife* 2017;6:e27679. [DOI PubMed PMC](#)

AUTHOR INSTRUCTIONS

1. Submission Overview

Before you decide to publish with *Ageing and Neurodegenerative Diseases (AND)*, please read the following items carefully and make sure that you are well aware of Editorial Policies and the following requirements.

1.1 Topic Suitability

The topic of the manuscript must fit the scope of the journal. Please refer to Aims and Scope for more information.

1.2 Open Access and Copyright

The journal adopts Gold Open Access publishing model and distributes content under the Creative Commons Attribution 4.0 International License. Copyright is retained by authors. Please make sure that you are well aware of these policies.

1.3 Publication Fees

AND is an open access journal. When a paper is accepted for publication, authors are required to pay Article Processing Charges (APCs) to cover its editorial and production costs. The APC for each submission is \$600. There are no additional charges based on color, length, figures, or other elements. For more details, please refer to OAE Publication Fees.

1.4 Language Editing

All submissions are required to be presented clearly and cohesively in good English. Authors whose first language is not English are advised to have their manuscripts checked or edited by a native English speaker before submission to ensure the high quality of expression. A well-organized manuscript in good English would make the peer review even the whole editorial handling more smoothly and efficiently.

If needed, authors are recommended to consider the language editing services provided by Charlesworth to ensure that the manuscript is written in correct scientific English before submission. Authors who publish with OAE journals enjoy a special discount for the services of Charlesworth via the following two ways.

Submit your manuscripts directly at <http://www.charlesworthauthorservices.com/~OAE>;

Open the link <http://www.charlesworthauthorservices.com/>, and enter Promotion Code “OAE” when you submit.

1.5 Work Funded by the National Institutes of Health

If an accepted manuscript was funded by National Institutes of Health (NIH), the authors may inform Editors of the NIH funding number. The Editors are able to deposit the paper to the NIH Manuscript Submission System on behalf of the authors.

2. Submission Preparation

2.1 Cover Letter

A cover letter is required to be submitted accompanying each manuscript. It should be concise and explain why the study is significant, why it fits the scope of the journal, and why it would be attractive to readers, etc.

Here is a guideline of a cover letter for authors' consideration:

In the first paragraph: include the title and type (e.g., Original Article, Review, Case Report, etc.) of the manuscript, a brief on the background of the study, the question the author sought out to answer and why;

In the second paragraph: concisely explain what was done, the main findings and why they are significant;

In the third paragraph: indicate why the manuscript fits the Aims and Scope of the journal, and why it would be attractive to readers;

In the fourth paragraph: confirm that the manuscript has not been published elsewhere and not under consideration of any other journal. All authors have approved the manuscript and agreed on its submission to the journal. Journal's specific requirements have been met if any.

If the manuscript is contributed to a Special Issue, please also mention it in the cover letter.

If the manuscript was presented partly or entirely in a conference, the author should clearly state the background information of the event, including the conference name, time and place in the cover letter.

2.2 Types of Manuscripts

There is no restriction on the length of manuscripts, number of figures, tables and references, provided that the manuscript is concise and comprehensive. The journal publishes Original Article, Review, Meta-Analysis, Case Report, Commentary, etc. For more details about paper type, please refer to the following table.

Manuscript Type	Definition	Abstract	Keywords	Main Text Structure
Original Article	An Original Article describes detailed results from novel research. All findings are extensively discussed.	Structured abstract including Aim, Methods, Results and Conclusion. No more than 250 words.	3-8 keywords	The main content should include four sections: Introduction, Methods, Results and Discussion.
Review	A Review paper summarizes the literature on previous studies. It usually does not present any new information on a subject.	Unstructured abstract. No more than 250 words.	3-8 keywords	The main text may consist of several sections with unfixed section titles. We suggest that the author includes an "Introduction" section at the beginning, several sections with unfixed titles in the middle part, and a "Conclusion" section in the end.
Case Report	A Case Report details symptoms, signs, diagnosis, treatment, and follows up an individual patient. The goal of a Case Report is to make other researchers aware of the possibility that a specific phenomenon might occur.	Unstructured abstract. No more than 150 words.	3-8 keywords	The main text consists of three sections with fixed section titles: Introduction, Case Report, and Discussion.
Meta-Analysis	A Meta-Analysis is a statistical analysis combining the results of multiple scientific studies. It is often an overview of clinical trials.	Structured abstract including Aim, Methods, Results and Conclusion. No more than 250 words.	3-8 keywords	The main content should include four sections: Introduction, Methods, Results and Discussion.
Systematic Review	A Systematic Review collects and critically analyzes multiple research studies, using methods selected before one or more research questions are formulated, and then finding and analyzing related studies and answering those questions in a structured methodology.	Structured abstract including Aim, Methods, Results and Conclusion. No more than 250 words.	3-8 keywords	The main content should include four sections: Introduction, Methods, Results and Discussion.
Technical Note	A Technical Note is a short article giving a brief description of a specific development, technique or procedure, or it may describe a modification of an existing technique, procedure or device applied in research.	Unstructured abstract. No more than 250 words.	3-8 keywords	/
Commentary	A Commentary is to provide comments on a newly published article or an alternative viewpoint on a certain topic.	Unstructured abstract. No more than 250 words.	3-8 keywords	/
Editorial	An Editorial is a short article describing news about the journal or opinions of senior editors or the publisher.	None required	None required	/
Letter to Editor	A Letter to Editor is usually an open post-publication review of a paper from its readers, often critical of some aspect of a published paper. Controversial papers often attract numerous Letters to Editor	Unstructured abstract (optional). No more than 250 words.	3-8 keywords (optional)	/
Opinion	An Opinion usually presents personal thoughts, beliefs, or feelings on a topic.	Unstructured abstract (optional). No more than 250 words.	3-8 keywords	/
Perspective	A Perspective provides personal points of view on the state-of-the-art of a specific area of knowledge and its future prospects. Links to areas of intense current research focus can also be made. The emphasis should be on a personal assessment rather than a comprehensive, critical review. However, comments should be put into the context of existing literature. Perspectives are usually invited by the Editors.	Unstructured abstract. No more than 150 words.	3-8 keywords	/

Clinical Observation	Clinical observation refers to records of the effects of treatment on hospitalized patients. It details symptoms, diagnosis and treatment of the disease to be reported. The characteristics of clinical reports include new or rare, complex adverse reactions, confusing symptoms or signs, examples of new theories, etc.	Unstructured abstract. No more than 250 words.	3-8 keywords	The main content should include four sections: Introduction, Methods, Results and Discussion.
----------------------	--	--	--------------	---

2.3 Manuscript Structure

2.3.1 Front Matter

2.3.1.1 Title

The title of the manuscript should be concise, specific and relevant, with no more than 16 words if possible. When gene or protein names are included, the abbreviated name rather than full name should be used.

2.3.1.2 Authors and Affiliations

Authors' full names should be listed. The initials of middle names can be provided. Institutional addresses and email addresses for all authors should be listed. At least one author should be designated as corresponding author. In addition, corresponding authors are suggested to provide their Open Researcher and Contributor ID upon submission. Please note that any change to authorship is not allowed after manuscript acceptance.

2.3.1.3 Abstract

The abstract should be a single paragraph with word limitation and specific structure requirements (for more details please refer to Types of Manuscripts). It usually describes the main objective(s) of the study, explains how the study was done, including any model organisms used, without methodological detail, and summarizes the most important results and their significance. The abstract must be an objective representation of the study: it is not allowed to contain results which are not presented and substantiated in the manuscript or exaggerate the main conclusions. Citations should not be included in the abstract.

2.3.1.4 Keywords

Three to eight keywords should be provided, which are specific to the article, yet reasonably common within the subject discipline.

2.3.2 Main Text

Manuscripts of different types are structured with different sections of content. Please refer to Types of Manuscripts to make sure which sections should be included in the manuscripts.

2.3.2.1 Introduction

The introduction should contain background that puts the manuscript into context, allow readers to understand why the study is important, include a brief review of key literature, and conclude with a brief statement of the overall aim of the work and a comment about whether the aim was achieved. Relevant controversies or disagreements in the field should be introduced as well.

2.3.2.2 Methods

Methods should contain sufficient details to allow others to fully replicate the study. New methods and protocols should be described in detail while well-established methods can be briefly described or appropriately cited. Experimental participants selected, the drugs and chemicals used, the statistical methods taken, and the computer software used should be identified precisely. Statistical terms, abbreviations, and all symbols used should be defined clearly. Protocol documents for clinical trials, observational studies, and other non-laboratory investigations may be uploaded as supplementary materials.

2.3.2.3 Results

This section contains the findings of the study. Results of statistical analysis should also be included either as text or as tables or figures if appropriate. Authors should emphasize and summarize only the most important observations. Data on all primary and secondary outcomes identified in the section Methods should also be provided. Extra or supplementary materials and technical details can be placed in supplementary documents.

2.3.2.4 Discussion

This section should discuss the implications of the findings in context of existing research and highlight limitations of the study. Future research directions may also be mentioned.

2.3.2.5 Conclusion

It should state clearly the main conclusions and include the explanation of their relevance or importance to the field.

2.3.3 Back Matter

2.3.3.1 Acknowledgments

Anyone who contributed towards the article but does not meet the criteria for authorship, including those who provided professional writing services or materials, should be acknowledged. Authors should obtain permission to acknowledge from all those mentioned in the Acknowledgments section. This section is not added if the author does not have anyone to acknowledge.

2.3.3.2 Authors' Contributions

Each author is expected to have made substantial contributions to the conception or design of the work, or the acquisition, analysis, or interpretation of data, or the creation of new software used in the work, or have drafted the work or substantively revised it.

Please use Surname and Initial of Forename to refer to an author's contribution. For example: made substantial contributions to conception and design of the study and performed data analysis and interpretation: Salas H, Castaneda WV; performed data acquisition, as well as provided administrative, technical, and material support: Castillo N, Young V.

If an article is single-authored, please include "The author contributed solely to the article." in this section.

2.3.3.3 Availability of Data and Materials

In order to maintain the integrity, transparency and reproducibility of research records, authors should include this section in their manuscripts, detailing where the data supporting their findings can be found. Data can be deposited into data repositories or published as supplementary information in the journal. Authors who cannot share their data should state that the data will not be shared and explain it. If a manuscript does not involve such issue, please state "Not applicable." in this section.

2.3.3.4 Financial Support and Sponsorship

All sources of funding for the study reported should be declared. The role of the funding body in the experiment design, collection, analysis and interpretation of data, and writing of the manuscript should be declared. Any relevant grant numbers and the link of funder's website should be provided if any. If the study is not involved with this issue, state "None." in this section.

2.3.3.5 Conflicts of Interest

Authors must declare any potential conflicts of interest that may be perceived as inappropriately influencing the representation or interpretation of reported research results. If there are no conflicts of interest, please state "All authors declared that there are no conflicts of interest." in this section. Some authors may be bound by confidentiality agreements. In such cases, in place of itemized disclosures, we will require authors to state "All authors declare that they are bound by confidentiality agreements that prevent them from disclosing their conflicts of interest in this work." If authors are unsure whether conflicts of interest exist, please refer to the "Conflicts of Interest" of *AND* Editorial Policies for a full explanation.

2.3.3.6 Ethical Approval and Consent to Participate

Research involving human subjects, human material or human data must be performed in accordance with the Declaration of Helsinki and approved by an appropriate ethics committee. An informed consent to participate in the study should also be obtained from participants, or their parents or legal guardians for children under 16. A statement detailing the name of the ethics committee (including the reference number where appropriate) and the informed consent obtained must appear in the manuscripts reporting such research.

Studies involving animals and cell lines must include a statement on ethical approval. More information is available at Editorial Policies.

If the manuscript does not involve such issue, please state "Not applicable." in this section.

2.3.3.7 Consent for Publication

Manuscripts containing individual details, images or videos, must obtain consent for publication from that person, or in the case of children, their parents or legal guardians. If the person has died, consent for publication must be obtained from the next of kin of the participant. Manuscripts must include a statement that a written informed consent for publication was obtained. Authors do not have to submit such content accompanying the manuscript. However, these documents must be available if requested. If the manuscript does not involve this issue, state "Not applicable." in this section.

2.3.3.8 Copyright

Authors retain copyright of their works through a Creative Commons Attribution 4.0 International License that clearly states how readers can copy, distribute, and use their attributed research, free of charge. A declaration "© The Author(s) 2022." will be added to each article. Authors are required to sign License to Publish before formal publication.

2.3.3.9 References

References should be numbered in order of appearance at the end of manuscripts. In the text, reference numbers should be placed in square brackets and the corresponding references are cited thereafter. If the number of authors is less than or equal

to six, we require to list all authors' names. If the number of authors is more than six, only the first three authors' names are required to be listed in the references, other authors' names should be omitted and replaced with "et al.". Abbreviations of the journals should be provided on the basis of Index Medicus. Information from manuscripts accepted but not published should be cited in the text as "Unpublished material" with written permission from the source. References should be described as follows, depending on the types of works:

Types	Examples
Journal articles by individual authors	Weaver DL, Ashikaga T, Krag DN, et al. Effect of occult metastases on survival in node-negative breast cancer. <i>N Engl J Med</i> 2011;364:412-21. [PMID: 21247310 DOI: 10.1056/NEJMoa1008108]
Organization as author	Diabetes Prevention Program Research Group. Hypertension, insulin, and proinsulin in participants with impaired glucose tolerance. <i>Hypertension</i> 2002;40:679-86. [PMID: 12411462]
Both personal authors and organization as author	Vallancien G, Emberton M, Harving N, van Moorselaar RJ, Alf-One Study Group. Sexual dysfunction in 1,274 European men suffering from lower urinary tract symptoms. <i>J Urol</i> 2003;169:2257-61. [PMID: 12771764 DOI: 10.1097/01.ju.0000067940.76090.73]
Journal articles not in English	Zhang X, Xiong H, Ji TY, Zhang YH, Wang Y. Case report of anti-N-methyl-D-aspartate receptor encephalitis in child. <i>J Appl Clin Pediatr</i> 2012;27:1903-7. (in Chinese)
Journal articles ahead of print	Odibo AO. Falling stillbirth and neonatal mortality rates in twin gestation: not a reason for complacency. <i>BJOG</i> 2018; Epub ahead of print [PMID: 30461178 DOI: 10.1111/1471-0528.15541]
Books	Sherlock S, Dooley J. Diseases of the liver and biliary system. 9th ed. Oxford: Blackwell Sci Pub; 1993. pp. 258-96.
Book chapters	Meltzer PS, Kallioniemi A, Trent JM. Chromosome alterations in human solid tumors. In: Vogelstein B, Kinzler KW, editors. The genetic basis of human cancer. New York: McGraw-Hill; 2002. pp. 93-113.
Online resource	FDA News Release. FDA approval brings first gene therapy to the United States. Available from: https://www.fda.gov/NewsEvents/Newsroom/PressAnnouncements/ucm574058.htm . [Last accessed on 30 Oct 2017]
Conference proceedings	Harnden P, Joffe JK, Jones WG, editors. Germ cell tumours V. Proceedings of the 5th Germ Cell Tumour Conference; 2001 Sep 13-15; Leeds, UK. New York: Springer; 2002.
Conference paper	Christensen S, Oppacher F. An analysis of Koza's computational effort statistic for genetic programming. In: Foster JA, Lutton E, Miller J, Ryan C, Tettamanzi AG, editors. Genetic programming. EuroGP 2002: Proceedings of the 5th European Conference on Genetic Programming; 2002 Apr 3-5; Kinsdale, Ireland. Berlin: Springer; 2002. pp. 182-91.
Unpublished material	Tian D, Araki H, Stahl E, Bergelson J, Kreitman M. Signature of balancing selection in Arabidopsis. <i>Proc Natl Acad Sci U S A</i> . Forthcoming 2002.

For other types of references, please refer to U.S. National Library of Medicine.

The journal also recommends that authors prepare references with a bibliography software package, such as EndNote to avoid typing mistakes and duplicated references.

2.3.3.10 Supplementary Materials

Additional data and information can be uploaded as Supplementary Materials to accompany the manuscripts. The supplementary materials will also be available to the referees as part of the peer-review process. Any file format is acceptable, such as data sheet (word, excel, csv, cdx, fasta, pdf or zip files), presentation (powerpoint, pdf or zip files), image (cdx, eps, jpeg, pdf, png or tiff), table (word, excel, csv or pdf), audio (mp3, wav or wma) or video (avi, divx, flv, mov, mp4, mpeg, mpg or wmv). All information should be clearly presented. Supplementary materials should be cited in the main text in numeric order (e.g., Supplementary Figure 1, Supplementary Figure 2, Supplementary Table 1, Supplementary Table 2, etc.). The style of supplementary figures or tables complies with the same requirements on figures or tables in main text. Videos and audios should be prepared in English and limited to a size of 500 MB.

2.4 Manuscript Format

2.4.1 File Format

Manuscript files can be in DOC and DOCX formats and should not be locked or protected.

2.4.2 Length

There are no restrictions on paper length, number of figures, or amount of supporting documents. Authors are encouraged to present and discuss their findings concisely.

2.4.3 Language

Manuscripts must be written in English.

2.4.4 Multimedia Files

The journal supports manuscripts with multimedia files. The requirements are listed as follows:

Videos or audio files are only acceptable in English. The presentation and introduction should be easy to understand. The

frames should be clear, and the speech speed should be moderate.

A brief overview of the video or audio files should be given in the manuscript text.

The video or audio files should be limited to a size of up to 500 MB.

Please use professional software to produce high-quality video files, to facilitate acceptance and publication along with the submitted article. Upload the videos in mp4, wmv, or rm format (preferably mp4) and audio files in mp3 or wav format.

2.4.5 Figures

Figures should be cited in numeric order (e.g., Figure 1, Figure 2) and placed after the paragraph where it is first cited;

Figures can be submitted in format of tiff, psd, AI or jpeg, with resolution of 300-600 dpi;

Figure caption is placed under the Figure;

Diagrams with describing words (including, flow chart, coordinate diagram, bar chart, line chart, and scatter diagram, *etc.*) should be editable in word, excel or powerpoint format. Non-English information should be avoided;

Labels, numbers, letters, arrows, and symbols in figure should be clear, of uniform size, and contrast with the background; Symbols, arrows, numbers, or letters used to identify parts of the illustrations must be identified and explained in the legend;

Internal scale (magnification) should be explained and the staining method in photomicrographs should be identified;

All non-standard abbreviations should be explained in the legend;

Permission for use of copyrighted materials from other sources, including re-published, adapted, modified, or partial figures and images from the internet, must be obtained. It is authors' responsibility to acquire the licenses, to follow any citation instruction requested by third-party rights holders, and cover any supplementary charges.

2.4.6 Tables

Tables should be cited in numeric order and placed after the paragraph where it is first cited;

The table caption should be placed above the table and labeled sequentially (e.g., Table 1, Table 2);

Tables should be provided in editable form like DOC or DOCX format (picture is not allowed);

Abbreviations and symbols used in table should be explained in footnote;

Explanatory matter should also be placed in footnotes;

Permission for use of copyrighted materials from other sources, including re-published, adapted, modified, or partial tables from the internet, must be obtained. It is authors' responsibility to acquire the licenses, to follow any citation instruction requested by third-party rights holders, and cover any supplementary charges.

2.4.7 Abbreviations

Abbreviations should be defined upon first appearance in the abstract, main text, and in figure or table captions and used consistently thereafter. Non-standard abbreviations are not allowed unless they appear at least three times in the text. Commonly-used abbreviations, such as DNA, RNA, ATP, *etc.*, can be used directly without definition. Abbreviations in titles and keywords should be avoided, except for the ones which are widely used.

2.4.8 Italics

General italic words like *vs.*, *et al.*, *etc.*, *in vivo*, *in vitro*; *t* test, *F* test, *U* test; related coefficient as *r*, sample number as *n*, and probability as *P*; names of genes; names of bacteria and biology species in Latin.

2.4.9 Units

SI Units should be used. Imperial, US customary and other units should be converted to SI units whenever possible. There is a space between the number and the unit (i.e., 23 mL). Hour, minute, second should be written as h, min, s.

2.4.10 Numbers

Numbers appearing at the beginning of sentences should be expressed in English. When there are two or more numbers in a paragraph, they should be expressed as Arabic numerals; when there is only one number in a paragraph, number < 10 should be expressed in English and number > 10 should be expressed as Arabic numerals. 12345678 should be written as 12,345,678.

2.4.11 Equations

Equations should be editable and not appear in a picture format. Authors are advised to use either the Microsoft Equation Editor or the MathType for display and inline equations.

2.5 Submission Link

Submit an article via <https://oaemesas.com/login?JournalId=and>.

3. Research and Publication Ethics

3.1 Research Involving Human Subjects

All studies involving human subjects must be in accordance with the Helsinki Declaration and seek approval to conduct the study from an independent local, regional, or national review body (e.g., ethics committee, institutional review board, *etc.*).

Such approval, including the names of the ethics committee, institutional review board, etc., must be listed in a declaration statement of Ethical Approval and Consent to Participate in the manuscript. If the study is judged exempt from ethics approval, related information (e.g., name of the ethics committee granting the exemption and the reason for the exemption) must be listed. Further documentation on ethics should also be prepared, as Editors may request more detailed information. Manuscripts with suspected ethical problems will be investigated according to COPE Guidelines.

3.1.1 Consent to Participate

For all studies involving human subjects, informed consent to participate in the studies must be obtained from participants, or their parents or legal guardians for children under 16. Statements regarding consent to participate should be included in a declaration statement of Ethical Approval and Consent to Participate in the manuscript. If informed consent is not required, the name of the ethics committee granting the exemption and the reason for the exemption must be listed. If any ethical violation is found at any stage of publication, the issue will be investigated seriously based on COPE Guidelines.

3.1.2 Consent for Publication

All articles published by *AND* are freely available on the Internet. All manuscripts that include individual participants' data in any form (i.e., details, images, videos, etc.) will not be published without Consent for Publication obtained from that person(s), or for children, their parents or legal guardians. If the person has died, Consent for Publication must be obtained from the next of kin. Authors must add a declaration statement of Consent for Publication in the manuscript, specifying written informed consent for publication has been obtained.

3.1.3 Trial Registration

AND requires all authors to register all relevant clinical trials that are reported in manuscripts submitted. *AND* follows the World Health Organization (WHO)'s definition of clinical trials: "A clinical trial is any research study that prospectively assigns human participants or groups of humans to one or more health-related interventions to evaluate the effects on health outcomes. Interventions include but are not restricted to drugs, cells, other biological products, surgical procedures, radiologic procedures, devices, behavioral treatments, process-of-care changes, preventive care, etc."

In line with International Committee of Medical Journal Editors (ICMJE) recommendation, *AND* requires the registration of clinical trials in a public trial registry at or before the time of first patient enrollment. *AND* accepts publicly accessible registration in any registry that is a primary register of the WHO International Clinical Trials Registry Platform or in ClinicalTrials.gov. The trial registration number should be listed at the end of the Abstract section.

Secondary data analyses of primary (parent) clinical trials should not be registered as a new clinical trial, but rather reference the trial registration number of the primary trial.

Editors of *AND* will consider carefully whether studies failed to register or had an incomplete trial registration. Because of the importance of prospective trial registration, if there is an exception to this policy, trials must be registered and the authors should indicate in the publication when registration was completed and why it was delayed. Editors will publish a statement indicating why an exception was allowed. Please note such exceptions should be rare, and authors failing to prospectively register a trial risk its inadmissibility to *AND*.

Authors who are not sure whether they need trial registration may refer to ICMJE FAQs for further information.

3.2 Research Involving Animals

Experimental research on animals should be approved by an appropriate ethics committee and must comply with institutional, national, or international guidelines. *AND* encourages authors to comply with the AALAS Guidelines, the ARRIVE Guidelines, and/or the ICLAS Guidelines, and obtain prior approval from the relevant ethics committee. Manuscripts must include a statement indicating that the study has been approved by the relevant ethical committee and the whole research process complies with ethical guidelines. If a study is granted an exemption from requiring ethics approval, the name of the ethics committee granting the exemption and the reason(s) for the exemption should be detailed. Editors will take account of animal welfare issues and reserve the right to reject a manuscript, especially if the research involves protocols that are inconsistent with commonly accepted norms of animal research.

3.3 Research Involving Cell Lines

Authors must describe what cell lines are used and their origin so that the research can be reproduced. For established cell lines, the provenance should be stated and references must also be given to either a published paper or to a commercial source. For de novo cell lines derived from human tissue, appropriate approval from an institutional review board or equivalent ethical committee, and consent from the donor or next of kin, should be obtained. Such statements should be listed on the Declaration section of Ethical Approval and Consent to Participate in the manuscript.

Further information is available from the International Cell Line Authentication Committee (ICLAC). *AND* recommends that authors check the NCBI database for misidentification and contamination of human cell lines.

3.4 Research Involving Plants

Experimental research on plants (either cultivated or wild), including collection of plant material, must comply with institutional, national, or international guidelines. Field studies should be conducted in accordance with local legislation, and the manuscript should include a statement specifying the appropriate permissions and/or licenses. *AND* recommends that authors comply with the IUCN Policy Statement on Research Involving Species at Risk of Extinction and the Convention on the Trade in Endangered Species of Wild Fauna and Flora.

For each submitted manuscript, supporting genetic information and origin must be provided for plants that were utilized. For research manuscripts involving rare and non-model plants (other than, e.g., *Arabidopsis thaliana*, *Nicotiana benthamiana*, *Oriza sativa*, or many other typical model plants), voucher specimens must be deposited in a public herbarium or other public collections providing access to deposited materials.

3.5 Publication Ethics Statement

OAE is a member of the Committee on Publication Ethics (COPE). We fully adhere to its Code of Conduct and to its Best Practice Guidelines.

The Editors of *AND* enforce a rigorous peer-review process together with strict ethical policies and standards to guarantee to add high-quality scientific works to the field of scholarly publication. Unfortunately, cases of plagiarism, data falsification, image manipulation, inappropriate authorship credit, and the like, do arise. The Editors of *AND* take such publishing ethics issues very seriously and are trained to proceed in such cases with zero tolerance policy.

Authors wishing to publish their papers in *AND* must abide to the following:

The author(s) must disclose any possibility of a conflict of interest in the paper prior to submission.

The authors should declare that there is no academic misconduct in their manuscript in the cover letter.

Authors should accurately present their research findings and include an objective discussion of the significance of their findings.

Data and methods used in the research need to be presented in sufficient detail in the manuscript so that other researchers can replicate the work.

Authors should provide raw data if referees and the Editors of the journal request.

Simultaneous submission of manuscripts to more than one journal is not tolerated.

Republishing content that is not novel is not tolerated (for example, an English translation of a paper that is already published in another language will not be accepted).

The manuscript should not contain any information that has already been published. If you include already published figures or images, please get the necessary permission from the copyright holder to publish under the CC-BY license.

Plagiarism, data fabrication and image manipulation are not tolerated.

Plagiarism is not acceptable in *AND*.

Plagiarism involves the inclusion of large sections of unaltered or minimally altered text from an existing source without appropriate and unambiguous attribution, and/or an attempt to misattribute original authorship regarding ideas or results, and copying text, images, or data from another source, even from your own publications, without giving credit to the source.

As to reusing the text that is copied from another source, it must be between quotation marks and the source must be cited. If a study's design or the manuscript's structure or language has been inspired by previous studies, these studies must be cited explicitly.

If plagiarism is detected during the peer-review process, the manuscript may be rejected. If plagiarism is detected after publication, we may publish a Correction or retract the paper.

Falsification is manipulating research materials, equipment, or processes, or changing or omitting data or results so that the findings are not accurately represented in the research record.

Image files must not be manipulated or adjusted in any way that could lead to misinterpretation of the information provided by the original image.

Irregular manipulation includes introduction, enhancement, moving, or removing features from the original image; grouping of images that should be presented separately, or modifying the contrast, brightness, or color balance to obscure, eliminate, or enhance some information.

If irregular image manipulation is identified and confirmed during the peer-review process, we will reject the manuscript. If irregular image manipulation is identified and confirmed after publication, we may publish a Retraction or retract the paper.

AND reserves the right to contact the authors' institution(s) to investigate possible publication misconduct if the Editors find conclusive evidence of misconduct before or after publication. OAE has a partnership with iThenticate, which is the most trusted similarity checker. It is used to analyze received manuscripts to avoid plagiarism to the greatest extent possible.

When plagiarism becomes evident after publication, we will retract the original publication or require modifications, depending on the degree of plagiarism, context within the published article, and its impact on the overall integrity of the published study. Journal Editors will act under the relevant COPE Guidelines.

4. Authorship

Authorship credit of *AND* should be solely based on substantial contributions to a published study, as specified in the following four criteria:

1. Substantial contributions to the conception or design of the work, or the acquisition, analysis, or interpretation of data for the work;
2. Drafting the work or revising it critically for important intellectual content;
3. Final approval of the version to be published;
4. Agreement to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

All those who meet these criteria should be identified as authors. Authors must specify their contributions in the section Authors' Contributions of their manuscripts. Contributors who do not meet all the four criteria (like only involved in acquisition of funding, general supervision of a research group, general administrative support, writing assistance, technical editing, language editing, proofreading, etc.) should be acknowledged in the section of Acknowledgement in the manuscript rather than being listed as authors.

If a large multiple-author group has conducted the work, the group ideally should decide who will be authors before the work starts and confirm authors before submission. All authors of the group named as authors must meet all the four criteria for authorship.

5. Reviewers Exclusions

You are welcome to exclude a limited number of researchers as potential Editors or reviewers of your manuscript. To ensure a fair and rigorous peer review process, we ask that you keep your exclusions to a maximum of three people. If you wish to exclude additional referees, please explain or justify your concerns—this information will be helpful for Editors when deciding whether to honor your request.

6. Editors and Journal Staff as Authors

Editorial independence is extremely important and *AND* does not interfere with editorial decisions. Editorial staff or Editors shall not be involved in the processing their own academic work. Submissions authored by editorial staff/Editors will be assigned to at least two independent outside reviewers. Decisions will be made by other Editorial Board members who do not have conflict of interests with the author. Journal staffs are not involved in the processing of their own work submitted to any OAE journals.

7. Conflict of Interests

AND require authors to declare any possible financial and/or non-financial conflicts of interest at the end of their manuscript and in the cover letter, as well as confirm this point when submitting their manuscript in the submission system. If no conflicts of interest exist, authors need to state “The authors declare no conflicts of interest”. We also recognize that some authors may be bound by confidentiality agreements, in which cases authors need to state “The authors declare that they are bound by confidentiality agreements that prevent them from disclosing their competing interests in this work”.

8. Editorial Process

8.1 Initial check

8.1.1 Initial manuscript check

New submissions are initially checked by the Managing Editor from the perspectives of originality, suitability, structure and formatting, conflicts of interest, background of authors, etc. Poorly-prepared manuscripts may be rejected at this stage. If your manuscript does not meet one or more of these requirements, we will return it for further revisions.

8.1.2 Publishing ethics

All manuscripts submitted to *AND* are screened using iThenticate powered by CrossCheck to identify any plagiarized content. Your study must also meet all ethical requirements as outlined in our Editorial Policies. If the manuscript does not pass any of these checks, we may return it to you for further revisions or decline to consider your study for publication.

8.2 Editorial assessment

Once your manuscript has passed the initial manuscript check, it will be assigned to an Assistant Editor, and then the Editor-in-Chief, or an Associate Editor in the case of a conflict of interest, will be notified of the submission and invited to review. Regarding Special Issue paper, after passing the initial check, the manuscript will be successively assigned to an

Assistant Editor, Guest Editor, and then to the Editor-in-Chief, or an Associate Editor in the case of conflict of interest for the Editor-in-Chief to review. The Editor-in-Chief, or the Associate Editor may reject manuscripts that they deem highly unlikely to pass peer review without further consultation. Once your manuscript has passed the editorial assessment, the Assistant Editor will start to organize peer-review.

8.3 Process

AND operates a single-blind review process. The technical quality of the research described in the manuscript is assessed by a minimum of two independent expert reviewers. The Editor-in-Chief is responsible for the final decision regarding acceptance or rejection of the manuscript. For controversial manuscripts, the Editor-in-Chief is responsible for making the final decision.

8.4 Decisions

Your research will be judged on technical soundness only, not on its perceived impact as judged by Editors or referees. There are three possible decisions: Accept (your study satisfies all publication criteria), Invitation to Revise (more work is required to satisfy all criteria), and Reject (your study fails to satisfy key criteria and it is highly unlikely that further work can address its shortcomings).

9. Contact Us

Journal Contact

Ageing and Neurodegenerative Diseases Editorial Office

Suite 1504, Plaza A, Xi'an National Digital Publishing Base, No. 996 Tiangu 7th Road, Gaoxin District, Xi'an 710077, Shaanxi, China.

Tel: +86 (0)29 8954 0089

Monica Wang
Managing Editor
editorialoffice@ageneudisjournal.com



www.oaepublish.com

Ageing and Neurodegenerative Diseases
(AND)

Los Angeles Office

245 E Main Street ste122, Alhambra,

CA 91801, USA

E-mail: editorialoffice@ageneudisjournal.com

Website: www.ageneudisjournal.com

