


Neuroimmunology and Neuroinflammation



EDITORIAL BOARD

Honorary Editor-in-Chief

V. Wee Yong (Canada)

Editors-in-Chief

Gang Zhao (China)
Athanassios P. Kyritsis (Greece)
Thomas Müller (Germany)

Associate Editors

Dale Ding (USA)
Dirk M. Hermann (Germany)
Michel Mittelbronn (Germany)
Domenico Praticò (USA)
Gianfranco Spalletta (Italy)

Editorial Board Members

Paulo Henrique Aguiar (Brazil)
Tomohiro Aoki (Japan)
Bi-Tao Bu (China)
Aswin Chari (UK)
Chu Chen (USA)
Xing-Yong Chen (China)
Yan Chen (China)
Steven J. Collins (Australia)
Randall L. Davis (USA)
Xue-Wen Fan (China)
Vicente Felipo (Spain)
Yara Dadalti Fragoso (Brazil)
Stephen D. Ginsberg (USA)

Ye Gong (China)
Elena Grebenciucova (USA)
Hong-Zhi Guan (China)
Jun-Ying He (China)
Li-Fang Hu (China)
Nunzio Iraci (Italy)
Wen Jiang (China)
Sofia Kasradze (Georgia)
Andis Klegeris (Russia)
Dana-Lynn Takeko Koomoa-Lange (USA)
Bin Li (China)
Li Li (China)
Zhu-Yi Li (China)
Sofia Markoula (Greece)
Salvadeeswaran Meenakshi-Sundaram (India)
Shekher Mohan (USA)
Daniele Orsucci (Italy)
Stéphane Peineau (France)
George Perry (USA)
Fabrizio Piazza (Italy)
Sara Grazia Maria Piccirillo (USA)
Xiao-Kun Qi (China)
Ji-Jun Shi (China)
Chrissa G. Sioka (Greece)
Michael J. Strong (USA)
Hua Su (USA)
Kyoungso Suk (South Korea)
Jun Tan (USA)
Rudolph E. Tanzi (USA)

Konstantinos Tsamis (Greece)
Kiran K. Velpula (USA)
Chao-Dong Wang (China)
Jia-Wei Wang (China)
Yan-Jiang Wang (China)
Kevin K.W. Wang (USA)
Sara Xapelli (Portugal)
Jun-Ping Xin (USA)
Guo-Yuan Yang (China)
Jun Yang (USA)
Yu-Feng Zang (China)
Xing-Hu Zhang (China)
Rui Zhao (China)

Language Editors

Shekher Mohan (UK)
Sara Grazia Maria Piccirillo (USA)
Huiling Tan (UK)
Randall L. Davis (USA)

Editorial Staffs

Lei Ma (China)
Rui-Rui Zhang (China)
Mei-Lu Jia (China)
Hai-Di Ding (China)
Guang-Zhe Zhu (China)
Jun-Yao Li (China)
Huan-Liang Wu (China)
Cai-Hong Wang (China)

GENERAL INFORMATION

About the Journal

Neuroimmunology and Neuroinflammation (NN), ISSN 2349-6142 (Online), ISSN 2347-8659 (Print), is a peer-reviewed online journal with print on demand compilation of articles published. The journal's full text is available online at www.nnjournal.net. The journal allows free access (Open Access) to its contents and permits authors to self-archive final accepted version of the articles on any OAI-compliant institutional/subject-based repository. The journal focuses on neuroimmunology and neuroinflammation, and the coverage extends to other basic and clinical studies related to neuroscience, including molecular biology, pharmacology, endocrinology, pathology, physiology, psychology, oncology, *etc.* The journal is indexed with CAS, Chaoxing "Domain" Publishing Platform, Cite Factor, CNKI, DRJI, EBSCO, Embase, Eurasian Scientific Journal Index, Google Scholar, Hinari, JournalGuide, JournalTOCs, J-Gate, ResearchBib, Root Indexing, SHERPA/RoMEO, Wanfang Data and Worldcat.

Information for Authors

Manuscripts must be prepared in accordance with Author Instructions.

Please check www.nnjournal.net/pages/view/author_instructions for details.

All manuscripts must be submitted online at www.editorialmanager.com/neurimm.

Copyright

The entire contents of the *NN* are protected under international copyrights. The journal, however, grants to all users a free, irrevocable, worldwide, perpetual right of access to, and a license to copy, use, distribute, perform and display the work publicly and to make and distribute derivative works in any digital medium for any reasonable purpose, subject to proper attribution of authorship and ownership of the rights. The journal also grants the right to make small numbers of printed copies for their personal use under the Creative Commons Attribution 4.0 License.

Copyright is reserved by © The Author(s) 2020.

Permissions

For information on how to request permissions to reproduce articles/information from this journal, please visit www.nnjournal.net.

Disclaimer

The information and opinions presented in the journal reflect the views of the authors and not of the journal or its Editorial Board or the Publisher. Publication does not constitute endorsement by the journal. Neither the *NN* nor its publishers nor anyone else involved in creating, producing or delivering the *NN* or the materials contained therein, assumes any liability or responsibility for the accuracy, completeness, or usefulness of any information provided in the *NN*, nor shall they be liable for any direct, indirect, incidental, special, consequential or punitive damages arising out of the use of the *NN*. The *NN*, nor its publishers, nor any other party involved in the preparation of material contained in the *NN* represents or warrants that the information contained herein is in every respect accurate or complete, and they are not responsible for any errors or omissions or for the results obtained from the use of such material. Readers are encouraged to confirm the information contained herein with other sources.

Published by

OAE Publishing Inc.
245 E Main Street ste122, Alhambra, CA 91801, USA
Website: www.oaepublish.com

Contacts

E-mail: nn_editor001@nnjournal.net
Website: www.nnjournal.net

CONTENTS

- 7 Favorable and unfavorable roles of microglia and macrophages in the pathologic central nervous system**
Junya Tanaka
Neuroimmunol Neuroinflammation 2020;7:73-91 <http://dx.doi.org/10.20517/2347-8659.2020.04>
- 8 Bilirubin and inflammation in neurodegenerative and other neurological diseases**
Sri Jayanti, Rita Moretti, Claudio Tiribelli, Silvia Gazzin
Neuroimmunol Neuroinflammation 2020;7:92-108 <http://dx.doi.org/10.20517/2347-8659.2019.14>
- 9 Cerebrospinal fluid biomarkers in idiopathic normal pressure hydrocephalus**
Xin-Jie Zhang, Jian Guo, Jun Yang
Neuroimmunol Neuroinflammation 2020;7:109-119 <http://dx.doi.org/10.20517/2347-8659.2019.018>
- 10 Cerebrospinal fluid amyloid beta and tau proteins in atypical Parkinsonism: a review**
Vasilios C. Constantinides, George P. Paraskevas, Fotini Boufidou, Mara Bourbouli, Panagiotis G. Paraskevas, Leonidas Stefanis, Elisabeth Kapaki
Neuroimmunol Neuroinflammation 2020;7:120-131 <http://dx.doi.org/10.20517/2347-8659.2019.22>
- 11 Assessment of cerebrospinal fluid α -synuclein as a potential biomarker in Parkinson's disease and synucleinopathies**
Ioanna Chalatsa, Katerina Melachroinou, Evangelia Emmanouilidou, Kostas Vekrellis
Neuroimmunol Neuroinflammation 2020;7:132-140 <http://dx.doi.org/10.20517/2347-8659.2020.01>
- 12 COVID-19 neurotropism and implications for therapy**
Mahasweta Das, Courtney Penn, Taylor Martinez, Karthick Mayilsamy, Andrew McGill, Alison Wiling, Shyam S. Mohapatra, Subhra Mohapatra
Neuroimmunol Neuroinflammation 2020;7:141-149 <http://dx.doi.org/10.20517/2347-8659.2020.36>
- 13 An association between mitochondria and microglia effector function: what do we think we know?**
Gaylia Jean Harry, Gabrielle Childers, Sahana Giridharan, Irisyunuel Lopez Hernandez
Neuroimmunol Neuroinflammation 2020;7:150-165 <http://dx.doi.org/10.20517/2347-8659.2020.07>
- 14 *Phf* a novel transcriptional repressor regulating inflammation in a mouse microglial cell line**
Sandra E. Muroy, Greg A. Timblin, Marcela K. Preininger, Paulina Cedillo, Kaoru Saijo
Neuroimmunol Neuroinflammation 2020;7:166-182 <http://dx.doi.org/10.20517/2347-8659.2020.16>

Review

Open Access



Favorable and unfavorable roles of microglia and macrophages in the pathologic central nervous system

Junya Tanaka

Department of Molecular and Cellular Physiology, Graduate School of Medicine, Ehime University, Toon, Ehime 791-0295, Japan.

Correspondence to: Prof. Junya Tanaka, Department of Molecular and Cellular Physiology, Ehime University, School of Medicine, Toon, Ehime 791-0295, Japan. E-mail: jtanaka@m.ehime-u.ac.jp

How to cite this article: Tanaka J. Favorable and unfavorable roles of microglia and macrophages in the pathologic central nervous system. *Neuroimmunol Neuroinflammation* 2020;7:73-91. <http://dx.doi.org/10.20517/2347-8659.2020.04>

Received: 8 Jan 2020 **First Decision:** 14 Feb 2020 **Revised:** 21 Feb 2020 **Accepted:** 12 Mar 2020 **Available online:** 10 Apr 2020

Science Editor: Jeffrey Bajramovic **Copy Editor:** Jing-Wen Zhang **Production Editor:** Jing Yu

Abstract

Resident microglia in the central nervous system (CNS) are activated rapidly in response to even minor pathologic changes in the CNS, releasing various cytokines, growth factors, reactive oxygen species and other bioactive substances, in addition to eliminating synapses and degenerating cells through phagocytosis. Monocytes in circulation invade the inflamed brain tissues and develop into macrophages that also produce several bioactive substances and engage in phagocytosis. This article introduces methods for distinguishing microglia and macrophages. The pathophysiological roles of resident microglia and macrophages are discussed in animal models with neuroinflammation in the brain either with or without disruption of the blood-brain barrier. Both cell types have ameliorating and aggravating effects on the pathologic CNS, and their different roles are addressed in this article. Furthermore, this article compares the effects of some pharmacological interventions to induce phenotypic cellular changes for improved outcomes of the pathologic CNS.

Keywords: Parkinson's disease, stroke, traumatic brain injury, axotomy, spinal cord, glucocorticoid, noradrenaline, bromovalerylurea

INTRODUCTION

Microglia and blood-borne macrophages play major roles in the pathophysiological processes in various kinds of pathologies of the central nervous system (CNS) by releasing numerous bioactive substances,



© The Author(s) 2020. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, sharing, adaptation, distribution and reproduction in any medium or format, for any purpose, even commercially, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.



including cytokines, growth factors, and reactive oxygen/nitrogen species, and phagocytosing degenerating cells and materials^[1-4]. In this article, we use “microglia” to denote resident microglia in the CNS, and “macrophages” to denote cells derived from circulating monocytes that have invaded (typically inflamed) CNS lesions with a disrupted blood-brain barrier (BBB). The two types of mesoderm-derived cells share many kinds of characteristics and surface antigens^[5,6]. In particular, when microglia become activated in response to severe pathologic events, such as stroke or traumatic brain injury (TBI), which accompany BBB breakdown, both cell types resemble each other in terms of morphology, functions, and cell marker expression. The term “amoeboid microglia”^[7] has long been used to denote extremely activated microglia displaying almost the same morphology as brain macrophages in severe lesions. Therefore, it is generally difficult to distinguish one from the other. This assumption originated from the historical observation by Rio-Hortega^[8], who reported that ramified microglia in the normal mature brain can, in severely damaged brains, turn into phagocytes exhibiting spherical shapes of the same morphology as that of blood-borne macrophages^[5]. Therefore, numerous studies have described CD11b-expressing (in the case of the rat brain, a monoclonal antibody OX-42-immunoreactive) spherical cells in severely injured brains in the acute phase as activated microglia, although most of them should be recognized as invading neutrophils^[9,10].

More recently, it is well-known that both cell types play major roles in pathophysiological processes^[2,11,12]. The neuroinflammatory processes influence outcomes of CNS diseases or injuries in both favorable and unfavorable ways. In this review, “favorable” is used to describe microglia and macrophages that are neuroprotective cells bringing about the better outcome in the pathologic CNS than “unfavorable” ones that exert deleterious effects on the survival of neurons and other parenchymal cells. Therefore, various kinds of interventions, including pharmacological treatment and rehabilitation, have been studied in laboratory and clinical settings to determine if they enhance favorable responses while also suppressing the deleterious effects of microglia and macrophages^[1,4,13,14]. This aim may be interpreted as attempting to induce M2-polarized or alternatively activated phenotypes of these cells^[15-17]. However, phenotypes of activated microglia and macrophages cannot be classified clearly into M1-polarized or M2-polarized cells^[18,19].

Diseases and injuries of the CNS can be categorized by the absence of BBB disruption^[20,21]. In the absence of BBB disruption, the infiltration of circulating leukocytes is limited, and microglia play a central role as immune cells. When the BBB is disrupted, infiltrating leukocytes play much more significant roles^[9,22]. In particular, invading monocytes increase hugely in number because of their strong proliferative nature and end up occupying almost the entire area of the lesion in either stroke or traumatic injury^[11,12]. Marked accumulation of blood-borne macrophages is seen in malignant brain tumor masses, which are termed tumor-associated macrophages^[23,24]. Tumor-associated macrophages contribute to tumor growth and angiogenesis by secreting many kinds of growth factors, including vascular endothelial growth factor. Tumor-associated macrophages are not discussed in this article.

Thus, microglia and macrophages are the critical cells in CNS diseases and injuries and have a profound impact on patient outcomes. Therefore, many studies have investigated interventions to both suppress the unfavorable effects of microglia and macrophages and induce their favorable functions. This article also deals with several interventions targeting microglia and macrophages. There are three types of resident macrophages that are distinct from microglia and the invaded blood-borne macrophages. Those are the meningeal, perivascular and choroid plexus macrophages. Although they play significant roles in health and pathology, they are not discussed in this review.

DISTINGUISHING BETWEEN MICROGLIA AND MACROPHAGES

Discriminating microglia from macrophages has been considered to be very difficult. This is partly due to the old notion that microglia in the brain are derived from circulating monocytes^[7,25]. Moreover, microglia

have long been thought to be able to display almost completely spherical or amoeboid morphology when they become fully activated in the core of severe brain insults such as stroke or trauma^[5]. Both cell types share numerous immune cell markers^[26]. Therefore, distinguishing between them has been very problematic. However, many methods have been now established for this purpose.

Morphological characteristics

Despite the historical view, even highly activated microglia do not display spherical morphology^[9,27]. In a rat stroke model prepared by transient occlusion of middle cerebral artery (tMCAO), activated microglia in the peri-ischemic regions exhibit enlarged somata and shortened processes that are distinct from spikes. The processes can be identified with immunoreactivity to microglia/macrophage markers such as Iba1 or CD11b. Conversely, blood-borne macrophages and neutrophils, the latter of which infiltrate in abundance, do not have processes even though they may have a polygonal shape and short spikes^[27]. Thus, microglia can be distinguished from blood-borne cells via morphological observation^[10]. This may be the simplest method for specific identification of microglia.

Specific markers to distinguish the two cell types in the pathologic brains

Iba1, CD11b, CD45, and CD68 have long been used to identify resident microglia; however, these markers are more strongly expressed by infiltrating macrophages than by activated microglia^[9]. As CD11b is a marker for myeloid cells and CD45 is a marker for all of the leukocytes, they are not suitable for identifying microglia in the pathologic CNS. To examine specific roles of resident microglia in CNS pathology, they must be distinguished from blood-borne cells that are macrophages, lymphocytes, and neutrophils. Thus, considerable effort has been dedicated to finding microglia-specific markers that are not expressed by macrophages and other blood-borne cells^[28]. Comprehensive gene analyses using RNAseq and/or microarray analyses have identified genes that are expressed predominantly by microglia rather than by macrophages such as *Cx3cr1*, *Gpr34*, *P2ry12*, *P2ry13*, *Siglech*, *Tmem119*, and *Trem2*^[29-32]. Of the specific marker candidates, TMEM119 and Siglec-H may be the most promising for immunohistochemical discrimination of microglia from macrophages.

Transmembrane protein 119, commonly known as TMEM119, was identified by Bennet *et al.*^[33] Specific expression of TMEM119 by microglia has been demonstrated by immunohistochemical staining, flow cytometry analyses, and *in situ* hybridization^[33,34]. TMEM119 was identified originally as a type I transmembrane protein expressed by murine osteoblasts and is responsible for their differentiation^[35]. It is not expressed by microglia in immature murine brains, but its expression increases along with their development or ramification^[33]. In aged human brains either with or without Alzheimer's disease (AD), microglia expressed TMEM119^[34]. Activated microglia with enlarged somata in close proximity to amyloid plaques are less immunoreactive to TMEM119 antibody than are resting (or homeostatic) microglia. Moreover, TMEM119-expressing microglia in AD brains either do not or only weakly express the polarized markers CD80, CD163, or CD206. In a TBI model, ramified or homeostatic microglia express TMEM119 at higher levels than do activated microglia^[10]. Therefore, TMEM119 is particularly suitable for identifying homeostatic microglia with a ramified shape in both the normal and the injured mature brain.

Sialic-acid-binding immunoglobulin-like lectin-H (Siglec-H) is another promising marker for immunohistochemical discrimination of microglia from macrophages^[36]. Siglec-H is a single-pass transmembrane protein that was identified originally as a member of a CD33-related Siglec family. Siglec-H is barely expressed by circulating monocytes and their derived macrophages. In contrast to TMEM119, Siglec-H is expressed continually by activated microglia and by microglia in immature brains. Siglec-H may mediate signals necessary for phagocytosis by microglia^[37].

Bone marrow transplantation

Bone marrow transplantation (BMT) has long been a reliable method for identifying blood-borne cells in the CNS with BBB breakdown^[28]. After ~10 Gy irradiation to cause near-total death of the host's bone

marrow, the bone marrow from transgenic animals that ubiquitously express fluorescent proteins such as enhanced green fluorescent protein is transplanted^[12]. In the transplanted brain, blood-borne macrophages bear fluorescence but not resident microglia^[27]. However, BMT usually leads to partial chimera, which can make it difficult to analyze the results. With reconstruction of the bone marrow, it is a long time before the anemia disappears. Radiation may cause degeneration of neural cells, such as NG2 glia and neurons, which may affect the results^[38]. As radiation disrupts the BBB while increasing monocyte infiltration, BMT produces donor-derived microglia or microglia-like ramified cells in the brain parenchyma^[28]. Thus, the BMT may provide firm evidence showing the presence of blood-borne cells in the CNS, but it should be noted that BMT itself will significantly change the brain functions due to the toxic effects on the neural cells.

Flow cytometry

Microglial cells belong to a group of myeloid leukocytes and express a macrophage colony-stimulating factor receptor known as CSF-1R. Therefore, microglia express a myeloid cell marker CD11b and a pan-leukocyte marker CD45. Based on this finding, microglia have been analyzed by flow cytometry using antibodies to CD11b and CD45^[10,39]. Because of faint expression of CD45 by homeostatic microglia, immunohistochemical detection of this expression tends to be difficult. Conversely, macrophages and neutrophils express CD45 rather strongly^[10]. However, weak CD45 expression can be detected easily by flow cytometry and is an advantage in flow cytometry analysis; microglia can be defined as CD11b⁺/CD45^{lo} and macrophages as CD11b⁺/CD45^{hi}. Even activated microglia express at lower levels of CD45 than do macrophages. Moreover, activated and homeostatic microglia can be distinguished by flow cytometry based on forward and side scatter value; activated microglia have larger somata (larger forward scatter values) and more intracellular organelles (larger side scatter values) than do homeostatic microglia^[10]. Flow cytometry analyses can be used to isolate the cells by cell sorting^[10,39]. If the cells are treated appropriately to prevent degradation of proteins, RNA, or DNA, the sorted cells can be used for either Western blotting or PCR.

Flow cytometry analyses of either cultured microglia or circulating monocytes are simple^[40,41]. Nevertheless, analyses of microglia and macrophages in brain tissues have been difficult because of difficulties in dissociating the tissues into single cells. However, dissociation kits and apparatus are now available for preparing neural cell suspensions and are designed appropriately for dissociation of rodent and human brains^[10,39]. The sorted cells can also be used for culturing and/or functional analyses.

RESPONSE OF MICROGLIA IN BRAIN PATHOLOGY IN THE ABSENCE OF BBB BREAKDOWN

Microglia become activated in response to even minor pathological events that do not involve BBB disruption. This section discusses microglia in Parkinson's disease (PD), peripheral nerve injury, and Carbon monoxide (CO) intoxication. Minute activation of homeostatic microglia accompanying circadian changes can be observed in the normal mature brain; microglia exhibit weakly activated phenotypes around the time of onset of sleep^[39].

PD

PD is the second most frequent neurodegenerative disorder after AD. As BBB breakdown is not apparent in PD, infiltration of leukocytes, including monocytes, is not often seen. In PD pathophysiology, dopaminergic (DA) neurons in the substantia nigra pars compacta (SNc) in the mesencephalon primarily undergo degeneration, leading to microglial activation in the vicinity of the degenerating neurons^[42-44]. The activated microglia release potentially neurotoxic substances, such as either reactive oxygen/nitrogen species or glutamate^[45,46]. The microglia-derived proinflammatory cytokines and chemokines may also contribute to aggravation. Injection of 6-hydroxydopamine (6-OHDA) into the striatum or medial forebrain bundle is used to prepare the PD rat model^[47,48]. In the model, DA neurons in the SNc primarily

undergo degeneration, and microglia then become activated in response to damage-associated molecular patterns (DAMPs), such as high mobility group box-1 protein (HMGB1) released from damaged neurons^[49]. The activated microglia may exacerbate the degenerative processes of DA neurons. Another rat PD model is prepared via injection of lipopolysaccharide (LPS) either direct into the SNc or in its vicinity^[48]. In this model, microglial activation is primarily induced followed by DA neuron degeneration, suggesting that activated microglia could be a key cause of neuronal degeneration.

The hematopoietic cytokines interleukin-3 (IL-3) and granulocyte/macrophage colony-stimulating factor (GM-CSF) have been shown to modulate the phenotype of microglia in the SNc while suppressing proinflammatory nature and increasing secretion of neurotrophic factors, insulin-like growth factor 1 (IGF-1), and hepatocyte growth factor (HGF)^[44]. Following this, DA neurons increase the expression of anti-apoptotic factor Bcl-xL, and the symptoms of PD are ameliorated. This suggests that further investigation of microglial phenotypes would lead to a potential intervention for neurological disorders.

Microglia in the SNc have been the sole focus of studies investigating microglia in PD pathology. However, upon immunostaining sections of the mesencephalon of PD model rats with antibodies to microglia markers, activation of microglia in the substantia nigra pars reticulata (SNr) was more apparent than that of those in the SNc^[47]. The activated microglia in the SNr bore large CD68⁺ phagosomes in their cytoplasm, in which synaptic proteins were included. In the PD pathology, glutamatergic neurons in the subthalamic nuclei (STN) become hyperactivated and release excess amounts of glutamate in the basal ganglia outputs that are the SNr and the internal segment of the globus pallidus (GPI). The change causes bradykinesia, rigidity, and other PD symptoms. Activated microglia with large phagosomes are present not only in the SNr but also in the GPI. They internalize the glutamatergic synapses from the hyperactive STN. As the neurological deficits do not manifest until most DA neurons are lost, it is likely that there are some significant compensatory mechanisms that prevent the symptoms from appearing^[50]. Microglia should contribute to this compensation by eliminating hyperactive glutamatergic synapses. Administration of a single high dose of a synthetic glucocorticoid dexamethasone (Dex) to the PD model rats aggravated their motor deficits. Dex suppresses CD68 expression in the SNr and GPI, suggesting suppression of microglial phagocytosis.

Conversely, chronic administration of glucocorticoid^[51,52] and other anti-inflammatory agents, such as bromovalerylurea (BU)^[43], has been shown in laboratory settings to ameliorate the outcome of the motor deficits, likely because of suppression of proinflammatory activation of microglia in the SNc. These findings reveal that microglia play both ameliorative and detrimental roles. There appear to be two forms of microglia activation: one is characterized by the production of proinflammatory mediators found in the SNc^[43,44], and the other is characterized by enhanced phagocytic ability in the SNr and GPI^[47]. The dual role of microglia in the PD pathophysiology is summarized in a schematic diagram [Figure 1].

Despite significant evidence showing the involvement of microglia in the DA neuron loss in animal PD models, it is not clear whether microglia actually affect the pathology of human PD cases. A few clinical trials have shown the positive effects of antiinflammatory drugs^[53], and activated microglia are found in PD patients' brains by *in vivo* imaging with positron emission tomography^[54]. However, there is still no firm evidence demonstrating that microglia actually induce DA neuron death in human PD cases. Most anti-inflammatory interventions are not ameliorative^[55,56]. This contrasts with animal model cases, in which anti-inflammatory interventions markedly suppress DA neuron death. The discrepancy between animal PD models and human cases may be partially attributable to the speed of the pathological processes. The animal model is acutely prepared, whereas human PD is a chronic disease. Furthermore, when PD is diagnosed based on motor symptoms, more than 60 % of DA neurons in the SNc are degenerated. In animal models, anti-inflammatory drugs are often given simultaneously with or even before administration




Figure 1. Both favorable and unfavorable activated microglia participate in the pathophysiology of PD. Unfavorable microglia present in the SNc release neurotoxic proinflammatory mediators while accelerating the degeneration of DA neurons. Favorable microglia present in the SNr and GPi engage in eliminating hyperactive glutamatergic synapses from the STN in the indirect pathway of the basal ganglia. This figure is based primarily on the study by Aono *et al.*^[47]. PD: Parkinson's disease; SNc: substantia nigra pars compacta; DA: dopaminergic; SNr: substantia nigra pars reticulata; GPi: globus pallidus; STN: subthalamic nuclei; GPe: globus pallidus pars externa

of neurotoxic substances causing DA neuron loss. Even if anti-inflammatory treatments could delay or prevent progressive neuronal loss in PD pathology, it is a prerequisite for the treatments that diagnosis is made at much earlier time points when most DA neurons in the SNc are still viable.

Peripheral nerve injury

Peripheral nerve injuries induce activation of microglia in close proximity to damaged neurons. In particular, facial nerve transection has been used frequently to observe the response of microglia^[57]. Microglia become activated and proliferate while attaching intimately to the axotomized neurons. Direct contact with neurons may be one of the direct causes of the microglial activation^[58]. The activated microglia detach afferent axonal endings through "synaptic stripping"^[57,59]. In the axotomy model, the activated microglia may be neuroprotective through releasing a plethora of neuroprotective factors^[60]. Axotomy and spinal cord ischemia both caused similar changes of microglia around the alpha motoneurons in the anterior horn^[61]. Synapses surrounding the neurons disappeared when activated microglia attached intimately to damaged neurons.

Constriction injury of the sciatic nerve is another model that is used often to study the responses of microglia in the spinal cord. This model is well known to cause chronic neuropathic pain^[62]. Microglia exhibit activated morphology in the posterior horn, and they phagocytose myelin elements^[63]. Constriction injury-induced hyperalgesia and the activation of microglia in the posterior horn are sustained chronically. Microglia in the anterior horn also become activated while surrounding damaged motoneurons and remove afferent synapses, as do microglia do in the facial nerve axotomy model. Different from the chronic sensory impairment, Constriction injury-induced motor deficits become ameliorated quite rapidly, indicating that the activated microglia in the anterior horn may be neuroprotective. Thus, microglia could become either protective/favorable or destructive/unfavorable cells.

CO intoxication

CO intoxication causes serious adverse effects in brain functions that are known as delayed encephalopathy. Compared with hypoxia-induced disorder, CO intoxication causes more severe memory impairment and more aggravated degeneration of neuronal cells in the hippocampus^[64]. Moreover, CO intoxication causes damage of oligodendrocytes, myelin and NG2 glia. Notably, CO intoxication induces profound loss of microglia. Expression of neurotrophic factors, such as IGF-1, HGF, platelet-derived growth factor, and

basic fibroblast growth factor (bFGF) is reduced considerably, whereas expression of pro-apoptotic factors, such as Bid, Bad, and Bax, is increased, leading to marked loss of the hippocampal neurons. These findings indicate that microglia contribute significantly to supporting neuronal survival.

RESPONSE OF MICROGLIA AND MACROPHAGES IN THE BRAIN WITH ISCHEMIC AND TRAUMATIC INJURIES THAT CAUSE BBB BREAKDOWN

Severe damage to brains and spinal cords disrupts the BBB, resulting in massive infiltration of blood-borne cells, such as monocytes and neutrophils^[9,22]. Discrimination of the roles of these immune cell types in pathophysiological processes is becoming possible, although further study is necessary^[10]. Different responses of microglia and macrophages are seen in the ischemic lesion core and the ischemic penumbra or peri-ischemic region^[11,27]. It should be noted that microglia are very vulnerable to various kinds of severe brain insults, including CO intoxication, as mentioned above^[64]. Six hours after reperfusion in tMCAO (90 min-occlusion) in a rat model of severe stroke, microglia undergo apoptotic degeneration in the lesion core, whereas neurons appear unchanged^[9,11]. Therefore, blood-borne macrophages and neutrophils are the main immune cells in the lesion cores and during the acute phase. However, microglia are both viable and become activated in the peri-region neighboring the core of the lesion and modulate pathological processes.

Roles of microglia

The activated microglia in the peri-ischemic regions bear large phagosomes that can be recognized by immunohistochemical staining of CD68^[27]. Moreover, they frequently express NG2 chondroitin sulfate proteoglycan (NG2), which may be another marker for phagocytosing microglia and macrophages^[27]. CD68⁺-phagocytosing microglia in the SNr and GPi in the rat PD model brains also express NG2^[47]. Such NG2⁺/CD68⁺ microglia engage in phagocytosis of degenerating neurons in the very limited region located along the border zone delineating the ischemic core and the penumbra^[27]. The region is termed the demarcation zone^[65] and is characterized by high expression of NG2. Neurodegeneration, known as delayed neuronal death, is still progressing in the zone in the subacute phase, which has long been a therapeutic target in stroke research to ameliorate the outcome of stroke.

Microglia are assumed to eliminate still viable neurons via phagocytosis in the ischemic penumbra^[66,67]. Because of decreased blood flow, neurons reduce their ATP synthesis and, thus, frequently externalize phosphatidylserine (PS) on their surface. PS is a typical eat-me-signal molecule that is recognized by molecules, such as either Milk fat globule EGF-like factor 8 (MFG-E8) or protein S, which are, in turn, recognized by either vitronectin receptor or Mer receptor tyrosine kinase (MerTK) expressed by microglia. Expression of MFG-E8 and MerTK is enhanced either in activated microglia or in macrophages in the ischemic lesions. Knocking out expression of MerTK or MFG-E8 by microglia prevented the delayed neuronal loss considerably. These findings may indicate that CD68⁺/NG2⁺-phagocytosing microglia are aggravating cells in stroke pathology.

Administration of the CSF1R antagonist PLX3397 to mice depletes microglia. This pharmacological intervention was used to study the overall effects of microglia on the outcome of ischemic brain insults. Eliminating microglia increases infarct volume, indicating that the overall effects of microglia on the ischemic brain are ameliorative. Microglia in the non-ischemic regions may maintain the neuronal circuitry, suppress proinflammatory activation of astrocytes, and prevent infiltration by various leukocytes, such as T cells, monocytes, and granulocytes^[68,69].

Activated microglia in the ischemic brain release a considerable quantity of transforming growth factor β 1 (TGF β 1)^[27], which is a strong immunosuppressive cytokine. Ischemic brain lesions should contain

abundant DAMPs, such as HMGB1^[70] and peroxiredoxin^[71], which are potential ligands for toll-like receptors (TLRs). DAMPs cause proinflammatory activation of microglia as lipopolysaccharide (LPS). However, expression of proinflammatory cytokine by both microglia and macrophages in the ischemic brain is not very remarkable^[72]. TLR ligands strongly induce expression of inducible nitric oxide synthase (iNOS) by microglia in culture. However, microglia in the ischemic brain either do not express iNOS protein or do so very faintly. TGFβ1 expression increases gradually until 7 days after the ischemic insults. Either LPS or DAMPs induce phosphorylation of IκB kinase (IKK), causing degradation of IκB, enabling the major proinflammatory transcription factor NFκB to translocate into nuclei and resulting in increased transcription of mRNA for proinflammatory mediators^[73]. However, after being incubated with TGFβ1 for ~24 h, primary cultured microglia do not respond to LPS treatments, even when the TGFβ1 is removed from the culture media. The TGFβ1-treated microglia cannot be classified into M2 polarized cells as they do not express M2 markers. TGFβ1 also inhibits phosphorylation of signal transducers and activators of transcription 1 (STAT1) and expression of interferon regulatory factor 1 (IRF1), both of which augment proinflammatory reactions of microglia in a sustained manner. When injected into the parenchyma of the ischemic brain, both microglia and macrophages lose their immunoreactivity to phosphorylated IKK (pIKK). Expression of other anti-inflammatory cytokines, such as IL-4, IL-10, or IL-13, is much weaker than is that of TGFβ1. These findings indicate that TGFβ1 is a key ameliorating factor released by both microglia and macrophages.

Roles of macrophages

Macrophages accumulate densely in the core of the ischemic and traumatic lesions where almost all microglia disappear^[11,12,19,74]. Therefore, macrophages play central roles in the immunological modulation of pathophysiological processes in these lesions. As most of the accumulated macrophages express NG2^[74], they have been called brain Iba1⁺/NG2⁺ cells (BINCs)^[11]. At least some are involved in removing degenerated materials in the core. The macrophage-precursor monocytes invade inflamed brain tissues in ischemic and traumatic injuries through recognizing chemokines that are typically either CCL2 or monocyte chemoattractant protein 1 and either CX3CL1 or fractalkine^[22]. Kinetic study in the rat tMCAO models showed that expression of CCL2 and CX3CL1 disappeared rapidly after tMCAO, within 2 days. When macrophage proliferation was inhibited by a single administration of the anti-cancer drug 5-fluorouracil 48 h after tMCAO, accumulation of macrophages at 7 dpr did not occur^[12]. Therefore, proliferation of monocytes that infiltrate within 2 days of the onset of the events causes the massive accumulation of macrophages in the core of the lesion^[11]. The reduction in number of the accumulated macrophages often causes death of the model rats. When BINCs were isolated from ischemic rat brains and transplanted into the core lesions of other stroke model rats, the BINCs proliferate hugely and the outcomes were ameliorated greatly. BINCs express various neuroprotective factors, such as IGF-1 and HGF, that may contribute to improved outcomes. Although BINCs were also found in lesions in aged human brains of stroke cases, however, the density was considerably less than that in the young rat model lesions^[12]. This may be one reason why human stroke cases become more severe than do the young rat models. Overall, the collective effects of macrophages on ischemic brains appear to be ameliorative.

However, the effects of monocyte-derived macrophages may not be favorable during the acute phase. Microglia and macrophages were isolated individually from TBI model rats at 1.5 days after injury using a fluorescence-activated cell sorter based on the different levels of CD45 expression^[10]. In this TBI model, oxidative injury may play a significant detrimental role in inducing neuronal degeneration, as an oxidative product 8-hydroxydeoxyguanine (8-OHdG) was found in neuronal nuclei. Reactive oxygen species (ROS) is the cause for this product, and most of the ROS may be derived from the mitochondria of macrophages^[75]. Infiltrating macrophages produce much higher levels of mitochondrial ROS than do microglia. ROS is also produced by NADPH oxidase activity, and macrophages express much higher levels of NADPH oxidase than do microglia. Moreover, expression of IL-1β and iNOS is more significant in macrophages than in

microglia. Although NO is occasionally described as a neuroprotective factor^[76], it can bind to superoxide O_2^- and form the highly toxic molecule peroxynitrite (ONOO⁻)^[77]. Conversely, microglia can save neurons from degeneration induced by oxidative stress^[78,79]. Collectively, the roles of blood-borne macrophages are more detrimental than are those of microglia in the earliest phase of severe brain injuries.

Some different phenotypes of macrophages accumulate in the traumatic and ischemic lesion cores^[10,19]. During the acute phase, 2 days after the onset of stroke, most macrophages exhibit a rather neurotoxic phenotype characterized by expression of IL-1 β and iNOS and release of ROS^[10]. However, most macrophages exhibit neuroprotective phenotypes characterized by expression of IGF-1 and HGF. This does not imply that a single population of macrophages turns into the two different (neuroprotective and neuro-destructive) populations depending on the pathological processes. As the neuroprotective macrophages characteristically express NG2, they are termed BINC. However, BINC cannot be classified as M2-polarized macrophages; they express a typical M1 marker CD86 and not an M2 marker CD163, whereas they do not express IL-1 β and iNOS in the ischemic lesion core. There is a minor subpopulation of Iba1⁺/CD68⁺ macrophages; though they do not express either NG2 or CD86, they express IL-1 β , CCL2, and iNOS, in addition to TLR4, which may be the source of the proinflammatory factors. Characteristically, they express CD200^[19,80], which mediates immunosuppressive signals primarily to myeloid cells expressing its receptor CD200R. Though the CD200⁺/NG2⁻ macrophages are likely the predominant population during the acute phase of severe brain injury, they undergo gradual apoptotic degeneration due to the oxidative injury, as evidenced by the accumulation of 8OHdG⁺ materials in their nuclei^[10]. Importantly, as they are not proliferative, they decline in number along with the pathological processes^[19]. Alternatively, BINC may be a small population during the acute phase, but they are highly proliferative and become the main macrophage population in the lesion core. Figure 2 summarizes the two types of macrophages.

In conclusion, macrophages also play both favorable and unfavorable roles in brain pathology. However, unlike the case with microglia, there are different macrophage populations that exert favorable and unfavorable effects independently.

PHARMACOLOGICAL INTERVENTIONS

The term “double-edged sword” has been used frequently to describe the favorable and unfavorable effects of microglia and macrophages^[81-83]. Pharmacological interventions have long been sought that would either induce or strengthen the favorable effects of these cells^[15,84]. However, in the case of macrophages, it may be more likely that there are different subpopulations with different effects on neurons rather than there being opposing effects of a single population of macrophages. Nevertheless, numerous pharmacological interventions have been shown to modulate the natures of both microglia and macrophages. Some of these are discussed below. Here it is better to once again confirm the notion that the favorable and unfavorable phenotypes for neuronal survival are incompatible with the M1 and M2 classifications^[18,19,72].

Glucocorticoids

Glucocorticoids have strong immunosuppressive effects on many kinds of cells and may be the most often-used anti-inflammatory agent in laboratory and clinical settings. Glucocorticoids may have stronger anti-inflammatory effects on microglia than do adrenergic agonists, minocycline, troglitazone, and antidepressants [Figure 3]. They cause microglial cells to shrink, reduce lysosomal enzyme activities, and suppress proliferation of primary cultured rat microglia^[85]. They suppress expression of the proinflammatory mediators by LPS-treated cultured microglia primarily at the transcription level^[43]. Glucocorticoids can bind not only to the glucocorticoid receptor but also to the mineralocorticoid receptor^[85]. Mineralocorticoid receptor may mediate activating effects on microglia rather than inhibitory effects. As the major rodent glucocorticoid corticosterone has an affinity for both mineralocorticoid

Two distinct brain macrophage populations in ischemic lesion core










Figure 2. Distinctions between two populations of blood-borne macrophages accumulated in ischemic core lesions of a rat stroke model prepared by tMCAO. CD200⁻/NG2⁺/Iba1⁺ macrophages (BINCs) are the dominant populations occupying the most ischemic core regions. The two populations cannot be classified into M1 and M2 categories. The data come mainly from the data Matsumoto *et al.*^[19] (2015)

receptor and glucocorticoid receptor, it exerts biphasic effects on the microglial proinflammatory actions - activation at low concentrations and inhibition at high concentrations.

A synthetic glucocorticoid dexamethasone (Dex), a specific ligand for glucocorticoid receptor, strongly inhibits LPS-induced NO release and expression of mRNA for IL-1 β and TNF α ^[43]. When primary cortical neurons were cocultured with microglia in the presence of LPS, neurons were degenerated. As this LPS-induced neurodegeneration was prevented by an NOS inhibitor, L-NMMA, the neurodegeneration is caused by NO derived from activated microglia. However, Dex, rather than the NOS inhibitor, suppresses the release of NO more strongly and prevents the degeneration almost completely [Figure 3]. Moreover, Dex increases the expression of mRNA for the neuroprotective factors HGF and IGF-1. The findings indicate that glucocorticoids can induce a neuroprotective microglia phenotype even in the presence of LPS in culture or of DAMPs in the damaged brain parenchyma.

Because of such favorable effects, glucocorticoids have been studied in various animal brain disease models with neuroinflammation. As mentioned, in the rat 6-OHDA-induced PD model, microglia in the SNc are activated in response to degeneration of DA neurons. Chronic administration of glucocorticoids appears to inhibit microglia activation and enhance both the viability of DA neurons and the amount of tyrosine hydroxylase, a rate-limiting enzyme for DA synthesis [Figure 3]. However, glucocorticoids have several adverse effects, including induction of osteoporosis, impaired glucose tolerance, and increased susceptibility to infection. Moreover, glucocorticoids potentially impair cognitive functions by damaging the circuitry of the hippocampus, in which glucocorticoid receptor is expressed abundantly^[86,87]. Nevertheless, as shown in Figure 3, chronic oral administration of glucocorticoid at a rather low dose of






Figure 3. Immunomodulatory effects of a synthetic glucocorticoid Dex on microglia. A: effects of 10 agents on NO release by rat primary microglia (A-a) and rat primary cortical neuron-microglia coculture (A-b) incubated for 48 h with LPS. Nitrite levels in conditioned media were determined. Dex suppressed NO release most effectively. $*P < 0.05$, $**P < 0.01$, $***P < 0.001$, respectively vs. LPS. B: effects of LPS and the agents on LPS-treated neuron-microglia cocultures. B-a: representative immunoblots showing the contents of MAP2 and iNOS proteins in the cocultures. (-LPS) denotes an absence of LPS; B-b: statistical comparison of suppressive effects on LPS-induced iNOS expression; B-c: neuroprotective effects of the 10 agents on the LPS-treated coculture. Strong iNOS expression was correlated with loss of MAP2-immunoreactivity. Only Dex protected neurons significantly from microglial neurotoxicity. Data from four independent cultures are expressed as mean \pm SEM. $*P < 0.05$, $**P < 0.01$, $***P < 0.001$, respectively, vs. LPS. C: microglia were incubated with an indicated agent for 16 h in the absence of LPS. Only Dex increased expression of mRNAs encoding IGF-1 and HGF significantly while suppressing expression of IL-1 β and TNF α mRNAs. $n = 4$, $*P < 0.05$, $**P < 0.01$, $***P < 0.001$, respectively, vs. control. D: dose-dependent ameliorative effects of Dex on a rat 6-OHDA-induced PD model. D-a: representative immunoblots showing dose-dependent effects of Dex on the content of TH, a DA neuron marker; D-b: statistical analyses of the dose-dependent effects of Dex. A dose 0.003 μ g/kg weight of Dex significantly prevented the reduction in TH immunoreactivity. Data from four rats for each dose were expressed as mean \pm SEM. $*P < 0.05$, $**P < 0.01$, $***P < 0.001$, respectively vs. 6-OHDA/Dex 0. The data are from unpublished ones by Wada *et al.* (unpublished data). Dex: dexamethasone; Ald: aldosterone; Imi: imipramine; Flu: fluvoxamine; Ind: indomethacin; Ibu: ibuprofen; Ter: terbutaline; Iso: isoproterenol; Min: minocycline; Tro: troglitazone; LPS: lipopolysaccharide; TH: tyrosine hydroxylase; DA: dopaminergic; PD: Parkinson's disease; HGF: hepatocyte growth factor; IGF-1: insulin-like growth factor 1; 6-OHDA: 6-hydroxydopamine; iNOS: inducible nitric oxide synthase

3 μ g/kg body weight for 6-OHDA-induced PD model rats prevented the reduction in tyrosine hydroxylase immunoreactivity in the SNc to a significant extent. It may be worth administering this dose in a clinical setting.

As mentioned above, microglia play a favorable role by eliminating hyperactive glutamatergic synapses from STN in the PD pathophysiology^[47]. When 6-OHDA-induced PD model rats are administered a high dose of glucocorticoid, the motor deficits are aggravated. This may be one example demonstrating that strong immunosuppressive interventions for activated microglia and macrophages have an aggravating effect, as they have both favorable and unfavorable roles.

Noradrenaline and other cAMP-elevating agents

As shown in Figure 3, noradrenaline (NA) and related agents have significant inhibitory effects on activated microglia. These effects have been attributed mainly to the adrenergic β_2 receptor that increases levels of intracellular cAMP^[73,88]. The β_2 agonist terbutaline strongly suppressed LPS-induced proinflammatory activation of microglia. cAMP-elevating agents, such as phosphodiesterase (PDE) inhibitors also exhibited strong inhibitory effects on LPS-treated microglia^[89]. Rolipram, an inhibitor of the cAMP-selective hydrolase PDE4, may be the PDE inhibitor with the strongest inhibitory effects on rat primary microglia. A cAMP-analogue, 8-bromo cAMP, and forskolin, an activator for adenylate cyclase, also have strong inhibitory effects on microglia *in vitro*. Furthermore, the cAMP-elevating agents inhibit proliferation of microglia *in vitro*^[90]. However, it should be noted that adrenergic α_1 receptor agonists such as phenylephrine can inhibit microglial activation to an extent similar to that of terbutaline^[73,88]. The inhibitory effects of NA cannot be eliminated by an inhibitor for cAMP-dependent protein kinase^[73]. NA, α_1 , and β_2 agonists strongly prevent LPS-induced translocation into the nuclei of NF κ B^[73]. This effect is mediated mainly by suppression of I κ B degradation. NA and the agonists suppressed LPS-induced phosphorylation of STAT1 and expression of IRF1. IRF1 may contribute significantly to microglia activation^[43]. However, the precise molecular mechanisms underlying the inhibitory effects of NA, in addition to the agonists, remain to be identified.

These studies may indicate that BBB-permeable adrenergic agonists are promising agents for suppressing neuroinflammatory processes in pathologic brains. The curative effects of BBB-permeable β_2 agonists, while preventing the proinflammatory nature of microglia in the SNc, have been shown in murine PD models^[91]. In AD pathology, NA neurons in the locus ceruleus, which is the most important NA source in the brain, undergo degeneration leading to microglia activation. This subsequently results in further neuronal degeneration^[92]. However, there are many conflicting studies demonstrating the stimulatory actions of NA and the adrenergic agonists on microglia in the brain^[93].

TGF β 1

TGF β 1 may be the most abundantly released cytokine from microglia and macrophages in severely damaged brains^[27,72]. TGF β 1 suppresses iNOS expression by LPS-treated primary rat microglia almost completely in culture at both mRNA and protein levels. The inhibitory effect of TGF β 1 is as strong as is 100 nM of Dex in culture experiments^[72]. As mentioned above, once incubated with TGF β 1, LPS cannot induce NF κ B translocation into nuclei in microglia. TGF β 1 also inhibits LPS-induced phosphorylation of STAT1 and expression of IRF1. Although TGF β 1 does not increase expression of the M2 markers CD206 and Ym1, it increases expression of the neuroprotective factors HGF and bFGF. When TGF β 1 was injected into ischemic tissues 48 h after MCAO, the immunoreactivity of pIKK in the tissue surrounding the injection site was suppressed markedly in a dose-dependent manner.

Severely damaged brain tissues contain DAMPs that should induce proinflammatory activation of microglia and macrophages. DAMPs bind to TLRs on both microglia and macrophages, resulting in translocation of NF κ B into nuclei. Microglia and macrophages should then display proinflammatory phenotypes expressing proinflammatory mediators. However, there were very few pIKK-bearing or iNOS-expressing microglia/macrophages in the lesion, especially during the subacute phase when TGF β 1 expression is high. Interventions to increase the actions of TGF β 1 for stroke models have been shown to ameliorate the outcomes for severely damaged brains^[94]. Thus, TGF β 1 may ameliorate severe CNS damage through inducing anti-inflammatory phenotypes of microglia and macrophages.

Bromovalerylurea

Bromovalerylurea (BU; C₆H₁₁BrN₂O₂, CAS: 496-67-3) is a hypnotic/sedative that was developed more than a century ago^[95]. It is not prescribed currently because of its weak actions as a hypnotic/sedative compared

with newer agents such as benzodiazepines and also due to dependency. Recently, it has been found to have strong anti-inflammatory effects^[10,43,96,97]. The effects of BU on LPS-treated primary rat microglia are as strong as those of Dex^[43]. Like Dex, BU increases the expression of neuroprotective factors such as HGF or IGF-1 by LPS-treated primary microglia while inhibiting the expression both of proinflammatory factors and of the proinflammatory transcription factors IRF1, IRF7, and IRF8^[43,98]. Although BU does not suppress LPS-induced NF κ B translocation into nuclei, its inhibitory effects are exerted at the transcriptional level. BU inhibited ATP synthesis in mitochondria; this may be related to its anti-inflammatory effects^[10,96]. Moreover, BU inhibits Janus kinase 1 activity, thus suppressing the phosphorylation of STAT1 and the subsequent expression of IRF1^[43].

BU prevented the death of rats with cecum ligation and puncture-induced sepsis by suppressing the proinflammatory activation of peritoneal macrophages^[97]. Oral administration of BU ameliorated PD in rat models while inhibiting the expression of proinflammatory mediators in the ventral midbrain^[43]. When BU is added to the LPS-treated microglia/neuron coculture, it can almost completely inhibit neuronal death by almost completely suppressing the release of NO. In rat TBI models, BU ameliorated neurological deficits considerably while inhibiting expression of chemokine CCL2 and suppressed monocyte infiltration of the lesion^[10]. Furthermore, BU inhibited mitochondrial ROS release by macrophages strongly. On the other hand, BU did not affect the accumulation of favorable macrophages at later time points.

As BU is a hypnotic/sedative, it crosses the BBB easily. It may hold promise as an agent for ameliorating a range of brain diseases and injuries. However, BU is associated with marked dependency, and it has been used for suicide. Nevertheless, recent research has shown that the sleep-inducing effects of BU necessitate a higher dose than that required to produce the anti-inflammatory effects. As a hypnotic, BU increases the total sleeping period while reducing rapid eye movement sleep at a dose of 250 mg/kg for rats (Takeda *et al.*, unpublished observation). Conversely, BU has been administered to sepsis, TBI, and PD model rats at a dose of 50 mg/kg. The findings suggest that BU could be used as a specific anti-inflammatory drug without significant dependency.

Agents that elevate expression of antiapoptotic factors in neurons

Microglia and macrophages can be activated by the presence of damaged neurons because of released DAMPs. Moreover, damaged neurons will reduce the activity of flippase, which is required for asymmetric distribution of phospholipids. Flippase reverses the translocation of PS to the exoplasmic face of the plasma membrane in an ATP-dependent manner. Therefore, even weak neuron damage that reduces intracellular ATP results in translocation of PS to the exoplasmic surface of the plasma membrane. The PS will be recognized by proteins such as MFG-E8 and also by complement C1^[99]. These changes stimulate phagocytic elimination of neurons by microglia and macrophages in a process called phagoptosis^[66]. Therefore, either suppression of the apoptotic changes or mitochondrial damage will suppress the unfavorable activation of microglia and macrophages^[100,101].

A hematopoietic cytokine IL-3 has been shown to increase Bcl-xL expression in neurons in ischemic hippocampi^[102] and in the SNc of PD model rats^[44]. Moreover, GM-CSF also increases Bcl-xL expression^[103]. Simultaneous administration of IL-3 and GM-CSF shows more marked ameliorative effects in TBI and PD model rats. The mixture of the cytokines also inhibited neuronal loss in a rat stroke model prepared by MCAO. Subcutaneous administration of the cytokine mixture to PD model rats inhibited IL-1 β and TNF α expression in the ventral midbrain, whereas the expression of IGF-1 and HGF was increased. The favorable changes in microglia may be mediated by the ameliorated survival of neurons. The cytokines also have specific direct effects on microglia and macrophages^[13,104]. Addition of cytokines to the culture of isolated macrophages (BINC)s from TBI lesions increased HGF and IGF-1 expression, whereas it did not affect IL-1 β expression^[13]. This indicates that inhibition of the proinflammatory nature of the cytokines may be

attributed to ameliorated neuronal survival rather than the direct effects on macrophages and microglia. Thus, an agent that can increase expression of anti-apoptotic factors in neurons would increase the number of favorable microglia and macrophages.

Ginsenosides are natural products isolated from the plant ginseng. Among the ginsenosides, ginsenoside Rb1 and its derivatives strongly ameliorate the outcome of stroke^[105] and TBI^[106] models, while increasing Bcl-xl expression. Along with their direct effects on neurons, ginsenosides have been shown to inhibit the proinflammatory activation of microglia. It remains to be determined whether they inhibit LPS-induced proinflammatory activation of microglia and macrophages^[107,108].

CONCLUSION

Microglia and macrophages have profound effects on the pathophysiological processes of several brain pathologies. They become activated in response to pathological changes of neurons that produce DAMPs and express PS on their surface. Microglia and macrophages can have ameliorative and/or deleterious effects on the CNS depending on the severity of the disease or injuries, time course, and BBB disruption. Even within the same pathology, the cells exert both different effects in completely different ways, as described in the cases of PD and the spinal cord of the peripheral nerve injury model. The development of pharmacological interventions to regulate the response of microglia and macrophages has long been anticipated. However much more research into their responses is required before that goal can be attained.

DECLARATIONS

Acknowledgments

The author would like to thank Enago (www.enago.jp) for the English language review.

Authors' contributions

The author contributed solely to the article.

Availability of data and materials

Not applicable.

Financial support and sponsorship

None.

Conflicts of interest

The author declared that there are no conflicts of interest.

Ethical approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Copyright

© The Author(s) 2020.

REFERENCES

1. Jian Z, Liu R, Zhu X, Smerin D, Zhong Y, et al. The involvement and therapy target of immune cells after ischemic stroke. *Front Immunol* 2019;10:2167.

2. Sevenich L. Brain-resident microglia and blood-borne macrophages orchestrate central nervous system inflammation in neurodegenerative disorders and brain cancer. *Front Immunol* 2018;9:697.
3. Hu X, Leak RK, Shi Y, Suenaga J, Gao Y, et al. Microglial and macrophage polarization - new prospects for brain repair. *Nat Rev Neurol* 2015;11:56-64.
4. Flores JJ, Klebe D, Tang J, Zhang JH. A comprehensive review of therapeutic targets that induce microglia/macrophage-mediated hematoma resolution after germinal matrix hemorrhage. *J Neurosci Res* 2020;98:121-8.
5. Kreutzberg GW. Microglia: a sensor for pathological events in the CNS. *Trends Neurosci* 1996;19:312-8.
6. Kettenmann H, Hanisch UK, Noda M, Verkhratsky A. Physiology of microglia. *Physiol Rev* 2011;91:461-553.
7. Ling EA, Wong WC. The origin and nature of ramified and amoeboid microglia: a historical review and current concepts. *Glia* 1993;7:9-18.
8. Rio-Hortega PD. The microglia. *Lancet* 1939;233:1023-6.
9. Matsumoto H, Kumon Y, Watanabe H, Ohnishi T, Shudou M, et al. Antibodies to CD11b, CD68, and lectin label neutrophils rather than microglia in traumatic and ischemic brain lesions. *J Neurosci Res* 2007;85:994-1009.
10. Abe N, Choudhury ME, Watanabe M, Kawasaki S, Nishihara T, et al. Comparison of the detrimental features of microglia and infiltrated macrophages in traumatic brain injury: a study using a hypnotic bromovalerylurea. *Glia* 2018;66:2158-73.
11. Matsumoto H, Kumon Y, Watanabe H, Ohnishi T, Shudou M, et al. Accumulation of macrophage-like cells expressing NG2 proteoglycan and Iba1 in ischemic core of rat brain after transient middle cerebral artery occlusion. *J Cereb Blood Flow Metab* 2008;28:149-63.
12. Smirkin A, Matsumoto H, Takahashi H, Inoue A, Tagawa M, et al. Iba1(+)/NG2(+) macrophage-like cells expressing a variety of neuroprotective factors ameliorate ischemic damage of the brain. *J Cereb Blood Flow Metab* 2010;30:603-15.
13. Nishihara T, Ochi M, Sugimoto K, Takahashi H, Yano H, et al. Subcutaneous injection containing IL-3 and GM-CSF ameliorates stab wound-induced brain injury in rats. *Exp Neurol* 2011;229:507-16.
14. Taguchi S, Choudhury ME, Miyaniishi K, Nakanishi Y, Kameda K, et al. Aggravating effects of treadmill exercises during the early-onset period in a rat traumatic brain injury model: when should rehabilitation exercises be initiated? *IBRO Rep* 2019;7:82-9.
15. Hu X, Li P, Guo Y, Wang H, Leak RK, et al. Microglia/macrophage polarization dynamics reveal novel mechanism of injury expansion after focal cerebral ischemia. *Stroke* 2012;43:3063-70.
16. Kigerl KA, Gensel JC, Ankeny DP, Alexander JK, Donnelly DJ, et al. Identification of two distinct macrophage subsets with divergent effects causing either neurotoxicity or regeneration in the injured mouse spinal cord. *J Neurosci* 2009;29:13435-44.
17. Cherry JD, Olschowka JA, O'Banion MK. Neuroinflammation and M2 microglia: The good, the bad, and the inflamed. *J Neuroinflammation* 2014;11:98.
18. Ransohoff RM. A polarizing question: do M1 and M2 microglia exist. *Nat Neurosci* 2016;19:987-91.
19. Matsumoto S, Tanaka J, Yano H, Takahashi H, Sugimoto K, et al. CD200+ and CD200- macrophages accumulated in ischemic lesions of rat brain: the two populations cannot be classified as either M1 or M2 macrophages. *J Neuroimmunol* 2015;282:7-20.
20. da Fonseca ACC, Matias D, Garcia C, Amaral R, Geraldo LH, et al. The impact of microglial activation on blood-brain barrier in brain diseases. *Front Cell Neurosci* 2014;8:362.
21. Pun PBL, Lu J, Mochhala S. Involvement of ROS in BBB dysfunction. *Free Radic Res* 2009;43:348-64.
22. Tei N, Tanaka J, Sugimoto K, Nishihara T, Nishioka R, et al. Expression of MCP-1 and fractalkine on endothelial cells and astrocytes may contribute to the invasion and migration of brain macrophages in ischemic rat brain lesions. *J Neurosci Res* 2013;91:681-93.
23. Inoue A, Tanaka J, Takahashi H, Kohno S, Ohue S, et al. Blood vessels expressing CD90 in human and rat brain tumors. *Neuropathology* 2016;36:168-80.
24. Kobayashi K, Yano H, Umakoshi A, Matsumoto S, Mise A, et al. A truncated form of CD200 (CD200S) expressed on glioma cells prolonged survival in a rat glioma model by induction of a dendritic cell-like phenotype in tumor-associated macrophages. *Neoplasia* 2016;18:229-41.
25. Perry VH, Hume DA, Gordon S. Immunohistochemical localization of macrophages and microglia in the adult and developing mouse brain. *Neuroscience* 1985;15:313-26.
26. Perry VH, Gordon S. Macrophages and microglia in the nervous system. *Trends Neurosci* 1988;11:273-7.
27. Sugimoto K, Nishioka R, Ikeda A, Mise A, Takahashi H, et al. Activated microglia in a rat stroke model express NG2 proteoglycan in peri-infarct tissue through the involvement of TGF-beta1. *Glia* 2014;62:185-98.
28. Cronk JC, Filiano AJ, Louveau A, Marin I, Marsh R, et al. Peripherally derived macrophages can engraft the brain independent of irradiation and maintain an identity distinct from microglia. *J Exp Med* 2018;215:1627-47.
29. Butovsky O, Jedrychowski MP, Moore CS, Cialic R, Lanser AJ, et al. Identification of a unique TGF-beta-dependent molecular and functional signature in microglia. *Nat Neurosci* 2014;17:131-43.
30. Gautier EL, Shay T, Miller J, Greter M, Jakubzick C, et al.; Immunological Genome Consortium. Gene-expression profiles and transcriptional regulatory pathways that underlie the identity and diversity of mouse tissue macrophages. *Nat Immunol* 2012;13:1118-28.
31. DePaula-Silva AB, Gorbea C, Doty DJ, Libbey JE, Sanchez JMS, et al. Differential transcriptional profiles identify microglial- and macrophage-specific gene markers expressed during virus-induced neuroinflammation. *J Neuroinflammation* 2019;16:152.
32. Gosselin D, Link VM, Romanoski CE, Fonseca GJ, Eichenfield DZ, et al. Environment drives selection and function of enhancers controlling tissue-specific macrophage identities. *Cell* 2014;159:1327-40.
33. Bennett ML, Bennett FC, Liddelow SA, Ajami B, Zamanian JL, et al. New tools for studying microglia in the mouse and human CNS. *Proc Natl Acad Sci U S A* 2016;113:E1738-46.

34. Satoh J, Kino Y, Asahina N, Takitani M, Miyoshi J, et al. TMEM119 marks a subset of microglia in the human brain. *Neuropathology* 2016;36:39-49.
35. Kanamoto T, Mizuhashi K, Terada K, Minami T, Yoshikawa H, et al. Isolation and characterization of a novel plasma membrane protein, osteoblast induction factor (obif), associated with osteoblast differentiation. *BMC Dev Biol* 2009;9:70.
36. Konishi H, Kobayashi M, Kunisawa T, Imai K, Sayo A, et al. Siglec-H is a microglia-specific marker that discriminates microglia from CNS-associated macrophages and CNS-infiltrating monocytes. *Glia* 2017;65:1927-43.
37. Siew JJ, Chern Y. Microglial lectins in health and neurological diseases. *Front Mol Neurosci* 2018;11:158.
38. Kumar MAS, Peluso M, Chaudhary P, Dhawan J, Beheshti A, et al. Fractionated radiation exposure of rat spinal cords leads to latent neuro-inflammation in brain, cognitive deficits, and alterations in apurinic endonuclease. *PLoS One* 2015;10:e0133016.
39. Choudhury ME, Miyanishi K, Takeda H, Islam A, Matsuoka N, et al. Phagocytic elimination of synapses by microglia during sleep. *Glia* 2020;68:44-59.
40. Yokoyama A, Yang L, Itoh S, Mori K, Tanaka J. Microglia, a potential source of neurons, astrocytes, and oligodendrocytes. *Glia* 2004;45:96-104.
41. Umakoshi K, Choudhury ME, Nishioka R, Matsumoto H, Abe N, et al. B lymphocytopenia and Bregs in a not-to-die murine sepsis model. *Biochem Biophys Res Commun* 2020;523:202-7.
42. McGeer PL, McGeer EG. Glial reactions in Parkinson's disease. *Mov Disord* 2008;23:474-83.
43. Higaki H, Choudhury ME, Kawamoto C, Miyamoto K, Islam A, et al. The hypnotic bromovalerylurea ameliorates 6-hydroxydopamine-induced dopaminergic neuron loss while suppressing expression of interferon regulatory factors by microglia. *Neurochem Int* 2016;99:158-68.
44. Choudhury ME, Sugimoto K, Kubo M, Nagai M, Nomoto M, et al. A cytokine mixture of GM-CSF and IL-3 that induces a neuroprotective phenotype of microglia leading to amelioration of (6-OHDA)-induced Parkinsonism of rats. *Brain Behav* 2011;1:26-43.
45. Banati RB, Gehrmann J, Schubert P, Kreutzberg GW. Cytotoxicity of microglia. *Glia* 1993;7:111-8.
46. Takeuchi H, Jin S, Suzuki H, Doi Y, Liang J, et al. Blockade of microglial glutamate release protects against ischemic brain injury. *Exp Neurol* 2008;214:144-6.
47. Aono H, Choudhury ME, Higaki H, Miyanishi K, Kigami Y, et al. Microglia may compensate for dopaminergic neuron loss in experimental Parkinsonism through selective elimination of glutamatergic synapses from the subthalamic nucleus. *Glia* 2017;65:1833-47.
48. Miyanishi K, Choudhury ME, Watanabe M, Kubo M, Nomoto M, et al. Behavioral tests predicting striatal dopamine level in a rat hemi-Parkinson's disease model. *Neurochem Int* 2019;122:38-46.
49. Kumar V. Toll-like receptors in the pathogenesis of neuroinflammation. *J Neuroimmunol* 2019;332:16-30.
50. Bezard E, Gross CE, Brotchie JM. Presymptomatic compensation in Parkinson's disease is not dopamine-mediated. *Trends Neurosci* 2003;26:215-21.
51. Kurkowska-Jastrzebska I, Litwin T, Joniec I, Ciesielska A, Przybylowski A, et al. Dexamethasone protects against dopaminergic neurons damage in a mouse model of Parkinson's disease. *Int Immunopharmacol* 2004;4:1307-18.
52. Castano A, Herrera AJ, Cano J, Machado A. The degenerative effect of a single intranigral injection of LPS on the dopaminergic system is prevented by dexamethasone, and not mimicked by rh-TNF-alpha, IL-1beta and IFN-gamma. *J Neurochem* 2002;81:150-7.
53. Chen H, Jacobs E, Schwarzschild MA, McCullough ML, Calle EE, et al. Nonsteroidal antiinflammatory drug use and the risk for Parkinson's disease. *Ann Neurol* 2005;58:963-7.
54. Gerhard A, Pavese N, Hotton G, Turkheimer F, Es M, et al. In vivo imaging of microglial activation with [11C](R)-PK11195 PET in idiopathic Parkinson's disease. *Neurobiol Dis* 2006;21:404-12.
55. Gagne JJ, Power MC. Anti-inflammatory drugs and risk of Parkinson disease: a meta-analysis. *Neurology* 2010;74:995-1002.
56. Leal MC, Casabona JC, Puntel M, Pitossi FJ. Interleukin-1beta and tumor necrosis factor-alpha: reliable targets for protective therapies in Parkinson's Disease? *Front Cell Neurosci* 2013;7:53.
57. Moran LB, Graeber MB. The facial nerve axotomy model. *Brain Res Brain Res Rev* 2004;44:154-78.
58. Sudo S, Tanaka J, Toki K, Desaki J, Matsuda S, et al. Neurons induce the activation of microglial cells in vitro. *Exp Neurol* 1998;154:499-510.
59. Kettenmann H, Kirchhoff F, Verkhratsky A. Microglia: new roles for the synaptic stripper. *Neuron* 2013;77:10-8.
60. Streit WJ. Microglia as neuroprotective, immunocompetent cells of the CNS. *Glia* 2002;40:133-9.
61. Nakata T, Kawachi K, Nagashima M, Yasugi T, Izutani H, et al. Transient ischemia-induced paresis and complete paraplegia displayed distinct reactions of microglia and macrophages. *Brain Res* 2011;1420:114-24.
62. Tsuda M, Beggs S, Salter MW, Inoue K. Microglia and intractable chronic pain. *Glia* 2013;61:55-61.
63. Nishihara T, Tanaka J, Sekiya K, Nishikawa Y, Abe N, et al. Chronic constriction injury of the sciatic nerve in rats causes different activation modes of microglia between the anterior and posterior horns of the spinal cord. *Neurochem Int* 2020;134:104672.
64. Sekiya K, Nishihara T, Abe N, Konishi A, Nandate H, et al. Carbon monoxide poisoning-induced delayed encephalopathy accompanies decreased microglial cell numbers: Distinctive pathophysiological features from hypoxemia-induced brain damage. *Brain Res* 2019;1710:22-32.
65. Claus HL, Walberer M, Simard ML, Emig B, Muesken SM, et al. NG2 and NG2-positive cells delineate focal cerebral infarct demarcation in rats. *Neuropathology* 2013;33:30-8.
66. Brown GC, Neher JJ. Microglial phagocytosis of live neurons. *Nat Rev Neurosci* 2014;15:209-16.
67. Neher JJ, Emmrich JV, Fricker M, Mander PK, Thery C, et al. Phagocytosis executes delayed neuronal death after focal brain ischemia. *Proc Natl Acad Sci U S A* 2013;110:E4098-107.

68. Szalay G, Martinecz B, Lenart N, Kornyei Z, Orsolits B, et al. Microglia protect against brain injury and their selective elimination dysregulates neuronal network activity after stroke. *Nat Commun* 2016;7:11499.
69. Jin WN, Shi SX, Li Z, Li M, Wood K, et al. Depletion of microglia exacerbates postischemic inflammation and brain injury. *J Cereb Blood Flow Metab* 2017;37:2224-36.
70. Kim JB, Sig Choi J, Yu YM, Nam K, Piao CS, et al. HMGB1, a novel cytokine-like mediator linking acute neuronal death and delayed neuroinflammation in the postischemic brain. *J Neurosci* 2006;26:6413-21.
71. Shichita T, Hasegawa E, Kimura A, Morita R, Sakaguchi R, et al. Peroxiredoxin family proteins are key initiators of post-ischemic inflammation in the brain. *Nat Med* 2012;18:911-7.
72. Islam A, Choudhury ME, Kigami Y, Utsunomiya R, Matsumoto S, et al. Sustained anti-inflammatory effects of TGF-beta1 on microglia/macrophages. *Biochim Biophys Acta* 2018;1864:721-34.
73. Ishii Y, Yamaizumi A, Kawakami A, Islam A, Choudhury ME, et al. Anti-inflammatory effects of noradrenaline on LPS-treated microglial cells: suppression of NF-kappaB nuclear translocation and subsequent STAT1 phosphorylation. *Neurochem Int* 2015;90:56-66.
74. Yokoyama A, Sakamoto A, Kameda K, Imai Y, Tanaka J. NG2 proteoglycan-expressing microglia as multipotent neural progenitors in normal and pathologic brains. *Glia* 2006;53:754-68.
75. Gao L, Laude K, Cai H. Mitochondrial pathophysiology, reactive oxygen species, and cardiovascular diseases. *Vet Clin North Am Small Anim Pract* 2008;38:137-55.
76. Sinz EH, Kochanek PM, Dixon CE, Clark RS, Carcillo JA, et al. Inducible nitric oxide synthase is an endogenous neuroprotectant after traumatic brain injury in rats and mice. *J Clin Invest* 1999;104:647-56.
77. Hall ED, Wang JA, Miller DM. Relationship of nitric oxide synthase induction to peroxynitrite-mediated oxidative damage during the first week after experimental traumatic brain injury. *Exp Neurol* 2012;238:176-82.
78. Toku K, Tanaka J, Yano H, Desaki J, Zhang B, et al. Microglial cells prevent nitric oxide-induced neuronal apoptosis in vitro. *J Neurosci Res* 1998;53:415-25.
79. Tanaka J, Toku K, Zhang B, Ishihara K, Sakanaka M, et al. Astrocytes prevent neuronal death induced by reactive oxygen and nitrogen species. *Glia* 1999;28:85-96.
80. Matsumoto H, Kumon Y, Watanabe H, Ohnishi T, Takahashi H, et al. Expression of CD200 by macrophage-like cells in ischemic core of rat brain after transient middle cerebral artery occlusion. *Neurosci Lett* 2007;418:44-8.
81. Dudvarski Stankovic N, Teodorczyk M, Ploen R, Zipp F, Schmidt MHH. Microglia-blood vessel interactions: a double-edged sword in brain pathologies. *Acta Neuropathol* 2016;131:347-63.
82. Jiang Z, Jiang JX, Zhang GX. Macrophages: a double-edged sword in experimental autoimmune encephalomyelitis. *Immunol Lett* 2014;160:17-22.
83. Patel AR, Ritzel R, McCullough LD, Liu F. Microglia and ischemic stroke: a double-edged sword. *Int J Physiol Pathophysiol Pharmacol* 2013;5:73-90.
84. Xu H, Wang Z, Li J, Wu H, Peng Y, et al. The polarization states of microglia in TBI: a new paradigm for pharmacological intervention. *Neural Plast* 2017;2017:5405104.
85. Tanaka J, Fujita H, Matsuda S, Toku K, Sakanaka M, et al. Glucocorticoid- and mineralocorticoid receptors in microglial cells: the two receptors mediate differential effects of corticosteroids. *Glia* 1997;20:23-37.
86. Kvarta MD, Bradbrook KE, Dantrassy HM, Bailey AM, Thompson SM. Corticosterone mediates the synaptic and behavioral effects of chronic stress at rat hippocampal temporoammonic synapses. *J Neurophysiol* 2015;114:1713-24.
87. Smith MA. Hippocampal vulnerability to stress and aging: possible role of neurotrophic factors. *Behav Brain Res* 1996;78:25-36.
88. Mori K, Ozaki E, Zhang B, Yang L, Yokoyama A, et al. Effects of norepinephrine on rat cultured microglial cells that express alpha1, alpha2, beta1 and beta2 adrenergic receptors. *Neuropharmacology* 2002;43:1026-34.
89. Zhang B, Yang L, Konishi Y, Maeda N, Sakanaka M, et al. Suppressive effects of phosphodiesterase type IV inhibitors on rat cultured microglial cells: comparison with other types of cAMP-elevating agents. *Neuropharmacology* 2002;42:262-9.
90. Fujita H, Tanaka J, Maeda N, Sakanaka M. Adrenergic agonists suppress the proliferation of microglia through beta 2-adrenergic receptor. *Neurosci Lett* 1998;242:37-40.
91. Qian L, Wu HM, Chen SH, Zhang D, Ali SF, et al. beta2-adrenergic receptor activation prevents rodent dopaminergic neurotoxicity by inhibiting microglia via a novel signaling pathway. *J Immunol* 2011;186:4443-54.
92. Heneka MT, Nadrigny F, Regen T, Martinez-Hernandez A, Dumitrescu-Ozimek L, et al. Locus ceruleus controls Alzheimer's disease pathology by modulating microglial functions through norepinephrine. *Proc Natl Acad Sci U S A* 2010;107:6058-63.
93. Sugama S, Takenouchi T, Hashimoto M, Ohata H, Takenaka Y, et al. Stress-induced microglial activation occurs through beta-adrenergic receptor: noradrenaline as a key neurotransmitter in microglial activation. *J Neuroinflammation* 2019;16:266.
94. Dhandapani KM, Brann DW. Transforming growth factor-beta: a neuroprotective factor in cerebral ischemia. *Cell Biochem Biophys* 2003;39:13-22.
95. Eeckhout Avd. Studien über die hypnotische Wirkung in der Vaierian-säuregruppe. *Arc Exp Path Pharmacol* 1907;57:338-57.
96. Kawasaki S, Abe N, Ohtake F, Islam A, Choudhury ME, et al. Effects of hypnotic bromovalerylurea on microglial BV2 cells. *J Pharmacol Sci* 2017;134:116-23.
97. Kikuchi S, Nishihara T, Kawasaki S, Abe N, Kuwabara J, et al. The ameliorative effects of a hypnotic bromovalerylurea in sepsis. *Biochem Biophys Res Commun* 2015;459:319-26.
98. Tanaka T, Murakami K, Bando Y, Yoshida S. Interferon regulatory factor 7 participates in the M1-like microglial polarization switch. *Glia* 2015;63:595-610.

99. Gyorffy BA, Kun J, Torok G, Bulyaki E, Borhegyi Z, et al. Local apoptotic-like mechanisms underlie complement-mediated synaptic pruning. *Proc Natl Acad Sci U S A* 2018;115:6303-8.
100. Nakatsuka H, Ohta S, Tanaka J, Toku K, Kumon Y, et al. Cytochrome c release from mitochondria to the cytosol was suppressed in the ischemia-tolerance-induced hippocampal CA1 region after 5-min forebrain ischemia in gerbils. *Neurosci Lett* 2000;278:53-6.
101. Nakatsuka H, Ohta S, Tanaka J, Toku K, Kumon Y, et al. Histochemical cytochrome c oxidase activity and caspase-3 in gerbil hippocampal CA1 neurons after transient forebrain ischemia. *Neurosci Lett* 2000;285:127-30.
102. Wen TC, Tanaka J, Peng H, Desaki J, Matsuda S, et al. Interleukin 3 prevents delayed neuronal death in the hippocampal CA1 field. *J Exp Med* 1998;188:635-49.
103. Schabitz WR, Kruger C, Pitzer C, Weber D, Laage R, et al. A neuroprotective function for the hematopoietic protein granulocyte-macrophage colony stimulating factor (GM-CSF). *J Cereb Blood Flow Metab* 2008;28:29-43.
104. Fujita H, Tanaka J, Toku K, Tateishi N, Suzuki Y, et al. Effects of GM-CSF and ordinary supplements on the ramification of microglia in culture: a morphometrical study. *Glia* 1996;18:269-81.
105. Zhang B, Hata R, Zhu P, Sato K, Wen TC, et al. Prevention of ischemic neuronal death by intravenous infusion of a ginseng saponin, ginsenoside Rb(1), that upregulates Bcl-x(L) expression. *J Cereb Blood Flow Metab* 2006;26:708-21.
106. Sakanaka M, Zhu P, Zhang B, Wen TC, Cao F, et al. Intravenous infusion of dihydroginsenoside Rb1 prevents compressive spinal cord injury and ischemic brain damage through upregulation of VEGF and Bcl-XL. *J Neurotrauma* 2007;24:1037-54.
107. Li DW, Zhou FZ, Sun XC, Li SC, Yang JB, et al. Ginsenoside Rb1 protects dopaminergic neurons from inflammatory injury induced by intranigral lipopolysaccharide injection. *Neural Regen Res* 2019;14:1814-22.
108. Gao XQ, Du ZR, Yuan LJ, Zhang WD, Chen L, et al. Ginsenoside Rg1 exerts anti-inflammatory effects via G protein-coupled estrogen receptor in lipopolysaccharide-induced microglia activation. *Front Neurosci* 2019;13:1168.

Review

Open Access



Bilirubin and inflammation in neurodegenerative and other neurological diseases

Sri Jayanti¹, Rita Moretti², Claudio Tiribelli¹, Silvia Gazzin¹

¹Italian Liver Foundation, Centro Studi Fegato, Trieste 34149, Italy.

²Neurology Clinic, Department of Medical, Surgical, and Health Sciences, University of Trieste, Trieste 34139, Italy.

Correspondence to: Dr. Silvia Gazzin, Italian Liver Foundation, Bldg. Q, AREA Science Park, SS14; Km 163,5, Trieste 34149, Italy.
E-mail: silvia.gazzin@fegato.it

How to cite this article: Jayanti S, Moretti R, Tiribelli C, Gazzin S. Bilirubin and inflammation in neurodegenerative and other neurological diseases. *Neuroimmunol Neuroinflammation* 2020;7:92-108. <http://dx.doi.org/10.20517/2347-8659.2019.14>

Received: 12 Nov 2019 **First Decision:** 2 Mar 2020 **Revised:** 14 Mar 2020 **Accepted:** 24 Mar 2020 **Available online:** 11 May 2020

Science Editor: Athanassios P. Kyritsis **Copy Editor:** Jing-Wen Zhang **Production Editor:** Jing Yu

Abstract

Inflammation links neurodegenerative, neuropsychiatric and other neurological diseases (NDs) with acute brain events. It is responsible for the alteration of neurotransmission and circuitry, brain architecture, and cell fate, affecting mood and personality (anxiety, depression and schizophrenia) and behavior (decline in cognitive, motor and speech abilities, altered sleep, fatigue, pain sensitivity and dementia). Inflammation is also a key component in systemic chronic diseases (cardiovascular disease, cancer, diabetes, and metabolic syndrome), in which bilirubin has been demonstrated to improve the diseases by acting as a multi-target antiinflammatory molecule, and where the evaluation of pharmacological modulation of the pigment level as a therapeutic approach has already started. While altered serum bilirubin levels have been reported in ND patients, the potential activity of bilirubin in the brain is vague. This review summarizes the available fragmentary information on the interplay of bilirubin with neuroinflammation, aiming to elucidate the pigment's role in the central nervous system environment.

Keywords: Neuroinflammatory diseases, bilirubin, brain, heme oxygenase, biliverdin reductase, inflammation, homeostasis

INTRODUCTION

Bilirubin (unconjugated - UCB) is the final metabolite of hemoglobin, which is processed in the liver (by conjugation to 1 or 2 molecules of glucuronic acid - CB: conjugated bilirubin) before



© The Author(s) 2020. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, sharing, adaptation, distribution and reproduction in any medium or format, for any purpose, even commercially, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.







Figure 1. Bilirubin & pathological conditions. Representation of the known CNS and not-CNS diseases where TSB is altered. For references of CNS-diseases, see text. For details on not-CNS diseases, see references (according to Gazzin *et al.*^[2] and Wagner *et al.*^[3])

elimination through urine and feces. Increased level of bilirubin (UCB and CB) in the blood is a well-recognized marker of hepatic damage. Recently, slightly elevated serum bilirubin concentration emerged as a biomarker of resistance versus chronic diseases^[1]. Epidemiological data have revealed a reduced prevalence of type 2 diabetes, obesity and metabolic syndrome, certain cancers, and especially cardiovascular diseases and related causes of mortality in Gilbert's syndrome patients, showing mild hyperbilirubinemia^[2,3] [Figure 1]. The effect of the pigment has been demonstrated to be due mainly to the unconjugated (or indirect) moiety of bilirubin (UCB). UCB, especially in its free form [free bilirubin (Bf), the portion of UCB exceeding the serum albumin binding capacity], enters tissues from blood, acting as a powerful antioxidant molecule at nanomolar concentrations, where it is able to counteract 10,000 times higher levels of hydrogen peroxide (H₂O₂)^[4]. This capability has been related to the UCB-biliverdin cycle [Figure 2], which is able to regenerate the pigment consumed by oxidants and acts complementary to cellular glutathione (GSH)^[4-6]. In the last years,



text) and the tissues able to produce *de novo* bilirubin, based on the cited literature (Takeda *et al.*)^[7]

possess the complete enzymatic apparatus necessary for producing UCB themselves [Figure 2]^[7,8]. A more expanded and intricate view has emerged from the discovery of the interplay of UCB (and the enzymes involved in its production and recycling, altogether called the “yellow players”) with cellular functions, signaling pathways, and defense/adaptation mechanisms (not restricted to redox state). Collectively, these findings suggest a greater role for the yellow players in cellular homeostasis and defense against diseases^[2,3]. The interplay of the yellow players with neurological and neurodegenerative diseases is still much less explored. In this article, we review the currently available evidence on the potential roles of bilirubin and other yellow players in neurological disease, with special emphasis on inflammation. We will also address the role of enzyme modulation in bilirubin metabolism, with the goal of increasing the systemic level of bilirubin and the protection it confers.

INFLAMMATION AND NEUROLOGICAL DISEASES: AN OVERVIEW

Chronic neurodegenerative pathologies are currently the most dominant clinical conditions. They comprise multiple sclerosis (MS), amyotrophic lateral sclerosis (ALS), Parkinson's disease (PD), Alzheimer's disease (AD) and vascular dementia. Consistent data show that the inflammatory process can be triggered by protein misfolding or protein accumulation, which are the initial events of a given pathology (i.e., amyloid for AD, tau protein for frontal dementia, alpha synuclein for PD, *etc.*)^[9].

Aging has been associated with a low-grade sterile inflammatory status of the immune system, in which interleukin-6 (IL6), IL1 β and tumor necrosis factor (TNF) are key players, more evident in an unhealthy state. Neuroinflammation can be considered one of the most important etiological factors in age-related neurodegeneration^[10], associated with a reduction in neuron number, a decrease in neuronal arborization, and loss of spines^[11]. What fits particularly in this theory, is the strong evidence that in the aging brain, both macrophages and microglia react with a prolonged and overactive response to stimuli^[12]. This overactivation induces the production of reactive oxygen species and attracts peripheral leukocytes, and both these conditions can activate glial cells^[13]. The activation of glial cells promotes telomere shortening, which can be a contributor in different neurological conditions, such as AD^[14]. Resulting impaired phagocytosis alters the removal of toxic compounds, such as amyloid-beta (A β) and alpha-synuclein (α Syn)^[15]. Microglial activation is believed to be involved in the occurrence of deterioration in various neurodegenerative diseases, such as AD, PD, ALS, and MS^[15]. Microglia are the main immune cells of the central nervous system (CNS), and as the first line of defense, microglia play an important role in the inflammatory reaction^[16]. In AD, microglia are known for their role in cleaning up A β . Meanwhile, in PD, a pathological α Syn aggregation can induce microglial activation and dysfunction^[17]. Microglia are also directly involved in MS by producing cytokines such as tumor necrosis factor- α (TNF- α) and interferon- γ (IFN- γ), superoxide (O $^{2-}$) or nitric oxide (NO), and release of proteolytic and lipolytic enzymes, which can destroy the myelin sheath^[15,18].

Neuroinflammation and mitochondrial dysfunction are common features of chronic neurodegenerative diseases. Both conditions can lead to increased oxidative stress, which leads to excess reactive oxygen species and reactive nitrogen species, resulting in a cascade of events, with injury to polarized bilayers, lipid peroxidation, lysosomal intrinsic activity and autophagy; all these events together, permit self-potentialization of an inflammatory cascade^[19].

Inflammation also plays a role in cerebral small-vessel disease (cSVD), which usually manifests as stroke, cognitive impairment, dementia, physical disability and depression^[20]. Inflammation in cSVD might be explained by the role of endothelial cells in the blood-brain barrier. Endothelial cells communicate with pericytes, astrocytes, microglia, and neural stem cells in the neurovascular unit. Damage to CNS tissues leads to the disruption of tight junctions that link endothelial cells^[21]. It is followed by the infiltration of neutrophils and monocytes, the activation of microglia and astrocytes and the invasion of T and B cells. This inflammatory response is mediated by the nuclear factor κ -light-chain-enhancer of activated B cells (NF κ B) pathway and strongly associated with neuroinflammation in the acute phase of various vascular injuries such as stroke, diabetic retinopathy, and AD^[22,23].

Bilirubin has been reported to act on all the above-mentioned molecular mechanisms. The following paragraphs review the state-of-the-art of what it is known about bilirubin and the brain.

BILIRUBIN: A MARKER FOR NEUROLOGICAL DISEASES

Cumulative clinical evidence demonstrates the alteration of total serum bilirubin (TSB) level in neuroinflammatory diseases, including schizophrenia, MS, PD, AD, ALS, stroke, diabetic retinopathy, *etc.* (for details, [Table 1](#)). For obvious reasons (limited CNS sample availability, especially in the early stages of diseases), few data on the potential molecular role of the pigment in these pathologies are available, and cause-effect studies are possible only by using experimental models.

Schizophrenia

Multiple contrasting studies exist on the potential correlation between the serum level of UCB and schizophrenia. Studies have reported an increase in UCB level, as in Gilbert's syndrome in schizophrenia compared to other psychiatric diseases (affective disorder and neuropsychosis) and healthy controls^[24].

Table 1. TSB level in neurological diseases

Dis.	Study design	# Subjects Dis/Ctrl	TSB Dis/Ctrl	CB Dis/Ctrl	UCB Dis/Ctrl	P	Ref.
AD	CC	101/101	10.26/15.39			< 0.001	[119]
AD	CC	pAD: 12/MC-AD: 12/HS: 12	4.62/3.93/12.48			< 0.05	[120]
ALS	CC	Short duration: 19/ long duration: 11	12.31/7.70			0.011	[115]
DR	CC	67/diabetic no DR: 35	10.1/15.1			< 0.01	[84]
DEM	CS /	Cogn Imp: 31/no Cogn Imp: 33	11.3/13.5			0.020	[117]
	CC	Cog. Imp. 31/40	11.3/15.0			0.003	
MS	CC	133/88	11.08/16.47	2.20/4.26	8.88/2.2	0.001	[87]
NMO	CC	67/98	12.25/16.15		9.17/13.22	< 0.001	[80]
NO	CC	42/48	11.8/15.5	2.7/4.1	8.7/11.4	< 0.01	[79]
PD	CC	420/435	9.57/7.70			< 0.001	[45]
PD	CS/CC	drug naïve PD: 75/75	12.65/8.72			< 0.001	[46]
PD	CC	LD-PD: 162/untr-PD: 93/ HS: 224	12.31/10.94/10.26			< 0.001	[48]
PD	CC	425/460	12.09/12.3	3.73/3.15	8.36/9.15	<0.05	[47]
SCI	CC	343/2522	8.3/10.7			< 0.001	[88]
SCZ	CC	34/114	9.74/11.14			0.04	[41]
SCZ	CC	72/65	6.5/12.2			10 ⁻¹³	[42]
SCZ vs. BD	CC	71/BD: 69	10.62/8.04*		7.87/5.47**	*0.02 **0.03	[26]
SCZ vs. BD	CS	50/BD: 43	8.89/7.01 ^a		6.84/4.45 ^b	^a 0.027 ^b 0.004	[27]
SCZ vs. BD	CS	44/BD: 56			7.01/4.95	< 0.0001	[25]
SCZ vs. SAD	CS	44/SAD: 99			7.01/5.81	< 0.03	

DDis: specification of the disease; Ctrl: specification of the controls or group of comparison (for detail see each study). When unspecified = healthy population; TSB: total serum bilirubin (μmol/L); CB: conjugated bilirubin or direct bilirubin (DB, μmol/L); UCB: unconjugated bilirubin or indirect bilirubin (IB, μmol/L); AD: alzheimer disease; ALS: amyotrophic lateral sclerosis; BD: bipolar disorders; CIS: clinically isolated syndrome; DEM: dementia, Cogn.Imp: subject with DEM and cognitive impairment; No Cogn.Imp: subject with DEM but not cognitive impairment; DR: diabetic retinopathy; HS: healthy subjects; MC-AD: mild cognitive Alzheimer disease; MS: multiple sclerosis; NO: neuritis optica; NMO: neuromyelitis optica; pAD: probable Alzheimer disease; PD: parkinson disease; LD-PD: L-Dopa treated PD patients; Untr.PD: untreated PD patients; SAD: schizoaffective disorder; SCI: silent cerebral infarct; SCZ: schizophrenia; CC: case control; CS: cross-sectional. ^aP value for TSB, ^bP value for UCB

More interestingly, after antipsychotic treatment, UCB decreases in 80% of subjects. Gama Marques *et al.*^[25] also reported that mean UCB levels are clearly higher in patients with schizophrenia than patients with schizoaffective disorder and bipolar disorder. Additionally, retrospective and recent prospective studies indicate that serum UCB level in patients with schizophrenia is higher than those with bipolar affective disorder (1.40- and 1.53-fold, respectively)^[26,27]. Notably, all these studies included only subjects with normal liver enzymes and no pre-existing liver disease or other confounding factors, further emphasizing the pivotal role of UCB increase. UCB can be viewed as a potential biomarker to distinguish schizophrenia from other psychiatric disorders^[28].

UCB level also shows a correlation with schizophrenia symptoms. Patients with hyperbilirubinemia have shown significantly higher scores on the positive and general psychiatric subscales of the PANSS (Positive and Negative Syndrome Scale)^[24]. In concordance with the previous finding, UCB elevation has been found frequently in psychotic episodes of schizophrenia^[25,29]. Radhakrishnan *et al.*^[26] reported that UCB levels are higher in paranoid schizophrenia than non-paranoid schizophrenia. Meanwhile, a recent study by Pradeep *et al.*^[27] found no association between serum UCB levels with the severity of psychopathology in schizophrenia subjects.

Schizophrenic subjects with unconjugated hyperbilirubinemia show a significant enlargement of cerebral ventricles^[30] and abnormalities of brain metabolism compared to both normobilirubinemic schizophrenia patients and normal controls^[31]. Neuroinflammation in schizophrenia is characterized by increased serum concentrations of proinflammatory cytokines, including IL1β, IL6, and transforming growth factor-β

(TGFβ) and microglial activation^[32,33]. This inflammatory picture is similar to that observed in kernicterus spectrum disorders, the neurological sequel due to severe neonatal hyperbilirubinemia^[34]. Dalman and Cullberg^[35] reported that neonates experiencing severe neonatal hyperbilirubinemia (TSB > 15 mg/dL) might later have an increased frequency of mental disorders. This clinical hypothesis is supported by the finding in the Gunn rat, the animal model for severe neonatal hyperbilirubinemia^[36], showing severe hyperbilirubinemia in the first weeks of life^[37]. The Gunn rat displays a stereotypical behaviour, a supposedly typical symptom of schizophrenia, associated with microglia activation (indicative of inflammation), decreased ribosomal protein synthesis activity in neuronal cells, decreased neurogenesis, and increased apoptosis^[38,39].

On the other hand, multiple studies have reported lower TSB levels among schizophrenia patients compared to patients suffering from some other psychiatric disorder and healthy controls^[40-42]. To reconcile these contradictory findings, Vitek *et al.*^[42] evaluated the correlation of bilirubin level with variability of the promoter of the gene for *UGT1A1* (uridine-diphospho-glucuronosyltransferase 1A1) (responsible for hepatic conjugation of UCB) between schizophrenia patients and controls. They noticed that an increase in bilirubin of 1 μM (0.06 mg/dL) could reduce the odds for schizophrenia status up to 19%.

Notably, UGTs participate in the metabolism of dopamine^[26], a critical neurotransmitter, the loss of which is responsible for motor PD, and dopaminergic over activity has recently been suspected responsible for psychosis in schizophrenia^[43]. An additional potential link between these two pathologies might be heme oxygenase-1 (HO-1). The overexpression of this enzyme in the astrocytes of glial fibrillary acidic protein-heme oxygenase-1 (GFAP-HMOX1) transgenic mice resulted in increased subcortical oxidative imbalance with the induction of mitochondrial damage and autophagy, augmented dopamine and serotonin levels in the basal ganglia, and reduced dopamine 1 receptor (D1) in the nucleus accumbens, and enzyme overexpression caused degeneration of axons in the hippocampus and hyperkinetic behavior^[44]. However, there is still no clear evidence of HO-1 alteration in the clinical setting for schizophrenia.

PD

PD is another neurological disease linked to UCB. A study comparing 420 PD patients and 435 healthy control showed not only a significant increase in bilirubin in PD patients but also a negative correlation between bilirubin level and progression from a less to more severe staging of the disease^[45]. Complementary with this study, Moccia *et al.*^[46] also reported higher TSB concentration in drug-naive PD patients compared to controls, and Qin *et al.*^[47] also observed an upregulation of levels of direct bilirubin (conjugated bilirubin, in clinical terminology), accompanied by reduced UCB (indirect bilirubin) levels in PD patients compared to the healthy group. Notably, L-DOPA treatment (the most used therapeutic approach to PD, able to improve symptomatology temporarily), has been found to increase TSB by about 20%^[48]. Because L-DOPA may increase the oxidative stress causing dopaminergic neuron loss in PD, and since bilirubin is a well-known antioxidant, an increased bilirubin level has been interpreted as a possible protective response to the disease^[48].

As noted in the Introduction, almost every cell in the body possesses the full enzymatic equipment for producing UCB itself^[7] [Figure 2]. Modulation of the enzymes responsible for UCB production in the brain of subjects with neurological/neurodegenerative diseases has been reported, including PD patients. An increased HO-1 signal has been detected in reactive astrocytes and affected dopaminergic neurons showing Lewy bodies^[49]. Although, the authors suggested that the HO-1 induction was uniquely due to its antioxidant properties, it is clear now that the elevation of HO-1 levels also represents an attempt to downregulate inflammation^[50].

A direct proof of the protective effect of HO-1 induction has been obtained in experimental models. The injection of adenovirus containing human heme oxygenase-1 gene (*Hmox1*) into the substantia nigra of

methyl-4-phenylpyridinium (MPP+) treated rats (a model for PD) revealed that the overexpression of *Hmox1* protects dopaminergic neurons by reducing the expression of TNF- α and IL1 β , and increasing brain-derived neurotrophic factor (BDNF) and glial cell line-derived neurotrophic factor (GDNF) expression^[51]. Both growth factors, BDNF and GDNF, are able to prevent injury to dopaminergic neurons and improve the behavioural deficits in PD^[52-54]. A recent work based on organotypic cultures of substantia nigra (challenged with rotenone to induce PD) demonstrated that inflammation and redox imbalance are early and simultaneous triggers for dopaminergic neuron loss^[55]. In this study, *Hmox1* upregulation occurs only at the very first phases of neurodegeneration (3 hours after rotenone challenging, a time which represents the pre-diagnosis stage in human PD based on dopaminergic neuron loss). Similarly, *Bdnf* expression initially increased but then rapidly declined below the control level accompanying dopaminergic neuron demise. The modulation trend of both markers (*Hmox1* and *Bdnf*) was interpreted as a failed tentative reaction to the insult^[55], supporting the hypothesis of the interplay between L-DOPA and bilirubin (see above). This interpretation may also agree with a recent finding of Song *et al.*^[56], who reported that the expression levels of HO-1 in the saliva of PD patients with early-stage (stage 1) based on the Hoehn and Yahr scale were significantly higher compared to control subjects or PD patients at stage 3 (advanced PD). The authors proposed an additional value of HO-1 as a potential biomarker in idiopathic PD patients. Notably, hyperactivation of HO-1 may exacerbate oxidative stress by the deposition of iron (Fe), one of its products [Figure 2], a phenomena frequently observed in neurological lesions and known to worsen the disease^[49]. This aspect is discussed in detail later on in the review.

Contrary to PD and schizophrenia, other neurological conditions have been associated with a reduced TSB level.

Multiple sclerosis

MS is a chronic inflammatory disease characterized by the destruction of myelin in the brain and spinal cord likely due to loss of immune system tolerance to myelin^[57]. TSB levels have been reported to be lower (vs. healthy controls) also in MS subjects with clinically isolated syndrome (predominately by neuroinflammation) and relapsing-remitting multiple sclerosis (predominately by neurodegenerative disorder), where predominate neurodegenerative disorders^[58]. Interestingly, bilirubin levels have been shown to be significantly lower in relapsing-remitting multiple sclerosis compared to clinically isolated syndrome patients^[59], suggesting a relationship between increased disease severity and decreased TSB. This hypothesis seems to be confirmed by the study of Ljubisavljevic *et al.*^[59] who found a significant correlation between higher TSB and lower disability status, fewer MRI lesions, and shorter disease duration in both study groups. In the brain tissue from MS patients, enhanced nitrotyrosine staining, one of the markers of oxidative damage^[60], was found in demyelinated regions, specifically in hypertrophic astrocytes and foamy macrophages on inflammatory lesions^[61]. The damaged sites also showed an upregulation of antioxidant enzymes, including HO-1, compared to normal-appearing white matter and white matter tissue from control brains with no neurological disease, and HO-1 immunoreactivity was particularly confined to microglia^[61]. These findings emphasize the close interaction between oxidative stress and inflammation in MS.

The effects of bilirubin on MS have also been studied in the animal model for the disease, the so-called experimental autoimmune encephalomyelitis (EAE) model, where the infiltration of lymphocytes, the activation of CD4⁺ T cells, and death of oligodendrocytes lead to the destruction of myelin sheaths^[62,63]. Figure 3 illustrates the most relevant mechanisms of action of bilirubin in EAE, one of the best-known models unravelling the interplay between MS and bilirubin. Bilirubin administration in the EAE model effectively prevented both acute and chronic EAE, even better than did glucocorticoid treatment, the most commonly used therapy for MS^[64,65].




Figure 3. Multiple ways of bilirubin as immunomodulator in neuroinflammation disease. Ahr: aryl hydrocarbon receptor; APC: antigen presenting cell; BDNF: brain-derived neurotrophic factor; DC: dendritic cell; FcR: Fc receptor; GDNF: glial cell line-derived neurotrophic factor; IFN γ : Interferon γ ; IL: interleukin; MHC II: major histocompatibility complex; PI3K/Akt: phosphatidylinositol 3-kinase; Th: T helper; TCR: T cell receptor

Histological analyses demonstrated that bilirubin interfered with the infiltration of inflammatory cells into the CNS by protecting the blood-brain barrier from free radical-induced permeability changes^[64], stressing the intimal connection between redox stress and inflammation in CNS diseases. Further study supported the antiinflammatory potential of bilirubin in the EAE model^[64]. *In vitro* experiments using spleen-harvested CD4⁺ T cells showed that bilirubin at non-apoptotic concentrations (20-150 μ M) inhibits CD4⁺ T cell proliferation through various mechanisms. It suppresses the production of T helper-1 (Th1) cytokines (IL2 and IFN γ) and Th2 cytokines (IL4 and IL10), reduces costimulatory molecule activity (CD28 on CD4⁺ T cell, the co-receptor B7-1 activity in macrophages and dendritic cells), inhibits NF κ B activation, which is a key transcription factor involved in T cell receptor-mediated signaling, and downregulates inducible MHC (major histocompatibility complex) class II expression. Bilirubin effectively downregulates EAE in SJL/J mice as confirmed by the reduction of the proliferation capacity of CD4⁺ T cell. Meanwhile, the reduction of endogenous bilirubin synthesis by zinc-protoporphyrin, a specific inhibitor of the bilirubin producing-enzyme HO-1, dramatically exacerbates this disease^[66]. In contrast, induction of HO-1 by cobalt protoporphyrin IX (CoPPIX) inhibits EAE effectively^[67].

According to the previous data, it seems beneficial to increase bilirubin synthesis through HO-1 induction. Several clinically used drugs have been reported to induce HO-1, among them are nonsteroidal antiinflammatory drugs (e.g., coxibs, acetylsalicylic acid) and hypolipidemic agents (e.g., niacin, fibrates, statins)^[68]. Atorvastatin and rosuvastatin treatment in mice demonstrated protection by increasing not only plasma bilirubin concentrations (up to 70%) but also cardiac tissue bilirubin content (up to 119%)^[69]. On the contrary, HO-1 induction in astroglia promotes oxidative mitochondrial membrane damage, iron sequestration, and mitophagy (macroautophagy)^[56]. These reasons then increase the doubt of bilirubin synthesis through HO-1 induction as a good strategy to counteract the neuroinflammatory process.

In another part of inflammation signalling pathways, both UCB and biliverdin are known as activators of aryl hydrocarbon receptor (Ahr), a ligand-activated transcription factor that plays critical modulatory roles in various immune cells during innate and adaptive immune responses^[2,70,71]. Ahr has critical roles

in MS by modulating IL17 producing T-helper (Th17) cells, and regulatory T cells^[72-74], as well as B cells, macrophages and dendritic cells. In macrophages, AhR regulates IL1 β production and also IL6, IL12 and TNF α expression^[73,75]. AhR also inhibits the transcriptional activity of NF κ B in stimulated macrophages^[76]. Meanwhile, in dendritic cells, AhR mediates the generation of T regulatory cells and Th17 cells from naive T cells and also IL10 production^[77]. Notably, the injection of low doses of UCB (20 μ g/kg body weight, injected into C57Bl/6 mice) increased the population of regulatory T cells by more than 50% compared to controls, prolonging graft survival^[78]. Regulatory T cells are also inducible by HO-1 activity, CO, and UCB^[78] and all “yellow players” [Figure 2].

Optic neuritis and neuromyelitis optica

Reduced TSB has been noticed in both optic neuritis, an acute inflammatory demyelinating disorder of the optic nerve, and neuromyelitis optica, also known as Devic disease, a severe autoimmune demyelinating disease that selectively affects the optic nerve and spinal cord^[79-82]. Although the main focus was on the potential antioxidant role of bilirubin, suggesting that the low TSB is a result of the overconsumption of bilirubin due to oxidative stress, immunohistochemistry of damaged nerves is similar as in MS^[83], with an evident inflammatory component. Thus, the interplay between bilirubin and inflammation described in MS also applies to optic neuritis and neuromyelitis optica. Further studies are needed.

Diabetic retinopathy and stroke

TSB levels have been found to be decreased also in patients with diabetes mellitus and diabetic retinopathy vs. subjects with diabetes mellitus without retinopathy, suggesting TSB as a biomarker of diabetic retinopathy^[84]. In the pathogenesis of diabetic retinopathy, particularly in the non-proliferative type, hyperglycemia exposure causes the alteration of retinal microvasculature with pericyte loss^[85], endothelial apoptosis and endothelial thickening of the basement membrane, which further lead to capillary occlusion and ischemia^[86].

Similarly, damage and even destruction of the cerebral vasculature are landmarks of stroke and related vascular events^[87]. In a large study conducted by Li *et al.*^[88], 343 subjects with silent cerebral infarct presented with lower TSB levels. In another study, patients with deep white matter lesions (DWMLs) had a decreased TSB level compared to non-DWMLs subjects. Again, the low and intermediate bilirubin groups showed a higher prevalence of severe DWMLs than did the group with higher bilirubin levels^[89]. Notably, DWMLs are a recognized predictor for the development of impaired cognitive function and stroke^[90,91].

A recent study also explored the association of UCB and intracranial atherosclerosis, which was found in approximately 50% of patients with transient ischemic attack and up to 47% of ischemic stroke patients in Asia^[92-94]. In this study, UCB was significantly negatively associated with intracranial atherosclerosis. The odds of intracranial atherosclerosis was 0.67-fold lower in participants in the high UCB concentration group (≥ 10.10 mMol/L) when compared with those in the low UCB concentration group^[94]. An increased bilirubin level has been proposed as a novel independent predictor for hemorrhagic transformation and symptomatic intracranial hemorrhage after mechanical thrombectomy^[95].

The role of bilirubin in all previously mentioned conditions might be explained by a large number of studies suggesting bilirubin as a protector against microvascular complications. Bilirubin exhibits potent antiinflammatory effects [via HO-1, nuclear factor erythroid 2-related factor 2 (Nrf2) and NFκB] by the inhibition of monocyte transmigration [through a decrease in TNFα-induced monocyte chemoattractant protein-1 (MCP-1) secretion], reducing endothelial vascular cell adhesion molecule-1 (VCAM-1) expression, and improving endothelial cell dysfunction and hyperproliferation after damage^[96-99]. UCB also appears to affect the immune system by inhibiting the activation of the complement cascade^[100], and also by modulating the phagocytic and antigen-presenting function of macrophages (by changing the expression of Fc receptors)^[101].

The potential protective role of HO-1 induction has been demonstrated in several studies. HO-1 increases early (24 h) after brain trauma^[102], and in intracerebral haemorrhage models, after infiltration of the brain by blood components activating the inflammatory response^[103]. HO-1 activity is observed in both endothelial cells and microglia surrounding the haematoma site. The protective effects of HO-1 in neurons was studied by Orozco-Ibarra *et al.*^[104], who revealed that the upregulation of HO-1 prevented the death of cerebellar granule neurons due to mitochondrial toxicity. Further investigation found that the HO-1 products, bilirubin and CORM2 (carbon monoxide releasing molecule), were involved in preventing cell death.

HO-1 has neuroprotective effects by regulating the phosphatidylinositol 3-kinase (PI3K/AKT) signalling pathway and by reducing apoptosis in rats with cerebral haemorrhage^[105]. Feng *et al.*^[106] showed that the PI3K/AKT and extracellular signal regulated kinase pathways are involved in oleanolic acid induced HO-1 expression by activating Nrf2 in vascular smooth muscle cells. As one of the signalling pathways for cell survival, the PI3K/AKT signal transduction pathway plays an important role in cell proliferation and differentiation and inhibition of neuronal apoptosis^[107]. Therefore, HO-1 may protect the nerves of rats with cerebral haemorrhage by regulating the PI3K/AKT signalling pathway.

A second isoform of HO exists in the brain, called heme oxygenase-2 (HO-2). Considered constitutive, it looks like HO-2 in the brain plays the vital function of maintaining adequate levels of UCB to guarantee normal cellular homeostasis, participating in brain protection^[108]. The protective activity of HO-2 has been reported in intracerebral haemorrhage models, where heme oxygenase-2 gene (*Hmox2*) deletion led to greater brain injury volumes and neurological deficits than in wild-type mice after intracerebral haemorrhage^[109]. In brain cultures, Doré *et al.*^[108] expanded our knowledge on the role of HO-2 and bilirubin as neuroprotective factors by showing increased neuronal death in cerebellar granule cultures of *Hmox2* knockout mice (*Hmox2*^{-/-}). On the contrary, *Hmox2* transfection in human embryonic kidney 293 (HEK293) cells rescued cells from apoptotic death. In another study, the induction of HO-2 activity by phorbol esters enhanced the production of bilirubin, which protected primary hippocampal and cortical neuronal cultures from the neurotoxicity of H₂O₂. Of note, HO-2 was immunolocalised in neurons both before and after traumatic brain injury, whereas HO-1 was highly upregulated in glia only after traumatic brain injury. Cell loss was significantly greater in *Hmox2*^{-/-} mice in areas including the cortex, hippocampus and lateral dorsal thalamus^[110].

Biliverdin (BV), another yellow player, also showed neuroprotective effects by ameliorating cerebral reperfusion injury in rats most probably via its antiinflammatory activity^[111]. Two pathways are known to be involved in the antiinflammatory mechanism of BV: by activating the nitric oxide-dependent biliverdin reductase, BV reduces the expression of toll-like receptor-4 (TLR-4) in murine macrophages^[112], and BV regulates the expression of complement C5a receptor^[113]. Furthermore, a study by Zou *et al.*^[111] found the downregulation of miR-204-5p and its target gene, ETS protooncogene 1 (*Ets1*), in cerebral ischemia-reperfusion injury rats following BV administration. *Ets1* is known to be responsible for inducing proinflammatory Th1 type response and causing neuronal death^[114]. Therefore, BV may play a crucial role in preventing injury in stroke by interfering in miRNAs levels.

Additional neurological conditions underpinned by reduced TSB levels

ALS

In ALS, the reduction of TSB levels correlates with both the clinical state and disease duration. Patients with long-lasting ALS (where motor neuron degeneration is noticed) have lower TSB levels than do patients with a shorter duration^[115].

Seasonal affective disorder

Nocturnal plasma bilirubin levels were evaluated in individuals with unipolar recurrent winter seasonal major depressive disorder and age-/gender-matched controls. Lower bilirubin levels were found in patients with seasonal affective disorder. The reduction in bilirubin level in this study was proposed as a vulnerable risk for depression, although only an associative and not a causative link was provided^[116].

Dementia

A significant reduction in TSB levels was reported in patients with cognitive impairment compared to those with normal cognitive function, although no significant correlation between bilirubin and disease duration was found^[117]. In a further study by Baierle *et al.*^[118], the authors showed that the significant increase in the proinflammatory cytokines IL1 β and TNF α correlated with the increase in oxidative stress and inversely associated with low cognitive performance in elderly patients.

AD

As in the majority of neurological conditions, TSB levels are also decreased in AD^[119]. The lower concentration of bilirubin is supposed to have a role in the transition from mild cognitive impairment to clear AD^[120]. Similarly, reduced *Hmox1* mRNA^[121] and protein^[122] levels were detected in the serum of full-blown AD subjects, and proposed to correlate with cognitive decline^[120]. Notably, the diagnosis of AD and cognitive decline is usually a late event and performed after the pathological alteration has already existed for some time. In the early stages, the brain tries to protect itself from the oxidative and inflammatory insult of AD by increasing the endogenous production of UCB. Both HO-1 and biliverdin reductase A (BLVR-A), the two key enzymes in biliverdin and bilirubin production [Figure 2], have been reported to be upregulated in the brain of AD patients^[103,123]. However, the upregulation of HO-1 will also cause Fe deposition in the brain, which worsens and accelerates disease progression by enhancing the redox imbalance in ongoing AD lesions^[49,124-126]. It should be recalled that the brain possesses very low antioxidant capacity compared to other organs, being much lower in neonatal life^[127,128] and in the elderly^[129,130]. Thus, the enhanced pro-oxidant milieu due to HO-1 hyperactivation leads to an increased oxidative and nitrosative post-translational modification of cellular enzymes, with their consequent inactivation^[131]. One of the targets of this mechanism is BLVRA, and thus, UCB production will be stopped^[123].

The chains of events described here, leading to the disruption of UCB-mediated protection and contributing to the progression of neurological damage fits well with what we described in PD.

IS THE BRAIN DIFFERENT FROM THE REST OF THE BODY?

Differently from extra-CNS diseases involving oxidative imbalance and inflammation, where a positive correlation between higher TSB level (Gilbert syndrome-like) and reduced disease incidence has been frequently found^[2,3,68], the most readily evident finding in this review is that brain diseases have lower TSB levels in common. Two major explanation have been given: (1) lower TSB levels in subjects with neurological disease may reflect the consumption of UCB due to oxidative stress, although it seems unlikely that the local redox imbalance occurring in the brain may affect the systemic levels of bilirubin; and (2) individuals with lower TSB, thus a lower systemic antioxidant status, might be more vulnerable to oxidative stress^[118] and related diseases (both in the brain and extra-CNS organs) [Figure 1].

Experimental evidence strongly supports the notion that an increased concentration of bilirubin in the CNS exerts antiinflammatory and antioxidant effects. On the other hand, it should be recalled that hyperactivation of HO-1 in the CNS, reported to be a tentative reaction against brain insults, may enhance the damage^[68,104,131-148]. This specificity of the CNS has to be taken into account in the growing research aimed at modulating the bilirubin players to increase levels of bilirubin and the protection it confers.

CONCLUSION

Studies assessing clearly the molecular events correlating bilirubin and neurological damage are still lacking, and many conclusions are based on clinical series. Understanding the role of bilirubin and all the other yellow players in the development and progression of the different neurological disorders may help in deciphering the efficacy of the modulation of bilirubin level to prevent CNS diseases. However, due to the side effects of the drugs inducing HO-1 activity, caution must be taken before their recommendation. More intriguing and promising is the *in situ* CNS elevation of UCB level achievable by the development of molecules modulating BLVR. The future will hopefully provide the much-needed answer due to the medical and social burden of neurological disorders.

DECLARATIONS

Authors' contributions

Made substantial contributions to the search of the literature, the writing of the manuscript, and the revision and discussion of the contents of the paper: Jayanti S, Moretti R, Tiribelli C, Gazzin S

Availability of data and materials

Not applicable.

Financial support and sponsorship

Jayanti S was supported in part by a fellowship from the Lembaga Pengelola Dana Pendidikan of Ministry of Finance of Indonesia and an internal grant from the Italian Liver Foundation. The funders had no role in data analysis and interpretation and also writing of the manuscript.

Conflicts of interest

All authors declare that there are no conflicts of interest.

Ethical approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Copyright

© The Author(s) 2020.

REFERENCES

1. Gazzin S, Masutti F, Vitek L, Tiribelli C. The molecular basis of jaundice: an old symptom revisited. *Liver Int* 2017;37:1094-102.
2. Gazzin S, Vitek L, Watchko J, Shapiro SM, Tiribelli C. A novel perspective on the biology of bilirubin in health and disease. *Trends Mol Med* 2016;22:758-68.
3. Wagner KH, Wallner M, Mölzer C, Gazzin S, Bulmer AC, et al. Looking to the horizon: the role of bilirubin in the development and prevention of age-related chronic diseases. *Clin Sci (Lond)* 2015;129:1-25.
4. Baranano DE, Rao M, Ferris CD, Snyder SH. Biliverdin reductase: a major physiologic cytoprotectant. *Proc Natl Acad Sci U S A* 2002;99:16093-8.
5. Sedlak TW, Snyder SH. Bilirubin benefits: cellular protection by a biliverdin reductase antioxidant cycle. *Pediatrics* 2004;113:1776-82.
6. Sedlak TW, Saleh M, Higginson DS, Paul BD, Juluri KR, et al. Bilirubin and glutathione have complementary antioxidant and cytoprotective roles. *Proc Natl Acad Sci U S A* 2009;106:5171-6.
7. Takeda TA, Mu A, Tai TT, Kitajima S, Taketani S. Continuous de novo biosynthesis of haem and its rapid turnover to bilirubin are necessary for cytoprotection against cell damage. *Sci Rep* 2015;5:10488.
8. Yuan X, Rietzschel N, Kwon H, Walter Nuno AB, Hanna DA, et al. Regulation of intracellular heme trafficking revealed by subcellular reporters. *Proc Natl Acad Sci U S A* 2016;113:E5144-52.
9. Chitnis T, Weiner HL. CNS inflammation and neurodegeneration. *J Clin Invest* 2017;127:3577-87.

10. Stephenson J, Nutma E, van der Valk P, Amor S. Inflammation in CNS neurodegenerative diseases. *Immunology* 2018;154:204-19.
11. Yankner BA, Lu T, Loerch P. The aging brain. *Annu Rev Pathol* 2008;3:41-66.
12. Rawji KS, Mishra MK, Michaels NJ, Rivest S, Stys PK, et al. Immunosenescence of microglia and macrophages: impact on the ageing central nervous system. *Brain* 2016;139:653-61.
13. Chinta SJ, Woods G, Rane A, Demaria M, Campisi J, et al. Cellular senescence and the aging brain. *Exp Gerontol* 2015;68:3-7.
14. Flanary BE, Sammons NW, Nguyen C, Walker D, Streit WJ. Evidence that aging and amyloid promote microglial cell senescence. *Rejuvenation Res* 2007;10:61-74.
15. Xu L, He D, Bai Y. Microglia-Mediated Inflammation and Neurodegenerative Disease. *Mol Neurobiol* 2016;53:6709-15.
16. Streit WJ. Microglia as neuroprotective, immunocompetent cells of the CNS. *Glia* 2002;40:133-9.
17. Zhang W, Wang T, Pei Z, Miller DS, Wu X, et al. Aggregated alpha-synuclein activates microglia: a process leading to disease progression in Parkinson's disease. *FASEB J* 2005;19:533-42.
18. Bsibsi M, Peferoen LA, Holtman IR, Nacken PJ, Gerritsen WH, et al. Demyelination during multiple sclerosis is associated with combined activation of microglia/macrophages by IFN- γ and alpha B-crystallin. *Acta Neuropathol* 2014;128:215-29.
19. Fischer R, Maier O. Interrelation of oxidative stress and inflammation in neurodegenerative disease: role of TNF. *Oxid Med Cell Longev* 2015;2015:610813.
20. Brown R, Benveniste H, Black SE, Charpak S, Dichgans M, et al. Understanding the role of the perivascular space in cerebral small vessel disease. *Cardiovasc Res* 2018;114:1462-73.
21. Choi YK, Kim KW. Blood-neural barrier: its diversity and coordinated cell-to-cell communication. *BMB Rep* 2008;41:345-52.
22. Lee H, Choi YK. Regenerative effects of heme oxygenase metabolites on neuroinflammatory diseases. *Int J Mol Sci* 2018;20:78.
23. Dong J, Jimi E, Zeiss C, Hayden MS, Ghosh S. Constitutively active NF-kappaB triggers systemic TNFalpha-dependent inflammation and localized TNFalpha-independent inflammatory disease. *Genes Dev* 2010;24:1709-17.
24. Miyaoka T, Seno H, Itoga M, Iijima M, Inagaki T, et al. Schizophrenia-associated idiopathic unconjugated hyperbilirubinemia (Gilbert's syndrome). *J Clin Psychiatry* 2000;61:868-71.
25. Gama Marques J, Pedro I, Ouakinin S. Unconjugated bilirubin and acute psychosis: a five years retrospective observational and controlled study in patients with schizophrenia, schizoaffective and bipolar disorders. *Int J Psychiatry Clin Pract* 2019;23:281-5.
26. Radhakrishnan R, Kanigere M, Menon J, Calvin S, Janish A, et al. Association between unconjugated bilirubin and schizophrenia. *Psychiatry Res* 2011;189:480-2.
27. Pradeep JR, Acharya MS, Radhakrishnan R, Srinivasan K. Elevated unconjugated bilirubin in schizophrenia compared to bipolar affective disorder. *Prim Care Companion CNS Disord* 2019;21:19m02448.
28. Pommerening Dornelles E, Gama Marques J, Ouakinin S. Unconjugated bilirubin and schizophrenia: a systematic review. *CNS Spectr* 2019;24:577-88.
29. Gama Marques J, Ouakinin S. Clinical profile in schizophrenia and schizoaffective spectrum: relation with unconjugated bilirubin in a prospective and controlled study with psychopathological and psychosocial variables. *CNS Spectr* 2019:1-8.
30. Miyaoka T, Seno H, Itoga M, Inagaki T, Horiguchi J. Structural brain changes in schizophrenia associated with idiopathic unconjugated hyperbilirubinemia (Gilbert's syndrome): a planimetric CT study. *Schizophr Res* 2001;52:291-3.
31. Miyaoka T, Yasukawa R, Mizuno S, Sukegawa T, Inagaki T, et al. Proton magnetic resonance spectroscopy (1H-MRS) of hippocampus, basal ganglia, and vermis of cerebellum in schizophrenia associated with idiopathic unconjugated hyperbilirubinemia (Gilbert's syndrome). *J Psychiatr Res* 2005;39:29-34.
32. Miller BJ, Buckley P, Seabolt W, Mellor A, Kirkpatrick B. Meta-analysis of cytokine alterations in schizophrenia: clinical status and antipsychotic effects. *Biol Psychiatry* 2011;70:663-71.
33. Steiner J, Mawrin C, Ziegeler A, Bielau H, Ullrich O, et al. Distribution of HLA-DR-positive microglia in schizophrenia reflects impaired cerebral lateralization. *Acta Neuropathol* 2006;112:305-16.
34. Le Pichon JB, Riordan SM, Watchko J, Shapiro SM. The neurological sequelae of neonatal hyperbilirubinemia: definitions, diagnosis and treatment of the kernicterus spectrum disorders (KSDs). *Curr Pediatr Rev* 2017;13:199-209.
35. Dalman C, Cullberg J. Neonatal hyperbilirubinaemia--a vulnerability factor for mental disorder? *Acta Psychiatr Scand* 1999;100:469-71.
36. Chowdhury JR, Kondapalli R, Chowdhury NR. Gunn rat: a model for inherited deficiency of bilirubin glucuronidation. *Adv Vet Sci Comp Med* 1993;37:149-73.
37. Gazzin S, Zelenka J, Zdrhalova L, Konickova R, Zabetta CC, et al. Bilirubin accumulation and Cyp mRNA expression in selected brain regions of jaundiced Gunn rat pups. *Pediatr Res* 2012;71:653-60.
38. Liaury K, Miyaoka T, Tsumori T, Furuya M, Wake R, et al. Morphological features of microglial cells in the hippocampal dentate gyrus of Gunn rat: a possible schizophrenia animal model. *J Neuroinflammation* 2012;9:56.
39. Hayashida M, Miyaoka T, Tsuchie K, Yasuda H, Wake R, et al. Hyperbilirubinemia-related behavioral and neuropathological changes in rats: a possible schizophrenia animal model. *Prog Neuropsychopharmacol Biol Psychiatry* 2009;33:581-8.
40. Pae CU, Paik IH, Lee C, Lee SJ, Kim JJ, et al. Decreased plasma antioxidants in schizophrenia. *Neuropsychobiology* 2004;50:54-6.
41. Yin XL, Jia QF, Zhang GY, Zhang JP, Shirao T, et al. Association between decreased serum TBIL concentration and immediate memory impairment in schizophrenia patients. *Sci Rep* 2019;9:1622.
42. Vitek L, Novotná M, Leníček M, Novotný L, Eberová J, et al. Serum bilirubin levels and UGT1A1 promoter variations in patients with schizophrenia. *Psychiatry Res* 2010;178:449-50.
43. Duan J, Göring HHH, Sanders AR, Moy W, Freda J, et al.; MGS. Transcriptomic signatures of schizophrenia revealed by dopamine perturbation in an ex vivo model. *Transl Psychiatry* 2018;8:158.

44. Song W, Zukor H, Lin SH, Hascaloivici J, Liberman A, et al. Schizophrenia-like features in transgenic mice overexpressing human HO-1 in the astrocytic compartment. *J Neurosci* 2012;32:10841-53.
45. Macías-García D, Méndez-Del Barrio C, Jesús S, Labrador MA, Adames-Gómez A, et al. Increased bilirubin levels in Parkinson's disease. *Parkinsonism Relat Disord* 2019;63:213-6.
46. Moccia M, Picillo M, Erro R, Longo K, Amboni M, et al. Increased bilirubin levels in de novo Parkinson's disease. *Eur J Neurol* 2015;22:954-9.
47. Qin XL, Zhang QS, Sun L, Hao MW, Hu ZT. Lower serum bilirubin and uric acid concentrations in patients with Parkinson's Disease in China. *Cell Biochem Biophys* 2015;72:49-56.
48. Scigliano G, Girotti F, Soliveri P, Musicco M, Radice D, et al. Increased plasma bilirubin in Parkinson patients on L-dopa: evidence against the free radical hypothesis? *Ital J Neurol Sci* 1997;18:69-72.
49. Schipper HM, Song W, Zukor H, Hascaloivici JR, Zeligman D. Heme oxygenase-1 and neurodegeneration: expanding frontiers of engagement. *J Neurochem* 2009;110:469-85.
50. Cuadrado A, Rojo AI. Heme oxygenase-1 as a therapeutic target in neurodegenerative diseases and brain infections. *Curr Pharm Des* 2008;14:429-42.
51. Hung SY, Liou HC, Kang KH, Wu RM, Wen CC, et al. Overexpression of heme oxygenase-1 protects dopaminergic neurons against 1-methyl-4-phenylpyridinium-induced neurotoxicity. *Mol Pharmacol* 2008;74:1564-75.
52. Gill SS, Patel NK, Hutton GR, O'Sullivan K, McCarter R, et al. Direct brain infusion of glial cell line-derived neurotrophic factor in Parkinson disease. *Nat Med* 2003;9:589-95.
53. Love S, Plaha P, Patel NK, Hutton GR, Brooks DJ, et al. Glial cell line-derived neurotrophic factor induces neuronal sprouting in human brain. *Nat Med* 2005;11:703-4.
54. Patel NK, Bunnage M, Plaha P, Svendsen CN, Heywood P, et al. Intraputamenal infusion of glial cell line-derived neurotrophic factor in PD: a two-year outcome study. *Ann Neurol* 2005;57:298-302.
55. Dal Ben M, Bongiovanni R, Tuniz S, Fioriti E, Tiribelli C, et al. Earliest mechanisms of dopaminergic neurons sufferance in a novel slow progressing ex vivo model of parkinson disease in rat organotypic cultures of substantia nigra. *Int J Mol Sci* 2019;20:2224.
56. Song W, Kothari V, Velly AM, Cressatti M, Liberman A, et al. Evaluation of salivary heme oxygenase-1 as a potential biomarker of early Parkinson's disease. *Mov Disord* 2018;33:583-91.
57. Ortiz GG, Pacheco-Moisés FP, Bitzer-Quintero OK, Ramírez-Anguiano AC, Flores-Alvarado LJ, et al. Immunology and oxidative stress in multiple sclerosis: clinical and basic approach. *Clin Dev Immunol* 2013;2013:708659.
58. Gonsette RE. Oxidative stress and excitotoxicity: a therapeutic issue in multiple sclerosis? *Mult Scler* 2008;14:22-34.
59. Ljubisavljevic S, Stojanovic I, Vojinovic S, Milojkovic M, Dunjic O, et al. Association of serum bilirubin and uric acid levels changes during neuroinflammation in patients with initial and relapsed demyelination attacks. *Metab Brain Dis* 2013;28:629-38.
60. Halliwell B, Zhao K, Whiteman M. Nitric oxide and peroxynitrite. The ugly, the uglier and the not so good: a personal view of recent controversies. *Free Radic Res* 1999;31:651-69.
61. van Horssen J, Schreibeit G, Drexhage J, Hazes T, Dijkstra CD, et al. Severe oxidative damage in multiple sclerosis lesions coincides with enhanced antioxidant enzyme expression. *Free Radic Biol Med* 2008;45:1729-37.
62. Artemiadis AK, Anagnostouli MC. Apoptosis of oligodendrocytes and post-translational modifications of myelin basic protein in multiple sclerosis: possible role for the early stages of multiple sclerosis. *Eur Neurol* 2010;63:65-72.
63. Kostic MS, Rajkovic JS, Floranovic MSP, Dimov ID, Pavlovic DD. Multiple sclerosis and oxidative stress - a clinical perspective. *Neurochem J* 2013;7:76-86.
64. Liu Y, Zhu B, Wang X, Luo L, Li P, et al. Bilirubin as a potent antioxidant suppresses experimental autoimmune encephalomyelitis: implications for the role of oxidative stress in the development of multiple sclerosis. *J Neuroimmunol* 2003;139:27-35.
65. Goodin DS. Chapter 21 - Glucocorticoid treatment of multiple sclerosis. In: Goodin DS, editor. *Handbook of Clinical Neurology*. Elsevier; 2014. pp. 455-64.
66. Liu Y, Li P, Lu J, Xiong W, Oger J, et al. Bilirubin possesses powerful immunomodulatory activity and suppresses experimental autoimmune encephalomyelitis. *J Immunol* 2008;181:1887-97.
67. Chora AA, Fontoura P, Cunha A, Pais TF, Cardoso S, et al. Heme oxygenase-1 and carbon monoxide suppress autoimmune neuroinflammation. *J Clin Invest* 2007;117:438-47.
68. Vitek L, Bellarosa C, Tiribelli C. Induction of mild hyperbilirubinemia: hype or real therapeutic opportunity? *Clin Pharmacol Ther* 2019;106:568-75.
69. Muchova L, Wong RJ, Hsu M, Morioka I, Vitek L, et al. Statin treatment increases formation of carbon monoxide and bilirubin in mice: a novel mechanism of in vivo antioxidant protection. *Can J Physiol Pharmacol* 2007;85:800-10.
70. Nguyen NT, Hanieh H, Nakahama T, Kishimoto T. The roles of aryl hydrocarbon receptor in immune responses. *Int Immunol* 2013;25:335-43.
71. Vitek L. Bilirubin as a signaling molecule. *Med Res Rev* 2020; Epub ahead of print [PMID: 32017160 DOI: 10.1002/med.21660]
72. Nakahama T, Hanieh H, Nguyen NT, Chinen I, Ripley B, et al. Aryl hydrocarbon receptor-mediated induction of the microRNA-132/212 cluster promotes interleukin-17-producing T-helper cell differentiation. *Proc Natl Acad Sci U S A* 2013;110:11964-9.
73. Kimura A, Naka T, Nohara K, Fujii-Kuriyama Y, Kishimoto T. Aryl hydrocarbon receptor regulates Stat1 activation and participates in the development of Th17 cells. *Proc Natl Acad Sci U S A* 2008;105:9721-6.
74. Quintana FJ, Basso AS, Iglesias AH, Korn T, Farez MF, et al. Control of T(reg) and T(H)17 cell differentiation by the aryl hydrocarbon receptor. *Nature* 2008;453:65-71.

75. Sekine H, Mimura J, Oshima M, Okawa H, Kanno J, et al. Hypersensitivity of aryl hydrocarbon receptor-deficient mice to lipopolysaccharide-induced septic shock. *Mol Cell Biol* 2009;29:6391-400.
76. Kimura A, Naka T, Nakahama T, Chinen I, Masuda K, et al. Aryl hydrocarbon receptor in combination with Stat1 regulates LPS-induced inflammatory responses. *J Exp Med* 2009;206:2027-35.
77. Nguyen NT, Kimura A, Nakahama T, Chinen I, Masuda K, et al. Aryl hydrocarbon receptor negatively regulates dendritic cell immunogenicity via a kynurenine-dependent mechanism. *Proc Natl Acad Sci U S A* 2010;107:19961-6.
78. Lee SS, Gao W, Mazzola S, Thomas MN, Csizmadia E, et al. Heme oxygenase-1, carbon monoxide, and bilirubin induce tolerance in recipients toward islet allografts by modulating T regulatory cells. *FASEB J* 2007;21:3450-7.
79. Deng J, Liang XM, Zhang XL, Ling SQ, Yang TT, et al. Relationship between serum bilirubin levels and optic neuritis. *Chin Med J (Engl)* 2013;126:3307-10.
80. Peng F, Yang Y, Liu J, Jiang Y, Zhu C, et al. Low antioxidant status of serum uric acid, bilirubin and albumin in patients with neuromyelitis optica. *Eur J Neurol* 2012;19:277-83.
81. Wang X, Jiao W, Lin M, Lu C, Liu C, et al. Resolution of inflammation in neuromyelitis optica spectrum disorders. *Mult Scler Relat Disord* 2019;27:34-41.
82. Wingerchuk DM, Lennon VA, Pittock SJ, Lucchinetti CF, Weinshenker BG. Revised diagnostic criteria for neuromyelitis optica. *Neurology* 2006;66:1485-9.
83. Tsoi VL, Hill KE, Carlson NG, Warner JE, Rose JW. Immunohistochemical evidence of inducible nitric oxide synthase and nitrotyrosine in a case of clinically isolated optic neuritis. *J Neuroophthalmol* 2006;26:87-94.
84. Cho HC. The Relationship among Homocysteine, Bilirubin, and Diabetic Retinopathy. *Diabetes Metab J* 2011;35:595-601.
85. Naruse K, Nakamura J, Hamada Y, Nakayama M, Chaya S, et al. Aldose reductase inhibition prevents glucose-induced apoptosis in cultured bovine retinal microvascular pericytes. *Exp Eye Res* 2000;71:309-15.
86. Beltramo E, Porta M. Pericyte loss in diabetic retinopathy: mechanisms and consequences. *Curr Med Chem* 2013;20:3218-25.
87. Peng F, Deng X, Yu Y, Chen X, Shen L, et al. Serum bilirubin concentrations and multiple sclerosis. *J Clin Neurosci* 2011;18:1355-9.
88. Li RY, Cao ZG, Zhang JR, Li Y, Wang RT. Decreased serum bilirubin is associated with silent cerebral infarction. *Arterioscler Thromb Vasc Biol* 2014;34:946-51.
89. Higuchi S, Kabeya Y, Uchida J, Kato K, Tsukada N. Low bilirubin levels indicate a high risk of cerebral deep white matter lesions in apparently healthy subjects. *Sci Rep* 2018;8:6473.
90. Garde E, Mortensen EL, Krabbe K, Rostrup E, Larsson HB. Relation between age-related decline in intelligence and cerebral white-matter hyperintensities in healthy octogenarians: a longitudinal study. *Lancet* 2000;356:628-34.
91. Dettke S, Markus HS. The clinical importance of white matter hyperintensities on brain magnetic resonance imaging: systematic review and meta-analysis. *BMJ* 2010;341:c3666.
92. Suwanwela NC, Chutinetr A. Risk factors for atherosclerosis of cervicocerebral arteries: intracranial versus extracranial. *Neuroepidemiology* 2003;22:37-40.
93. Wong KS, Li H, Chan YL, Ahuja A, Lam WW, et al. Use of transcranial Doppler ultrasound to predict outcome in patients with intracranial large-artery occlusive disease. *Stroke* 2000;31:2641-7.
94. Zhong K, Wang X, Ma X, Ji X, Sang S, et al. Association between serum bilirubin and asymptomatic intracranial atherosclerosis: results from a population-based study. *Neurol Sci* 2020; Epub ahead of print. doi: 10.1007/s10072-020-04268-x
95. Jian Y, Zhao L, Wang H, Li T, Zhang L, et al. Bilirubin: a novel predictor of hemorrhagic transformation and symptomatic intracranial hemorrhage after mechanical thrombectomy. *Neurol Sci* 2019; Epub ahead of print. doi:10.1007/s10072-019-04182-x
96. Ishikawa K, Navab M, Leitinger N, Fogelman AM, Lusis AJ. Induction of heme oxygenase-1 inhibits the monocyte transmigration induced by mildly oxidized LDL. *J Clin Invest* 1997;100:1209-16.
97. Kawamura K, Ishikawa K, Wada Y, Kimura S, Matsumoto H, et al. Bilirubin from heme oxygenase-1 attenuates vascular endothelial activation and dysfunction. *Arterioscler Thromb Vasc Biol* 2005;25:155-60.
98. Ollinger R, Bilban M, Erat A, Froio A, McDaid J, et al. Bilirubin: a natural inhibitor of vascular smooth muscle cell proliferation. *Circulation* 2005;112:1030-9.
99. Pae HO, Oh GS, Lee BS, Rim JS, Kim YM, et al. 3-Hydroxyanthranilic acid, one of L-tryptophan metabolites, inhibits monocyte chemoattractant protein-1 secretion and vascular cell adhesion molecule-1 expression via heme oxygenase-1 induction in human umbilical vein endothelial cells. *Atherosclerosis* 2006;187:274-84.
100. Basiglio CL, Arriaga SM, Pelusa HF, Almará AM, Roma MG, et al. Protective role of unconjugated bilirubin on complement-mediated hepatocytolysis. *Biochim Biophys Acta* 2007;1770:1003-10.
101. Větvicka V, Miler I, Šima P, Táborský L, Fornůšek L. The effect of bilirubin on the Fc receptor expression and phagocytic activity of mouse peritoneal macrophages. *Folia Microbiol (Praha)* 1985;30:373-80.
102. Dohi K, Satoh K, Ohtaki H, Shioda S, Miyake Y, et al. Elevated plasma levels of bilirubin in patients with neurotrauma reflect its pathophysiological role in free radical scavenging. *In Vivo* 2005;19:855-60.
103. Wang J, Doré S. Heme oxygenase-1 exacerbates early brain injury after intracerebral haemorrhage. *Brain* 2007;130:1643-52.
104. Orozco-Ibarra M, Estrada-Sánchez AM, Massieu L, Pedraza-Chaverri J. Heme oxygenase-1 induction prevents neuronal damage triggered during mitochondrial inhibition: role of CO and bilirubin. *Int J Biochem Cell Biol* 2009;41:1304-14.
105. Zhao Q, Qu R, Teng L, Yin C, Yuan Y. HO-1 protects the nerves of rats with cerebral hemorrhage by regulating the PI3K/AKT signaling pathway. *Neuropsychiatr Dis Treat* 2019;15:1459-68.
106. Feng J, Zhang P, Chen X, He G. PI3K and ERK/Nrf2 pathways are involved in oleanolic acid-induced heme oxygenase-1 expression in

- rat vascular smooth muscle cells. *J Cell Biochem* 2011;112:1524-31.
107. Li Q, Huai L, Zhang C, Wang C, Jia Y, et al. Icaritin induces AML cell apoptosis via the MAPK/ERK and PI3K/AKT signal pathways. *Int J Hematol* 2013;97:617-23.
108. Doré S, Goto S, Sampei K, Blackshaw S, Hester LD, et al. Heme oxygenase-2 acts to prevent neuronal death in brain cultures and following transient cerebral ischemia. *Neuroscience* 2000;99:587-92.
109. Wang J, Zhuang H, Doré S. Heme oxygenase 2 is neuroprotective against intracerebral hemorrhage. *Neurobiol Dis* 2006;22:473-6.
110. Chang EF, Wong RJ, Vreman HJ, Igarashi T, Galo E, et al. Heme oxygenase-2 protects against lipid peroxidation-mediated cell loss and impaired motor recovery after traumatic brain injury. *J Neurosci* 2003;23:3689-96.
111. Zou ZY, Liu J, Chang C, Li JJ, Luo J, et al. Biliverdin administration regulates the microRNA-mRNA expressional network associated with neuroprotection in cerebral ischemia reperfusion injury in rats. *Int J Mol Med* 2019;43:1356-72.
112. Wegiel B, Gallo D, Ciszmadia E, Roger T, Kaczmarek E, et al. Biliverdin inhibits Toll-like receptor-4 (TLR4) expression through nitric oxide-dependent nuclear translocation of biliverdin reductase. *Proc Natl Acad Sci U S A* 2011;108:18849-54.
113. Holst B, Raby AC, Hall JE, Labéta MO. Complement takes its Toll: an inflammatory crosstalk between Toll-like receptors and the receptors for the complement anaphylatoxin C5a. *Anaesthesia* 2012;67:60-4.
114. Pulliam JV, Xu Z, Ford GD, Liu C, Li Y, et al. Computational identification of conserved transcription factor binding sites upstream of genes induced in rat brain by transient focal ischemic stroke. *Brain Res* 2013;1495:76-85.
115. Ilzecka J, Stelmasiak Z. Serum bilirubin concentration in patients with amyotrophic lateral sclerosis. *Clin Neurol Neurosurg* 2003;105:237-40.
116. Oren DA, Desan PH, Boutros N, Anand A, Charney DS. Effects of light on low nocturnal bilirubin in winter depression: a preliminary report. *Biol Psychiatry* 2002;51:422-5.
117. Raszewski G, Gustaw K, Chwedorowicz R. Endogenous antioxidant status in dementia patients with cognitive impairment and normal cognitive function. *Clin Endocrinol* 2011;1:13-23.
118. Baierle M, Nascimento SN, Moro AM, Brucker N, Freitas F, et al. Relationship between inflammation and oxidative stress and cognitive decline in the institutionalized elderly. *Oxid Med Cell Longev* 2015;2015:804198.
119. Kim TS, Pae CU, Yoon SJ, Jang WY, Lee NJ, et al. Decreased plasma antioxidants in patients with Alzheimer's disease. *Int J Geriatr Psychiatry* 2006;21:344-8.
120. Di Domenico F, Barone E, Mancuso C, Perluigi M, Cocciolo A, et al. HO-1/BVR-a system analysis in plasma from probable Alzheimer's disease and mild cognitive impairment subjects: a potential biochemical marker for the prediction of the disease. *J Alzheimers Dis* 2012;32:277-89.
121. Ishizuka K, Kimura T, Yoshitake J, Akaike T, Shono M, et al. Possible assessment for antioxidant capacity in Alzheimer's disease by measuring lymphocyte heme oxygenase-1 expression with real-time RT-PCR. *Ann N Y Acad Sci* 2002;977:173-8.
122. Barone E, Di Domenico F, Sultana R, Coccia R, Mancuso C, et al. Heme oxygenase-1 posttranslational modifications in the brain of subjects with Alzheimer disease and mild cognitive impairment. *Free Radic Biol Med* 2012;52:2292-301.
123. Barone E, Di Domenico F, Mancuso C, Butterfield DA. The Janus face of the heme oxygenase/biliverdin reductase system in Alzheimer disease: it's time for reconciliation. *Neurobiol Dis* 2014;62:144-59.
124. Liu B, Moloney A, Meehan S, Morris K, Thomas SE, et al. Iron promotes the toxicity of amyloid beta peptide by impeding its ordered aggregation. *J Biol Chem* 2011;286:4248-56.
125. Schipper HM. Heme oxygenase expression in human central nervous system disorders. *Free Radic Biol Med* 2004;37:1995-2011.
126. Schipper HM. Heme oxygenase-1: transducer of pathological brain iron sequestration under oxidative stress. *Ann N Y Acad Sci* 2004;1012:84-93.
127. Buonocore G, Perrone S, Bracci R. Free radicals and brain damage in the newborn. *Biol Neonate* 2001;79:180-6.
128. Matyas M, Zaharie G. Antioxidants at newborns. *Antioxidants* 2019. Available from: <https://doi.org/10.5772/intechopen.85175> [Last accessed on 27 Mar 2020]
129. Finkel T, Holbrook NJ. Oxidants, oxidative stress and the biology of ageing. *Nature* 2000;408:239-47.
130. Gemma C, Vila J, Bachstetter A, Bickford PC. Oxidative stress and the aging brain: from theory to prevention. In: Riddle DR, editor. *Brain aging: models, methods, and mechanisms*. FL: CRC Press/Taylor & Francis; 2007.
131. Nitti M, Piras S, Marinari UM, Moretta L, Pronzato MA, et al. HO-1 induction in cancer progression: a matter of cell adaptation. *Antioxidants (Basel)* 2017;6:29.
132. Attucks OC, Jasmer KJ, Hannink M, Kassiss J, Zhong Z, et al. Induction of heme oxygenase I (HMOX1) by HPP-4382: a novel modulator of Bach1 activity. *PLoS One* 2014;9:e101044.
133. Campbell NK, Fitzgerald HK, Malara A, Hambly R, Sweeney CM, et al. Naturally derived Heme-Oxygenase 1 inducers attenuate inflammatory responses in human dendritic cells and T cells: relevance for psoriasis treatment. *Sci Rep* 2018;8:10287.
134. Croft KD, Zhang D, Jiang R, Ayer A, Shengule S, et al. Structural requirements of flavonoids to induce heme oxygenase-1 expression. *Free Radic Biol Med* 2017;113:165-75.
135. George EM, Stout JM, Stec DE, Granger JP. Heme oxygenase induction attenuates TNF- α -induced hypertension in pregnant rodents. *Front Pharmacol* 2015;6:165.
136. Gerbitz A, Ewing P, Wilke A, Schubert T, Eissner G, et al. Induction of heme oxygenase-1 before conditioning results in improved survival and reduced graft-versus-host disease after experimental allogeneic bone marrow transplantation. *Biol Blood Marrow Transplant* 2004;10:461-72.
137. Hamamura RS, Ohyashiki JH, Kurashina R, Kobayashi C, Zhang Y, et al. Induction of heme oxygenase-1 by cobalt protoporphyrin

- enhances the antitumour effect of bortezomib in adult T-cell leukaemia cells. *Br J Cancer* 2007;97:1099-105.
138. Kirby RJ, Divlianska DB, Whig K, Bryan N, Morfa CJ, et al. Discovery of novel small-molecule inducers of heme oxygenase-1 that protect human iPSC-derived cardiomyocytes from oxidative stress. *J Pharmacol Exp Ther* 2018;364:87-96.
139. Maamoun H, Zachariah M, McVey JH, Green FR, Agouni A. Heme oxygenase (HO)-1 induction prevents Endoplasmic Reticulum stress-mediated endothelial cell death and impaired angiogenic capacity. *Biochem Pharmacol* 2017;127:46-59.
140. Mikkelsen RB, Rabender CS, Graves P, Anscher MS. Induction of HO-1 and CO synthesis radiosensitizes tumor epithelial cells with CO acting as a NO mimetic. *Int J Radiat Oncol Biol Phys* 2014;90:S105.
141. Muchova L, Vanova K, Suk J, Micuda S, Dolezelova E, et al. Protective effect of heme oxygenase induction in ethinylestradiol-induced cholestasis. *J Cell Mol Med* 2015;19:924-33.
142. Pittala V, Vanella L, Salerno L, Di Giacomo C, Acquaviva R, et al. Novel caffeic acid phenethyl ester (Cape) analogues as inducers of heme oxygenase-1. *Curr Pharm Des* 2017;23:2657-64.
143. Ramma W, Ahmed A. Therapeutic potential of statins and the induction of heme oxygenase-1 in preeclampsia. *J Reprod Immunol* 2014;101-2:153-60.
144. Schipper HM. Is glial heme oxygenase-1 suppression in neurodegenerative disorders permissive for neural repair? *Neural Regen Res* 2015;10:208-10.
145. Simon T, Pogu J, Rémy S, Brau F, Pogu S, et al. Inhibition of effector antigen-specific T cells by intradermal administration of heme oxygenase-1 inducers. *J Autoimmun* 2017;81:44-55.
146. Skrzypek K, Tertilt M, Golda S, Ciesla M, Weglarczyk K, et al. Interplay between heme oxygenase-1 and miR-378 affects non-small cell lung carcinoma growth, vascularization, and metastasis. *Antioxid Redox Signal* 2013;19:644-60.
147. Strasky Z, Zemankova L, Nemeckova I, Rathouska J, Wong RJ, et al. *Spirulina platensis* and phycocyanobilin activate atheroprotective heme oxygenase-1: a possible implication for atherogenesis. *Food Funct* 2013;4:1586-94.
148. Won AN, Kim SA, Ahn JY, Han JH, Kim CH, et al. HO-1 Induction by *Selaginella tamariscina* Extract Inhibits Inflammatory Response in Lipopolysaccharide-Stimulated RAW 264.7 Macrophages. *Evid Based Complement Alternat Med* 2018;2018:7816923.

Review

Open Access



Cerebrospinal fluid biomarkers in idiopathic normal pressure hydrocephalus

Xin-Jie Zhang[#], Jian Guo[#], Jun Yang[#]

Department of Neurosurgery, Xiang'an Hospital of Xiamen University, School of Medicine, Xiamen University, Xiamen 361102, Fujian, China.

[#]Authors contributed equally.

Correspondence to: Dr. Jun Yang, Department of Neurosurgery, Xiang'an Hospital of Xiamen University, School of Medicine, Xiamen University, Xiamen 361102, Fujian, China. E-mail: junyangyang@hotmail.com

How to cite this article: Zhang XJ, Guo J, Yang J. Cerebrospinal fluid biomarkers in idiopathic normal pressure hydrocephalus. *Neuroimmunol Neuroinflammation* 2020;7:109-19. <http://dx.doi.org/10.20517/2347-8659.2019.018>

Received: 28 Nov 2019 **First Decision:** 1 Feb 2020 **Revised:** 15 Feb 2020 **Accepted:** 16 Mar 2020 **Available online:** 11 May 2020

Science Editor: George P. Paraskevas **Copy Editor:** Jing-Wen Zhang **Production Editor:** Jing Yu

Abstract

Idiopathic normal pressure hydrocephalus (iNPH) is characterized by abnormal cerebrospinal fluid (CSF) flow and consequent cerebral ventricular enlargement due to imbalance of CSF production and absorption. The typical triad symptoms, namely cognitive decline, gait disturbance, and urinary incontinence, are thought to be caused by disruption of CSF circulation. However, some patients may still experience symptomatic progression after functional shunting, suggesting that iNPH is far more complicated than a simple disorder of CSF circulation. Moreover, the diagnostic workup of iNPH can be challenging due to symptomatic and neuroimaging overlaps with other neurological disorders, such as Alzheimer's disease. Furthermore, accumulating studies indicate that the pathogenesis of iNPH might relate to multiple mechanisms, including abnormalities of brain development, brain extracellular matrix, synaptic function, blood flow, and cerebral metabolism. Therefore, iNPH is not an isolated entity in occurrence and development. Nevertheless, different pathogeneses may result in protein content changes in CSF, and the biomarkers in CSF may reflect the possible mechanisms involving the etiology of iNPH and are potentially useful in assisting the diagnosis and treatment selection. In this review, we summarize the main findings of CSF biomarkers and aim to outline a possible synthetic profile in assisting iNPH diagnosis and therapeutic options.

Keywords: Idiopathic normal pressure hydrocephalus, cerebrospinal fluid, biomarkers, neurodegenerative diseases



© The Author(s) 2020. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, sharing, adaptation, distribution and reproduction in any medium or format, for any purpose, even commercially, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.



INTRODUCTION

Idiopathic normal pressure hydrocephalus (iNPH) is one of the disabling neurological disorders whose potential treatability is significantly impacted by the timeliness of unequivocal diagnosis. iNPH is characterized by ventriculomegaly that is caused by an imbalance between cerebrospinal fluid (CSF) production and absorption. The characteristic triad symptoms of dementia, gait disturbance, and urinary incontinence are thought to be caused by a disruption of CSF dynamics^[1,2]. Therefore, the triad symptoms of iNPH could be surgically treatable with a diversion of CSF into peritoneal cavity or heart^[3]. However, the diagnostic workup of iNPH can be a challenge due to neuroimaging and symptomatic overlaps with other neurological disorders, such as Alzheimer's disease (AD) and subcortical ischemic vascular disease, especially at early stage. Moreover, although the cognitive decline could be proceeded by these diseases, they are not equally responsive to the treatment of CSF shunting. Therefore, further effort to improve the diagnosis of iNPH would benefit the current imaging and symptomatic diagnostic criteria. Increasing studies indicate that the pathogenesis of iNPH involves multiple mechanisms, including abnormalities of brain development, brain extracellular matrix, synaptic function, blood flow, and cerebral metabolism, which could result in protein content changes in CSF. On the other hand, impaired CSF absorption could lead to a pathological flow of CSF into the periventricular tissues to initiate a cascade of pathological processes such as edema and consequent neuronal degenerative changes^[4]. Therefore, measurements of different biomarkers in CSF may reflect the underlying neuropathological changes of the brain and could play an important role in revealing the possible etiological mechanisms. Furthermore, its detection may facilitate the timeliness and accuracy of iNPH diagnosis, and thus becomes potentially useful for therapeutic selection and treatment response monitoring. In addition, the biomarkers could help to differentiate iNPH from other neurological disorders, which might mimic iNPH symptomatology but show unsatisfactory outcomes after shunting^[5,6]. Despite a growing interest, the CSF biomarker profile in iNPH has not yet been identified definitively. In this review, we summarize the main findings of CSF biomarkers regarding iNPH and outline a rough CSF profile in order to assist iNPH diagnosis and provide adequate treatment. It is notable that, due to the etiological complexity of iNPH, most biomarkers might lack specificity for iNPH diagnosis and are possibly coincidental, confounding with other overlapping neurological diseases. In addition, in comparison with a cortical brain biopsy or neuropsychological testing, biomarkers may also have limitations in distinguishing iNPH from comorbid iNPH plus AD^[7], as well as in predicting clinical cognitive outcome post shunting^[6,8,9]. However, a combination of more than one biomarker may enhance the predictive value and provide more viable and accurate solutions. Ideally, the dynamic changes of biomarker measured before and after surgical diversion of CSF would supply useful clinical information for the diagnosis and assistance in monitoring disease progression. The biomarkers could be categorized as AD discrimination, neurodegeneration and demyelination, neuroinflammation, neuropeptides and cerebral metabolites, and as biomarkers in response to cerebral and vascular insulting, among others^[1,2,10,11].

BIOMARKERS FOR AD DISCRIMINATION

Dementia in iNPH is potentially reversible if adequately treated. However, it often resembles the clinical appearance of patients with AD, such as memory decline, as well as attention and executive impairment^[12]. Urinary incontinence and gait disturbance may also occur in both diseases due to disturbed subcortical network caused by vascular pathology. Moreover, ventricular enlargement may have been observed in AD patients as a result of cerebral atrophy rather than CSF circulation impairment^[13]. Furthermore, the pathological examination of cortical brain biopsies performed during placement of CSF shunts revealed AD neurodegenerative changes in 24% of iNPH patients, suggesting a high comorbidity of both diseases. Thus, cortical brain biopsy may provide a valuable predictive way for outcome evaluation^[6,8]. However, cortical brain biopsy is not always available or appropriate in some cases. Moreover, both iNPH and AD diseases may manifest sleep disturbances, which correlate with dysfunction of the glia-lymphatic (glymphatic)

system, consequently building-up of brain metabolic wasters, favoring dementia development^[14,15]. Therefore, it is always a challenge to discriminate iNPH and AD diseases in clinical practice.

The glymphatic system facilitates cerebral metabolite and brain fluid clearance during sleep via glia-supported perivascular channels. This system facilitates efflux of cerebrospinal and interstitial fluid via the perivascular spaces to the meningeal and cervical lymphatic vessels, assisting the draining/clearing of metabolic wastes from the central nervous. The glymphatic flux is proposed to be driven by cardiac-induced arterial pulsation^[16], and may be possibly manipulated through change of intracranial pressure pulsatility with our cardiac-gated device^[17]. Most interestingly, the action of glymphatic flux is predominant during sleep^[18], and up to 90% of iNPH patients are associated with obstructive sleep apnea, a common sleep disorder^[19]. Blockage of the airway in obstructive sleep apnea causes increased awakenings and decreased quality of sleep, resulting in glymphatic dysfunction and increased cerebral A β aggregation^[20]. Patients with obstructive sleep apnea encounter reduced oxygen intake due to intermittent airway obstruction. Excessive breathing against a closed airway induces negative intrathoracic pressure, sufficient to cause atrial distortion and reduced venous return to the heart^[19] and ultimately affect arterial pulsation, resulting in dysfunction of glymphatic flux.

Many studies have shown impaired glymphatic function in both iNPH and AD. Furthermore, iNPH and AD patients share multiple clinical and pathologic features such as A β deposition, cerebrovascular inflammation, impaired localization of perivascular astrocyte aquaporin-4 (AQP4), and sleep disturbances^[15]. Therefore, it is a diagnostic challenge in daily practice for iNPH and AD. Although many biomarkers have been investigated for their discrimination, amyloid- β 42 (A β_{42}), total-tau (t-tau), and phosphorylated tau (p-tau) are the most robust candidate markers to discriminate iNPH from AD patients^[1,2]. A β_{42} is lower in both iNPH and AD patients compared with healthy control, and A β_{42} does not separate iNPH and AD. Tau protein is a microtubule-associated protein and is a marker for neuronal degeneration^[21]. The levels of t-tau and p-tau are higher in AD patients compared with iNPH patients and controls, whereas the levels of t-tau and p-tau are within normal range in iNPH patients. The combination of these biomarkers, i.e., the reduced A β_{42} with concomitant normal or reduced t-tau and p-tau levels in iNPH coupled with reduced A β_{42} with concomitant increased both t-tau and p-tau levels in AD, may significantly improve the accuracy of differential diagnosis between AD and iNPH patients^[22]. The mechanism of lower A β_{42} level in iNPH patients is unknown. However, the reduced production of A β_{42} due to a decline in brain metabolism in the periventricular zone in iNPH patients^[23,24] and interstitial A β deposition due to impaired glymphatic function may be possible reasons^[15]. Meanwhile, the low concentrations of CSF t-tau and p-tau do not support the major cortical degenerative process in iNPH^[24,25], whereas, in AD patients, the core pathological changes are the accumulation of abnormally folded beta-amyloid and tau proteins in the plaques and neuronal tangles^[26], and the progressive deposition of amyloid plaques lowers A β_{42} level. Moreover, concurrent axonal degenerations and neurofibrillary tangle formation further increase t-tau and p-tau CSF levels in AD patients^[27]. The representative information and main biomarkers for assisting differential diagnosis of iNPH and AD are summarized in the attached Table 1.

NEURODEGENERATION AND DEMYELINATION

The disturbance of CSF circulation could lead to a potentially hostile milieu for cerebral structures, especially periventricular areas and subcortical structures, and could result in vascular lesions, destruction of periventricular white matter, and subsequent neurodegeneration and demyelination^[28-30]. Such pathological changes could be estimated with the examination of CSF contents, such as neurofilament light chains (NFL), myelin basic protein (MBP), and leucine-rich- α 2-glycoprotein (LRG)^[31-33]. NFL is a cytoskeletal element in nerve axons and dendrites, and therefore could be considered as a biomarker for axonal damage in patients with iNPH^[31,34]. Although some studies did not find difference of CSF NFL

Table 1. The representative information and biomarkers in iNPH and AD

	iNPH	AD	References	Comments
Etiology	Multiple	Multiple	[1-4]	
Dementia	10%	60%-70%	[12,15]	
Ventriculomegaly	↑↑	↑	[6,13]	18-42% of iNPH also had AD brain biopsy findings
OSA	65%-90%	44%	[15,19]	OSA: obstructive sleep apnea
GFD	Yes	Yes	[14-15,18]	GFD: Glymphatic flux dysfunction
WMLs	↑	↑	[64-66]	WMLs: cerebral white matter lesions
PWMD	↑	↑	[28-30]	PWMD: periventricular white matter damage
FSO	↑↑	↑	[1-3]	FSO: favorable surgical outcome
*Aβ42	↓	↓	[1-2,22-24]	Amyloid-beta-42. No difference vs. AD, ↓ vs. control
*t-tau	↓/-	↑	[1-2,21-24,27]	Total tau. ↓ vs. AD, no difference vs. control
*p-tau	↓/-	↑	[1-2,22,27]	Phosphorylated tau. ↓ vs. AD, no difference vs. control
NFL	↑	N/A	[12,31-34]	Neurofilament light chains. Correlated with PWMD and FSO
MBP	↑	↑	[31-33,39-41]	Myelin basic protein. Correlated with PWMD and FSO
LRG	↑	↑	[31-33,43]	Leucine-rich-α2-glycoprotein
TNF-α	↑	N/A	[45-46]	Tumor-necrosis factor α. Correlated with FSO
TGF-β1	↑	N/A	[47-49]	Transforming growth factor β1
IL-1β	↑	↑	[44,50-52]	Pro-inflammatory cytokines, interleukin-1β
IL-6	↑	↑	[50-52]	Pro-inflammatory cytokines, interleukin-6
IL-10	↑	↑	[50-52]	Anti-inflammatory cytokine, interleukin-10
TFPI-2	↑	↑	[50-52]	Tissue factor pathway inhibitor 2
YKL-40	↑	↑	[50-53]	Chitinase-3-like protein-1
MCP-1	↑	↑	[50-52]	Monocyte chemoattractant protein-1
SOM	↑/↓	N/A	[10,54-55]	Somatostatin
VIP	↓	N/A	[10,54-55,57]	Vasoactive intestinal peptide
NPY	↓	N/A	[10,54-55]	Neuropeptide Y
DSIP	↓	N/A	[10,54-55]	Delta-sleep inducing peptide
NGF	↑↑	N/A	[69,70]	Nerve growth factor
VEGF	↑	N/A	[59,71-73]	Vascular endothelial growth factor. Correlated with FSO
GFAP	↑	N/A	[34,76]	Glial fibrillary acidic protein
PGDS	↓	-	[77]	Prostaglandin D synthase

*Strengths; other weaknesses. ↑: increased; ↓: decreased; -: normal; N/A: not available; iNPH: Idiopathic normal pressure hydrocephalus; AD: Alzheimer's disease

levels between iNPH and AD patients^[11,32], as well as controls^[35], other studies demonstrated increased CSF NFL levels, and the increase paralleled the degeneration of large myelinated axons in iNPH^[31,36]. In addition, some studies observed that the ventricular NFL level directly correlated with altered signals in periventricular white matter in brain MRI^[37]. Moreover, one study demonstrated that high preoperative NFL level was associated with favorable surgical outcomes, and suggested that NFL could possibly be used as an indicator for neurodegeneration and a marker of ongoing axonal damage^[38].

Demyelination of the periventricular white matter could occur in hydrocephalus due to the result of mechanical stretching. MBP is an oligodendroglial structural protein of myelin and sulfatide is a glycosphingolipid component of myelin, and they are essential for the maintenance of central nervous system myelin and axon structure^[32,39]. Both MBP and sulfatide are well known indicators for ongoing demyelination and therefore are attractive markers for the pathological process^[40]. However, the CSF levels of MBP are higher in many different neurologic disorders, including iNPH and cerebrovascular diseases, leading to lack of specification for iNPH diagnosis^[32,36], whereas it is demonstrated that changes of MBP levels are correlated with periventricular white matter damage^[41]. When comparing the levels of MBP pre- and post-shunting, the results showed that the levels of MBP decreased post-shunting, suggesting that MBP could be used for evaluation of brain damage and shunting effect^[42].

LRG is an astrocytic protein and could be induced by inflammation. The LRG level in CSF increases with age in iNPH and other dementia diseases. It was speculated that the accumulation of LRG in the brains is

one of the causes of neurodegeneration, therefore its level in CSF could be an anticipated marker for early diagnosis of iNPH and other dementia diseases^[33,43].

Taking together, all these markers allow tracking the integrity of periventricular and subcortical structures. Although they are not disease specific, their changes in CSF directly reflect cerebral damage, and they may be useful indicators in comparative analyses between iNPH and other neurodegenerative diseases.

NEUROINFLAMMATION

Cytokines mediate inflammatory response and often correlate with neurodegeneration in neurological diseases. The profile of CSF cytokines provides access to explore the pathogenic mechanisms of different neurological diseases and therapeutic approaches^[44]. Abundant CSF cytokines have been investigated in iNPH patients, but a more definite profile still needs to be clarified^[32,36].

Tumor-necrosis factor (TNF- α) is a cytokine of inflammatory mediator and its level in CSF is significantly high in iNPH patients^[45,46]. Most interestingly, the CSF level of TNF- α returned to the control level in the patients with shunt improvement. Because of its short half-life, the increased CSF TNF- α may be caused by increased production rather than the accumulation due to CSF stagnation, which suggests that TNF- α in CSF might be used as a candidate marker for the evaluation of demyelination and disease progression in iNPH patients. More studies are needed for validation.

Transforming growth factor β 1 (TGF- β 1), one of the three cytokines in the TGF family, plays a role in cell differentiation and tissue modification during brain development. It could be released from microglia and astrocytes in response to cerebral insult to initiate neuroinflammation and neurodegeneration through the induction of fibrosis, vascular hypertrophy, accumulation of extracellular matrix components, and neuronal apoptosis^[47-49]. TGF- β level was found to be higher in iNPH patients than controls, and was considered to be a reliable index of cerebral damage in iNPH^[49].

Other increased inflammatory biomarkers measured in iNPH patients include IL-1 β and IL-6 (pro-inflammatory cytokines), IL-10 (anti-inflammatory cytokine), tissue factor pathway inhibitor 2 (TFPI-2), chitinase-3-like protein-1 (YKL-40), and monocyte chemoattractant protein (MCP-1)^[50-52]. However, as similar changes are also observed in AD and Parkinson's disease, these changes only reflect an underlying neuroinflammatory processes of pro-inflammatory reaction (IL-1 β and IL-6) and compensatory anti-inflammatory reaction (IL-10), rather than disease-specific indicators^[44,51,52]. TFPI-2 is involved in inflammatory process by recruiting astrocytes and microglia to the injury site^[50]. YKL-40 is then released from astrocyte and/or microglia in response to neuroinflammation. The increased CSF YKL-40 levels seem to be correlated with cognitive decline and therefore to predict progression of dementia^[53]. However, more studies are deserved on the clinical use of this novel promising neuroinflammation biomarker^[35,48].

NEUROPEPTIDES AND CEREBRAL METABOLITES

Neuropeptides, including somatostatin, vasoactive intestinal peptide, neuropeptide Y, and delta-sleep inducing peptide, have been evaluated by various groups^[10,54,55]. Decreased CSF somatostatin levels suggest damage to the hypothalamus and the cortical neurons that normally have high concentrations of somatostatin^[54]. Higher level of somatostatin correlates with better visual memory and mental condition in iNPH patients, proposing that somatostatin may have a modulatory role in cognition^[10]. Vasoactive intestinal peptide is a potent vasodilator and therefore may play a role in chronic ischemia, and the CSF level is usually higher in iNPH patients with cerebrovascular disease^[55-57]. Delta-sleep inducing peptide is a nine-amino acid peptide with a role in sleep-wakefulness regulation. iNPH patients with lower delta-sleep inducing peptide level show worse psychomotor performance^[56]. Several studies also reported reduced levels of neuropeptide Y in iNPH patients^[54-56].

Cerebral metabolism changes may occur in iNPH patients. iNPH patients were also reported to have altered levels of lactate, an end product of anaerobic glycolysis underlying a presence of chronic ischemia^[58,59]. Free-radical peroxidation could result in cellular dysfunction and may therefore be implicated in the pathogenesis of iNPH and dementia. A study showed that the levels of free-radical peroxidation products significantly increased in iNPH patients^[60]. The authors implied that peroxidation of cytoplasmic membranes might be involved in the development of cognitive dysfunction in iNPH.

BLOOD-BRAIN BARRIER CHANGE AND BIOMARKERS RESPONDING TO CEREBRAL AND VASCULAR INSULTING IN INPH

Blood-brain barrier is a physically powerful gateway that strictly monitors and controls the interchange of substances between central nervous system and blood flow^[61]. Its function is strictly dependent on the integrity of microvascular endothelium and thus affected by many pathophysiological risk factors, including vascular/hemodynamic changes, inflammation, *etc.*, and in turn affects the homeostasis of central nervous system^[62]. The “CSF/blood albumin ratio” represents a reliable index of blood-brain barrier function. Blood-brain barrier impairment was reported in different neurodegenerative diseases, including AD and cerebral vascular disease^[63]. Nowadays, it has been scarcely evaluated in iNPH patients, but available reports indicate a substantial preservation of the blood-brain barrier^[22,36].

Vascular risk factor may be a component of subcortical neuropathology in the development of iNPH^[2]. As key components, cerebral white matter lesions and hypertension were reported to be related to the pathophysiology of iNPH^[64-66]. White matter lesions, involved in different cognitive processes and/or clinical outcomes, are associated with small vessel disease and white matter ischemia. The association between iNPH and white matter lesions indicates the involvement of microvascular disturbances in the white matter and in the pathological processes of iNPH. In addition, hypertension increases the risk of iNPH through the mechanisms of involved small vessel diseases, including hypertension induced endothelial damage and resultant extravasation of blood products into white matter, impaired blood flow with reduced metabolism, and direct mechanical effect on ventricular size^[66,67]. Therefore, identifications of vascular related risk factors may improve diagnostic accuracy and address the underlying pathology regarding the development of iNPH, and ultimately provide suitable intervention for iNPH management. Overall, the dynamic and morphological alterations in subcortical structure of iNPH brain could be resulted from white matter lesions, hypertension related vascular lesions, destruction of periventricular white matter axons and gliosis, and impaired CSF circulation^[28]. Such pathological alterations could affect CSF protein contents and biomarkers in CSF could mirror the underlying pathologic alterations. As markers of subcortical damage, at least three proteins have been measured in iNPH patients, including NFL, LRG, and MBP^[1]. The functions and clinical application of these proteins are discussed above. In summary, NFL is a cytoskeletal protein for maintenance of axonal architecture and is considered as a marker for neuronal morphological integrity^[31]. Although it has also been assessed as a biomarker for inflammatory and neurodegenerative diseases, it has been observed that ventricular NFL levels in iNPH patients directly correlate with more extensive altered signals in periventricular white matter in brain MRI^[1]. LRG is an astrocytic protein and is increased in CSF of iNPH patients, suggesting a potential biomarker for iNPH, but it also changes with aging and non-specific inflammation^[68]. MBP is an oligodendroglial structural protein of myelin. Its CSF levels are increased in iNPH patients and other cerebrovascular and neurodegenerative diseases, indicating the damage of periventricular white matter^[1,32].

In addition, nerve growth factor (NGF) play an important role in neuro-regeneration in response to brain injury and age-related atrophy. NGF is scarcely detectable in innervated tissues, but denervation of cerebral tissue could lead to the production of NGF and become measurable in the target tissues^[69]. The CSF level of NGF was found to be significantly higher in hydrocephalus patients compared with the controls^[70], which

suggests the possibility that the increased NGF levels could represent an increased cerebral regeneration after shunting.

Vascular endothelial growth factor plays roles in many cerebral physiological and pathological modifications, and its level in CSF is respondent to ischemic condition involved in different neurological disorders^[59,71-73]. Our group demonstrated that the CSF levels of vascular endothelial growth factor in iNPH patients have circadian variations and exercise induced increasing^[74]. The higher concentration of vascular endothelial growth factor level in CSF is associated with less response to shunting and worse clinical outcome, suggesting a possible concurrent ischemic or vascular injury in iNPH patients^[73,75].

Glial fibrillary acidic protein is a marker for gliosis^[34,76]. In iNPH patients, the CSF level of glial fibrillary acidic protein was increased when compared with controls, and correlated with disease progression^[38]. The increased glial fibrillary acidic protein level in CSF suggests an irreversible damage to astrocytes, since glial fibrillary acidic protein is not secreted by astrocytes.

All of these markers suggest the involvement of vascular risk factors and consequent subcortical white matter lesions in the development of iNPH; however, further studies are needed to explore their predictive value in clinical application.

OTHER BIOMARKERS AND METHODOLOGICAL IMPACT ON CSF BIOMARKER DETECTION

The level of prostaglandin D synthase was found to be significantly lower in iNPH patients compared with controls and other dementia patients, such as Lewy body dementia, vascular dementia, and AD^[77]. This enzyme is secreted into CSF by the leptomeninges and the trabecular cells of the arachnoid membrane. The authors speculated that the decreased level of prostaglandin D synthase was probably due to a degenerative change of the arachnoid membrane in iNPH patients.

Finally, the methodology of CSF biomarker detection may also affect the ability to reliably evaluate biological biomarkers for the differentiation and prognosis of cognitive impairment diseases^[78]. Many factors may affect the reliability and sensitivity of biomarker detection, for example, the systematic difference between different assays, different pre-analytical protocol for sample preparation and storage, analytical variability of measurement procedures, *etc.*^[79,80]. When interpreting measurement results, these factors should be considered. In addition, some biomarkers exhibit periodic concentration patterns. Therefore, the most appropriate time for sample collection must also be considered when designing a protocol^[79].

CONCLUSION

The overlap of neuroimaging and symptomatic manifestations leads to diagnostic confusion between iNPH and other neurodegeneration diseases, such as AD and subcortical ischemic vascular disease. Despite the absence of definite pathological hallmarks, the biomarkers altered in CSF might serve as targets for diagnosis and therapeutic intervention. Furthermore, the biomarkers in CSF could reflect the adjacent cerebral pathophysiological status, therefore are potentially useful to provide insight into the pathological changes in the brain milieu and underlying pathogenesis. Although many CSF biomarkers have been analyzed in iNPH patients, the significant findings include the reduced $A\beta_{42}$ with concomitant normal or reduced t-tau and p-tau levels in iNPH coupled with reduced $A\beta_{42}$ with concomitant increased both t-tau and p-tau levels in AD. This characteristic alteration may significantly improve the accuracy of differential diagnosis between AD and iNPH patients. Other biomarkers may lack specification in differential diagnosis, but the definite changes may mirror the underlying pathogenesis mechanisms, such as demyelination, neurodegeneration, and neuroinflammation, and provide valuable information to further explore the pathogenesis mechanisms and optical therapeutic manipulations.

DECLARATIONS

Authors' contributions

Conceived of the presented idea: Yang J

Underwent literature review and synthesized a draft: Yang J, Zhang XJ, and Guo J

Contributed ideas throughout the process and approved the final draft: Yang J

Looked over and edited draft: Yang J, Zhang XJ, and Guo J

Availability of data and materials

Not applicable.

Financial support and sponsorship

None.

Conflicts of interest

All authors declared that there are no conflicts of interest.

Ethical approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Copyright

© The Author(s) 2020.

REFERENCES

1. Schirinzi T, Sancesario GM, Di Lazzaro G, D'Elia A, Imbriani P, et al. Cerebrospinal fluid biomarkers profile of idiopathic normal pressure hydrocephalus. *J Neural Transm (Vienna)* 2018;125:673-9.
2. Manniche C, Hejl AM, Hasselbalch SG, Simonsen AH. Cerebrospinal fluid biomarkers in idiopathic normal pressure hydrocephalus versus Alzheimer's disease and subcortical ischemic vascular disease: a systematic review. *J Alzheimers Dis* 2019;68:267-79.
3. Ghosh S, Lippa C. Diagnosis and prognosis in idiopathic normal pressure hydrocephalus. *Am J Alzheimers Dis Other Dement* 2014;29:583-9.
4. Keong NC, Pena A, Price SJ, Czosnyka M, Czosnyka Z, et al. Imaging normal pressure hydrocephalus: theories, techniques, and challenges. *Neurosurg Focus* 2016;41:E11.
5. Savolainen S, Hurskainen H, Paljärvi L, Alafuzoff I, Vapalahti M. Five-year outcome of normal pressure hydrocephalus with or without a shunt: predictive value of the clinical signs, neuropsychological evaluation and infusion test. *Acta Neurochir (Wien)* 2002;144:515-23.
6. Pomeraniec IJ, Bond AE, Lopes MB, Jane JA Sr. Concurrent Alzheimer's pathology in patients with clinical normal pressure hydrocephalus: correlation of high-volume lumbar puncture results, cortical brain biopsies, and outcomes. *J Neurosurg* 2016;124:382-8.
7. Graff-Radford NR. Alzheimer CSF biomarkers may be misleading in normal-pressure hydrocephalus. *Neurology* 2014;83:1573-5.
8. Pomeraniec IJ, Taylor DG, Bond AE, Lopes MB. Concurrent Alzheimer's pathology in patients with clinical normal pressure hydrocephalus. *J Neurosurg Sci* 2018.
9. McGovern RA, Nelp TB, Kelly KM, Chan AK, Mazzone P, et al. Predicting cognitive improvement in normal pressure hydrocephalus patients using preoperative neuropsychological testing and cerebrospinal fluid biomarkers. *Neurosurgery* 2019;85:E662-9.
10. Tarnaris A, Watkins LD, Kitchen ND. Biomarkers in chronic adult hydrocephalus. *Cerebrospinal Fluid Res* 2006;3:11.
11. Agren-Wilsson A, Lekman A, Sjöberg W, Rosengren L, Blennow K, et al. CSF biomarkers in the evaluation of idiopathic normal pressure hydrocephalus. *Acta Neurol Scand* 2007;116:333-9.
12. Saito M, Nishio Y, Kanno S, Uchiyama M, Hayashi A, et al. Cognitive profile of idiopathic normal pressure hydrocephalus. *Dement Geriatr Cogn Dis Extra* 2011;1:202-11.
13. Espay AJ, Da Prat GA, Dwivedi AK, Rodriguez-Porcel F, Vaughan JE, et al. Deconstructing normal pressure hydrocephalus: ventriculomegaly as early sign of neurodegeneration. *Ann Neurol* 2017;82:503-13.
14. Nassar BR, Lippa CF. Idiopathic normal pressure hydrocephalus: a review for general practitioners. *Gerontol Geriatr Med* 2016;2:2333721416643702.
15. Reeves BC, Karimy JK, Kundishora AJ, Mestre H, Cerci HM, et al. Glymphatic system impairment in Alzheimer's disease and idiopathic

- normal pressure hydrocephalus. *Trends Mol Med* 2020;26:285-95.
16. Iliff JJ, Wang MH, Zeppenfeld DM, Venkataraman A, Plog BA, et al. Cerebral arterial pulsation drives paravascular CSF-interstitial fluid exchange in the murine brain. *J Neurosci* 2013;33:18190-9.
 17. Luciano MG, Dombrowski SM, Qvarlander S, El-Khoury S, Yang J, et al. Novel method for dynamic control of intracranial pressure. *J Neurosurg* 2017;126:1629-40.
 18. Xie LL, Kang HY, Xu QW, Chen MJ, Liao YH, et al. Sleep drives metabolite clearance from the adult brain. *Science* 2013;342:373-7.
 19. Roman GC, Jackson RE, Fung SH, Zhang YJ, Verma AK. Sleep-disordered breathing and idiopathic normal-pressure hydrocephalus: recent pathophysiological advances. *Curr Neurol Neurosci Rep* 2019;19:39.
 20. Ju YE, Zangrilli MA, Finn MB, Fagan AM, Holtzman DM. Obstructive sleep apnea treatment, slow wave activity, and amyloid-beta. *Ann Neurol* 2019;85:291-5.
 21. Gao Y, Tan L, Yu JT. Tau in Alzheimer's disease: mechanisms and therapeutic strategies. *Curr Alzheimer Res* 2018;15:283-300.
 22. Schirizzi T, Sancesario GM, Ialongo C, Imbriani P, Madeo G, et al. A clinical and biochemical analysis in the differential diagnosis of idiopathic normal pressure hydrocephalus. *Front Neurol* 2015;6:86.
 23. Momjian S, Owler BK, Czosnyka Z, Czosnyka M, Pena A, et al. Pattern of white matter regional cerebral blood flow and autoregulation in normal pressure hydrocephalus. *Brain* 2004;127:965-72.
 24. Jeppsson A, Wikkelsø C, Blennow K, Zetterberg H, Constantinescu R, et al. CSF biomarkers distinguish idiopathic normal pressure hydrocephalus from its mimics. *J Neurol Neurosurg Psychiatry* 2019;90:1117-23.
 25. Moriya M, Miyajima M, Nakajima M, Ogino I, Arai H. Impact of cerebrospinal fluid shunting for idiopathic normal pressure hydrocephalus on the amyloid cascade. *PLoS One* 2015;10:e0119973.
 26. Scheltens P, Blennow K, Breteler MM, de Strooper B, Frisoni GB, et al. Alzheimer's disease. *Lancet* 2016;388:505-17.
 27. Blennow K, Biscetti L, Eusebi P, Parnetti L. Cerebrospinal fluid biomarkers in Alzheimer's and Parkinson's diseases-from pathophysiology to clinical practice. *Mov Disord* 2016;31:836-47.
 28. Leinonen V, Koivisto AM, Savolainen S, Rummukainen J, Sutela A, et al. Post-mortem findings in 10 patients with presumed normal-pressure hydrocephalus and review of the literature. *Neuropathol Appl Neurobiol* 2012;38:72-86.
 29. Silverberg GD. Normal pressure hydrocephalus (NPH): ischaemia, CSF stagnation or both. *Brain* 2004;127:947-8.
 30. Kudo T, Mima T, Hashimoto R, Nakao K, Morihara T, et al. Tau protein is a potential biological marker for normal pressure hydrocephalus. *Psychiatry Clin Neurosci* 2000;54:199-202.
 31. Magdalinou N, Lees AJ, Zetterberg H. Cerebrospinal fluid biomarkers in parkinsonian conditions: an update and future directions. *J Neurol Neurosurg Psychiatry* 2014;85:1065-75.
 32. Pyykko OT, Lumela M, Rummukainen J, Nerg O, Seppala TT, et al. Cerebrospinal fluid biomarker and brain biopsy findings in idiopathic normal pressure hydrocephalus. *PLoS One* 2014;9:e91974.
 33. Miyajima M, Nakajima M, Motoi Y, Moriya M, Sugano H, et al. Leucine-rich alpha2-glycoprotein is a novel biomarker of neurodegenerative disease in human cerebrospinal fluid and causes neurodegeneration in mouse cerebral cortex. *PLoS One* 2013;8:e74453.
 34. Malmstrom C, Haghighi S, Rosengren L, Andersen O, Lycke J. Neurofilament light protein and glial fibrillary acidic protein as biological markers in MS. *Neurology* 2003;61:1720-5.
 35. Jeppsson A, Hölttä M, Zetterberg H, Blennow K, Wikkelsø C, et al. Amyloid mis-metabolism in idiopathic normal pressure hydrocephalus. *Fluids Barriers CNS* 2016;13:13.
 36. Jeppsson A, Zetterberg H, Blennow K, Wikkelsø C. Idiopathic normal-pressure hydrocephalus: pathophysiology and diagnosis by CSF biomarkers. *Neurology* 2013;80:1385-92.
 37. Tullberg M, Blennow K, Mansson JE, Fredman P, Tisell M, et al. Ventricular cerebrospinal fluid neurofilament protein levels decrease in parallel with white matter pathology after shunt surgery in normal pressure hydrocephalus. *Eur J Neurol* 2007;14:248-54.
 38. Tullberg M, Rosengren L, Blomsterwall E, Karlsson JE, Wikkelsø C. CSF neurofilament and glial fibrillary acidic protein in normal pressure hydrocephalus. *Neurology* 1998;50:1122-7.
 39. Marcus J, Honigbaum S, Shroff S, Honke K, Rosenbluth J, et al. Sulfatide is essential for the maintenance of CNS myelin and axon structure. *Glia* 2006;53:372-81.
 40. Miller A, Glass-Marmor L, Abraham M, Grossman I, Shapiro S, et al. Bio-markers of disease activity and response to therapy in multiple sclerosis. *Clin Neurol Neurosurg* 2004;106:249-54.
 41. Lamers KJB, Vos P, Verbeek MM, Rosmalen F, van Geel WJA, et al. Protein S-100B, neuron-specific enolase (NSE), myelin basic protein (MBP) and glial fibrillary acidic protein (GFAP) in cerebrospinal fluid (CSF) and blood of neurological patients. *Brain Res Bull* 2003;61:261-4.
 42. Longatti PL, Canova G, Guida F, Carniato A, Moro M, et al. The CSF myelin basic protein: a reliable marker of actual cerebral damage in hydrocephalus. *J Neurosurg Sci* 1993;37:87-90.
 43. Nakajima M, Miyajima M, Ogino I, Watanabe M, Miyata H, et al. Leucine-rich alpha2-glycoprotein is a marker for idiopathic normal pressure hydrocephalus. *Acta Neurochir (Wien)* 2011;153:1339-46.
 44. Kempuraj D, Thangavel R, Natteru PA, Selvakumar GP, Saeed D, et al. Neuroinflammation induces neurodegeneration. *J Neurol Neurosurg Spine* 2016;1.
 45. Tarkowski E, Tullberg M, Fredman P, Wikkelsø C. Normal pressure hydrocephalus triggers intrathecal production of TNF-alpha. *Neurobiol Aging* 2003;24:707-14.
 46. Castaneyra-Ruiz L, Gonzalez-Marrero I, Carmona-Calero EM, Abreu-Gonzalez P, Lecuona M, et al. Cerebrospinal fluid levels of tumor

- necrosis factor alpha and aquaporin 1 in patients with mild cognitive impairment and idiopathic normal pressure hydrocephalus. *Clin Neurol Neurosurg* 2016;146:76-81.
47. Li X, Miyajima M, Jiang C, Arai H. Expression of TGF-betas and TGF-beta type II receptor in cerebrospinal fluid of patients with idiopathic normal pressure hydrocephalus. *Neurosci Lett* 2007;413:141-4.
48. Pfanner T, Henri-Bhargava A, Borchert S. Cerebrospinal fluid biomarkers as predictors of shunt response in idiopathic normal pressure hydrocephalus: a systematic review. *Can J Neurol Sci* 2018;45:3-10.
49. Zhang X, Huang WJ, Chen WW. TGF-beta1 factor in the cerebrovascular diseases of Alzheimer's disease. *Eur Rev Med Pharmacol Sci* 2016;20:5178-85.
50. Crawley JT, Goulding DA, Ferreira V, Severs NJ, Lupu F. Expression and localization of tissue factor pathway inhibitor-2 in normal and atherosclerotic human vessels. *Arterioscler Thromb Vasc Biol* 2002;22:218-24.
51. Sosvorova L, Vcelak J, Mohapl M, Vitku J, Bicikova M, et al. Selected pro- and anti-inflammatory cytokines in cerebrospinal fluid in normal pressure hydrocephalus. *Neuro Endocrinol Lett* 2014;35:586-93.
52. Sosvorova L, Mohapl M, Vcelak J, Hill M, Vitku J, et al. The impact of selected cytokines in the follow-up of normal pressure hydrocephalus. *Physiol Res* 2015;64:S283-90.
53. Kester MI, Teunissen CE, Sutphen C, Herries EM, Ladenson JH, et al. Cerebrospinal fluid VILIP-1 and YKL-40, candidate biomarkers to diagnose, predict and monitor Alzheimer's disease in a memory clinic cohort. *Alzheimers Res Ther* 2015;7:59.
54. Poca MA, Mataró M, Sahuquillo J, Catalán R, Ibañez J, et al. Shunt related changes in somatostatin, neuropeptide Y, and corticotropin releasing factor concentrations in patients with normal pressure hydrocephalus. *J Neurol Neurosurg Psychiatry* 2001;70:298-304.
55. Tisell M, Tullberg M, Mansson JE, Fredman P, Blennow K, et al. Differences in cerebrospinal fluid dynamics do not affect the levels of biochemical markers in ventricular CSF from patients with aqueductal stenosis and idiopathic normal pressure hydrocephalus. *Eur J Neurol* 2004;11:17-23.
56. Wikkelsö C, Ekman R, Westergren I, Johansson B. Neuropeptides in cerebrospinal fluid in normal-pressure hydrocephalus and dementia. *Eur Neurol* 1991;31:88-93.
57. Tullberg M, Mansson JE, Fredman P, Lekman A, Blennow K, et al. CSF sulfatide distinguishes between normal pressure hydrocephalus and subcortical arteriosclerotic encephalopathy. *J Neurol Neurosurg Psychiatry* 2000;69:74-81.
58. Nooijen PT, Schoonderwaldt HC, Wevers RA, Hommes OR, Lamers KJ. Neuron-specific enolase, S-100 protein, myelin basic protein and lactate in CSF in dementia. *Dement Geriatr Cogn Disord* 1997;8:169-73.
59. Tarnaris A, Toma AK, Pullen E, Chapman MD, Petzold A, et al. Cognitive, biochemical, and imaging profile of patients suffering from idiopathic normal pressure hydrocephalus. *Alzheimers Dement* 2011;7:501-8.
60. Fersten E, Gordon-Krajcer W, Glowacki M, Mroziak B, Jurkiewicz J, et al. Cerebrospinal fluid free-radical peroxidation products and cognitive functioning patterns differentiate varieties of normal pressure hydrocephalus. *Folia Neuropathol* 2004;42:133-40.
61. Abbott NJ, Ronnback L, Hansson E. Astrocyte-endothelial interactions at the blood-brain barrier. *Nat Rev Neurosci* 2006;7:41-53.
62. Chow BW, Gu C. The molecular constituents of the blood-brain barrier. *Trends Neurosci* 2015;38:598-608.
63. Janelidze S, Hertz J, Nagga K, Nilsson K, Nilsson C, et al. Increased blood-brain barrier permeability is associated with dementia and diabetes but not amyloid pathology or APOE genotype. *Neurobiol Aging* 2017;51:104-12.
64. Eide PK, Pripp AH. Increased prevalence of cardiovascular disease in idiopathic normal pressure hydrocephalus patients compared to a population-based cohort from the HUNT3 survey. *Fluids Barriers CNS* 2014;11:19.
65. Krauss JK, Regel JP, Vach W, Orszagh M, Jungling FD, et al. White matter lesions in patients with idiopathic normal pressure hydrocephalus and in an age-matched control group: a comparative study. *Neurosurgery* 1997;40:491-5.
66. Jaraj D, Agerskov S, Rabiei K, Marlow T, Jensen C, et al. Vascular factors in suspected normal pressure hydrocephalus: a population-based study. *Neurology* 2016;86:592-9.
67. Graff-Radford NR, Knopman DS, Penman AD, Coker LH, Mosley TH. Do systolic BP and pulse pressure relate to ventricular enlargement? *Eur J Neurol* 2013;20:720-4.
68. Miyajima M, Nakajima M, Ogino I, Miyata H, Motoi Y, et al. Soluble amyloid precursor protein alpha in the cerebrospinal fluid as a diagnostic and prognostic biomarker for idiopathic normal pressure hydrocephalus. *Eur J Neurol* 2013;20:236-42.
69. Mashayekhi F, Salehi Z. Expression of nerve growth factor in cerebrospinal fluid of congenital hydrocephalic and normal children. *Eur J Neurol* 2005;12:632-7.
70. Yang JT, Chang CN, Hsu YH, Wei KC, Lin TK, et al. Increase in CSF NGF concentration is positively correlated with poor prognosis of postoperative hydrocephalic patients. *Clin Biochem* 1999;32:673-5.
71. Del Bigio MR. Neuropathological changes caused by hydrocephalus. *Acta Neuropathol* 1993;85:573-85.
72. Li X, Miyajima M, Mineki R, Taka H, Murayama K, et al. Analysis of cerebellum proteomics in the hydrocephalic H-Tx rat. *Neuroreport* 2005;16:571-4.
73. Yang J, Dombrowski SM, Krishnan C, Krajcir N, Deshpande A, et al. Vascular endothelial growth factor in the CSF of elderly patients with ventriculomegaly: variability, periodicity and levels in drainage responders and non-responders. *Clin Neurol Neurosurg* 2013;115:1729-34.
74. Yang J, Shanahan KJ, Shriver LP, Luciano MG. Exercise-induced changes of cerebrospinal fluid vascular endothelial growth factor in adult chronic hydrocephalus patients. *J Clin Neurosci* 2016;24:52-6.
75. Huang H, Yang J, Luciano M, Shriver LP. Longitudinal metabolite profiling of cerebrospinal fluid in normal pressure hydrocephalus links brain metabolism with exercise-induced VEGF production and clinical outcome. *Neurochem Res* 2016;41:1713-22.
76. Bartosik-Psujek H, Stelmasiak Z. Biochemical markers of damage of the central nervous system in multiple sclerosis. *Ann Univ Mariae*

- Curie Skłodowska Med 2001;56:389-92.
77. Mase M, Yamada K, Shimazu N, Seiki K, Oda H, et al. Lipocalin-type prostaglandin D synthase (beta-trace) in cerebrospinal fluid: a useful marker for the diagnosis of normal pressure hydrocephalus. *Neurosci Res* 2003;47:455-9.
78. Shaw LM, Hansson O, Manuilova E, Masters CL, Doecke JD, et al. Method comparison study of the Elecsys(R) beta-Amyloid (1-42) CSF assay versus comparator assays and LC-MS/MS. *Clin Biochem* 2019;72:7-14.
79. Yang J, Dombrowski SM, Deshpande A, Krajcir N, El-Khoury S, et al. Stability analysis of vascular endothelial growth factor in cerebrospinal fluid. *Neurochem Res* 2011;36:1947-54.
80. Schindler SE, Gray JD, Gordon BA, Xiong C, Batrla-Utermann R, et al. Cerebrospinal fluid biomarkers measured by Elecsys assays compared to amyloid imaging. *Alzheimers Dement* 2018;14:1460-9.

Review

Open Access



Cerebrospinal fluid amyloid beta and tau proteins in atypical Parkinsonism: a review

Vasilios C. Constantinides¹, George P. Paraskevas¹, Fotini Boufidou¹, Mara Bourbouli², Panagiotis G. Paraskevas³, Leonidas Stefanis¹, Elisabeth Kapaki¹

¹1st Department of Neurology, National and Kapodistrian University of Athens, Athens, P.C. 11528, Greece.

²Department of Neurology, University of Crete, Heraklion 70013, Greece.

³Department of Nursing, Technological Educational Institute of Crete, Heraklion 71500, Greece.

Correspondence to: Dr. Vasilios C. Constantinides, 1st Department of Neurology, National and Kapodistrian University of Athens, School of Medicine, Eginition Hospital, 72 Vas. Sofias Ave, Athens 11528, Greece.

E-mail: vconstan@med.uoa.gr; vassilis.kon@hotmail.com

How to cite this article: Constantinides VC, Paraskevas GP, Boufidou F, Bourbouli M, Paraskevas PG, Stefanis L, Kapaki E. Cerebrospinal fluid amyloid beta and tau proteins in atypical Parkinsonism: a review. *Neuroimmunol Neuroinflammation* 2020;7:120-31. <http://dx.doi.org/10.20517/2347-8659.2019.22>

Received: 17 Dec 2019 **First Decision:** 2 Mar 2020 **Revised:** 25 Mar 2020 **Accepted:** 7 Apr 2020 **Available online:** 16 May 2020

Science Editor: Athanassios P. Kyritsis **Copy Editor:** Jing-Wen Zhang **Production Editor:** Jing Yu

Abstract

Progressive supranuclear palsy, corticobasal degeneration, multiple system atrophy and dementia with Lewy bodies are the most common causes of atypical Parkinsonism and enter the differential diagnosis of Parkinson's disease. Multiple system atrophy, dementia with Lewy bodies and Parkinson's disease are synucleinopathies, whereas progressive supranuclear palsy and corticobasal degeneration are tauopathies. Multiple cerebrospinal fluid markers have been applied on cohorts of patients with Parkinsonism, with the aim to develop biomarkers for these disorders. Total tau (τ_T), phosphorylated tau at threonine 181 (τ_p -181) and amyloid-beta with 42 amino acids ($A\beta_{42}$) are considered classical biomarkers for Alzheimer's disease. The aim of the present study is to review the literature regarding these classical cerebrospinal fluid biomarkers in cohorts with Parkinsonism, as well as present data on novel approaches regarding analysis of these proteins.

Keywords: Biomarkers, cerebrospinal fluid, progressive supranuclear palsy, corticobasal degeneration, multiple system atrophy, dementia with Lewy bodies, tau protein, phosphorylated tau protein, amyloid beta



© The Author(s) 2020. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, sharing, adaptation, distribution and reproduction in any medium or format, for any purpose, even commercially, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.



INTRODUCTION

Parkinson's disease (PD) is the most common neurodegenerative movement disorder. The pathologic hallmarks of PD are Lewy bodies, which consist of intraneuronal cytoplasmic depositions of pathological α -synuclein^[1]. Thus, PD is considered a synucleinopathy. Clinical diagnosis of PD is straightforward in typical cases. However, it can be problematic in patients with atypical clinical features. Accuracy of clinical diagnosis of PD is suboptimal, since as many as 25% of patients can be misdiagnosed^[2].

Progressive supranuclear palsy (PSP), corticobasal degeneration (CBD), multiple system atrophy (MSA) and dementia with Lewy bodies (DLB) are the most common causes of atypical Parkinsonism. All of these diseases manifest with Parkinsonism, which is poorly or only transiently responsive to dopaminergic treatment.

MSA is a synucleinopathy, like PD. Its pathologic hallmark is glial cytoplasmic inclusions, which consist of abnormal α -synuclein deposition in oligodendrocytes^[3]. DLB is also a synucleinopathy, characterized by predominantly cortical Lewy bodies^[4]. PSP and CBD on the other hand are considered tauopathies, since their main pathologic findings (tufted astrocytes and astrocytic plaques, respectively) consist of abnormal tau protein aggregates in astrocytes^[5,6].

Tau protein can present in six isoforms, depending on the alternate splicing of the microtubule associated protein tau (*MAPT*) gene^[7]. This results in the variable expression of no, one or two oligonucleotides (N1 and N2) coded by exons 2 and 3 of the *MAPT* gene, as well as the presence of 3-repeat (3R) or 4-repeat (4R) microtubule binding regions coded by exon 10. Depending on the predominance of 3R- or 4R-tau isoforms, tauopathies are further divided into 4R- or 3R-tauopathies. PSP and CBD are considered 4-repeat (4R) tauopathies, Alzheimer's disease (AD) is a mixed 3R- and 4R-tauopathy, whereas Pick's disease is a 3R-tauopathy.

Total tau protein (τ_T), phosphorylated tau protein at threonine 181 (τ_{P-181}) and amyloid beta with 42 amino acids ($A\beta_{42}$) are well-characterized cerebrospinal fluid (CSF) biomarkers of AD. These biomarkers have been incorporated into the most recent AD diagnostic criteria^[8] and are the basis of the recently proposed AT(N) taxonomy system, which introduces biomarkers (according to their molecular specificity) for the *in vivo* pathological characterization of patients with AD^[9].

The aim of this review is to present data on the utility of these three classical CSF biomarkers (τ_T , τ_{P-181} and $A\beta_{42}$) in the differential diagnosis of atypical Parkinsonism from PD. To this end, only studies which included patients with atypical Parkinsonism vs. PD or healthy controls are included.

This review includes representative studies which either have established our current knowledge on CSF biomarkers in Parkinsonism or provide new insights on the subject [Table 1].

STUDIES IN PSP

Most studies do not report any differences in CSF $A\beta_{42}$ between PSP and other causes of Parkinsonism or controls^[10-16]. Interestingly, however, some studies have reported lower $A\beta_{42}$ values in PSP compared to controls^[17-19]. Moreover, a single study reported lower $A\beta_{42}$ levels in PSP vs. PD^[19]. According to this study, $A\beta_{42}$ could differentiate PSP from PD with 83% sensitivity and 67% specificity.

Regarding CSF τ_T , several studies could not establish any differences between PSP patients and controls^[13-17]. Likewise, no difference in τ_T was evident between PSP and CBD^[13,15-17], MSA^[13,15,16] or PD^[15,16] in most studies. Two studies have supported that PSP patients exhibit lower τ_T levels when compared to

Table 1. Overview of the studies with data on classical CSF biomarkers in cohorts with Parkinsonism

	A β ₄₂	τ _T	τ _{P-181}	Comparison group
Studies in PSP				
Holmberg <i>et al.</i> ^[10]	↔			Ctrl, MSA, PD
Verbeek <i>et al.</i> ^[11]	↔			Ctrl, MSA, PD
Aerts <i>et al.</i> ^[12]	↔ ^a	↓ ^b ↑ ^c	↔ ^a	a. CBD, PD, ctrl b. CBD c. PD
Hall <i>et al.</i> ^[13]	↔ ^a ↑ ^b	↔ ^c	↓ ^d	a. MSA, CBD, PD, PDD, ctrl b. DLB c. ctrl, CBD, MSA d. ctrl
Schoonenboom <i>et al.</i> ^[14]	↔	↔	↔	DLB, CBD, ctrl
Magdalinou <i>et al.</i> ^[15]	↔	↔	↔	CBD, MSA, PD, ctrl
Constantinides <i>et al.</i> ^[16]	↔	↔	↔	MSA, PD, CBD, ctrl
Noguchi <i>et al.</i> ^[17]	↓ ^a	↔ ^b	↔ ^b	a. ctrl b. ctrl, CBD
Wagshal <i>et al.</i> ^[18]	↓	↓	↔	ctrl
Schirinzi <i>et al.</i> ^[19]	↓	↓	↓	ctrl
Süssmuth <i>et al.</i> ^[20]	↔	↔	↔	MSA, PD
Studies in CBD				
Aerts <i>et al.</i> ^[12]	↔ ^a	↑ ^a	↑ ^b ↔ ^c	a. ctrl, PSP, PD b. PD c. ctrl, PSP
Constantinides <i>et al.</i> ^[16]	↓ ^a	↑ ^b ↔ ^c	↔ ^d	a. PD b. ctrl c. PSP, MSA, PD d. PSP, MSA, PD, ctrl
Mitani <i>et al.</i> ^[21]		↑		ctrl
Urakami <i>et al.</i> ^[22]		↑		ctrl, PSP
Urakami <i>et al.</i> ^[23]		↑		ctrl, PSP
Hall <i>et al.</i> ^[13]	↔ ^a ↑ ^b	↑ ^c ↔ ^d	↔ ^e ↑ ^f	a. MSA, PSP, PD, PDD, ctrl b. DLB c. PD d. PSP e. PSP, PD, ctrl f. MSA
Schoonenboom <i>et al.</i> ^[14]	↓	↔	↔	ctrl
Magdalinou <i>et al.</i> ^[15]	↔	↔	↔	MSA, PSP, PD, ctrl
Noguchi <i>et al.</i> ^[17]	↔ ^a ↓ ^b	↔ ^c	↔ ^c	a. PSP b. ctrl c. ctrl, PSP
Studies in MSA				
Holmberg <i>et al.</i> ^[10]	↓			PD, PSP, ctrl
Shi <i>et al.</i> ^[26]	↓ ^a ↔ ^b	↔ ^c	↔ ^c	a. ctrl b. PD c. ctrl, PD
Verbeek <i>et al.</i> ^[11]	↔			Ctrl, PD
Hall <i>et al.</i> ^[13]	↔ ^a ↑ ^b	↑ ^c	↓ ^d	a. MSA, CBD, PD, PDD, ctrl b. DLB c. PD, PDD d. PD, ctrl
Magdalinou <i>et al.</i> ^[15]	↔	↔	↔	CBD, PSP, PD, ctrl
Constantinides <i>et al.</i> ^[16]	↔	↔	↔	PSP, CBD, PD, ctrl
Süssmuth <i>et al.</i> ^[20]	↔ ^a	↑ ^b	↔ ^c	a. PD, PSP b. ctrl, PD c. PSP, PD, ctrl
Mollenhauer <i>et al.</i> ^[27]	↔	↔		PSP, PD
Abdo <i>et al.</i> ^[28]		↑		ctrl
Herbert <i>et al.</i> ^[29]		↑		Ctrl, PD
Abdo <i>et al.</i> ^[30]		↑		PD
Herbert <i>et al.</i> ^[31]		↑		PD
Studies in DLB				
Schoonenboom <i>et al.</i> ^[14]	↓ ^a ↑ ^b	↓ ^b ↑ ^a	↓ ^b ↑ ^a	a. ctrl b. AD

Mollenhauer <i>et al.</i> ^[32]	↓ ^a ↔ ^b	↓ ^b ↔ ^a	↑ ^a	a. ctrl b. AD
Mollenhauer <i>et al.</i> ^[33]	↓ ^a ↔ ^b	↓ ^b ↑ ^a	↔ ^b	a. ctrl b. AD
Mulugeta <i>et al.</i> ^[34]	↓ ^a ↔ ^b	↔ ^c	↔ ^c	a. ctrl b. AD, PDD c. ctrl, PDD
Llorens <i>et al.</i> ^[35]	↓ ^a	↑ ^b		a. ctrl, PD, PDD b. PD, MSA
Van Steenoven <i>et al.</i> ^[36]	↓			ctrl
Parnetti <i>et al.</i> ^[37]	↓ ^a ↔ ^b	↓ ^b ↑ ^c	↓ ^b	a. PD, PDD b. AD c. PD
Kaerst <i>et al.</i> ^[38]	↓	↑		PD
Gmitterova <i>et al.</i> ^[39]	↓	↑		PDD
Gómez-Tortosa <i>et al.</i> ^[40]	↑ ^a ↔ ^b	↓ ^a ↔ ^b		a. AD b. ctrl
Boström <i>et al.</i> ^[41]	↑ ^a	↓ ^a	↑ ^b	a. AD b. ctrl
Abdelnour <i>et al.</i> ^[42]	↑			ctrl
Mollenhauer <i>et al.</i> ^[43]		↔		PDD

↑: elevated CSF levels; ↓: decreased CSF levels; ↔: no difference in CSF levels; CSF: cerebrospinal fluid; AD: Alzheimer's disease; PD: Parkinson's disease; MSA: multiple system atrophy; PSP: progressive supranuclear palsy; CBD: corticobasal degeneration; DLB: dementia with Lewy bodies; PDD: parkinson's disease dementia

controls^[18,19], whereas a single study posited that patients with PSP have lower τ_T levels compared to CBD, but higher τ_T levels compared to PD^[12].

CSF τ_{P-181} does not seem to be useful in the differentiation of PSP from other causes of Parkinsonism or controls^[12,14-18]. Two studies have reported lower τ_{P-181} levels compared to controls^[13,19].

Few studies include data on ratios of CSF biomarkers. In a large study, which included PSP, CBD and AD patients, elevated $\tau_{P-181}/A\beta_{42}$ ratio values could differentiate AD from PSP and CBD^[17]. Another study posited that lower τ_{P-181}/τ_T ratio values could discriminate patients with atypical Parkinsonism (PSP and MSA) from PD^[20]. This ratio was optimal for discriminating PSP from CBD with a reported sensitivity of 86% and specificity of 75%^[12]. PSP patients could also be differentiated from controls by lower $A\beta_{42}/\tau_T$ ^[12] and higher τ_T/τ_{P-181} ^[16] ratios.

A single study examined classical CSF biomarkers in different phenotypes of PSP. To this end, patients with classic Richardson's syndrome (RS) were compared to patients with PSP-Parkinsonism (PSP-P). Interestingly, only patients with PSP-P had elevated total τ_T levels, compared to RS, PD and controls. $A\beta_{42}$ on the other hand was significantly lower in RS patients compared to PSP-P^[20].

Few PSP patients seem to harbor a CSF-AD profile (generally defined as decreased $A\beta_{42}$ with elevated τ_T or τ_{P-181}). In a large cohort, including diverse neurodegenerative disorders, 10% of PSP patients had a CSF-AD profile, as determined by an index incorporating CSF $A\beta_{42}$ and τ_{P-181} values^[14]. Likewise, only one of 19 PSP patients (~5%) had a typical CSF-AD profile, as determined by abnormal $A\beta_{42}$, τ_T and τ_{P-181} values, in a cohort of patients with Parkinsonism^[16].

Lastly, the possible relationship between CSF biomarkers and clinical characteristics has also been extensively studied in PSP. Most studies agree that there is no clinical-biochemical correlation in PSP^[12-15,17]. Two studies have correlated low $A\beta_{42}$ levels with higher disease severity, as measured by Hoehn and Yahr score^[10] or the PSP Rating Scale^[19].

As a general conclusion, there are no significant abnormalities in the classical CSF biomarkers in PSP. Despite being a 4R-tauopathy, CSF tau proteins do not seem to differ between PSP and other Parkinsonian disorders. CSF total tau protein, which is considered a non-specific marker of neurodegeneration, would be expected to increase in PSP, as is the case in AD. A plausible explanation would be that tau protein in PSP is concentrated intracellularly (in astrocytic plaques, tufted astrocytes and neurofibrillary tangles), and for this reason it does not enter the cerebrospinal fluid.

Additionally, no obvious clinical-biochemical correlation has been observed, while an AD biomarker profile in PSP (indicating an AD co-occurrence or possibly a PSP-like phenotype of AD) is a rarity. The presence of a CSF-AD profile in PSP patients is more likely to be indicative of the presence of dual underlying pathologies (AD and PSP), since pathological studies have not established a correlation of AD pathology and Richardson syndrome (the most common phenotype of PSP).

STUDIES IN CBD

Most studies have focused on CSF tau proteins in CBD, since CBD is a tauopathy. Several of these studies have reported elevated levels of τ_T in CBD, compared to controls^[12,16,21-23], PSP^[12,22,23] and PD^[12,13]. Interestingly, τ_T could differentiate CBD from PSP with 80% specificity and sensitivity in one study^[23]. Likewise, CSF τ_T provided 75% sensitivity and 90% specificity in the discrimination of CBD from PD in another study^[12]. However, no difference in τ_T between CBD and controls^[14,15,17], PSP^[13,15,16,20], MSA^[15,16] and PD^[15,16] has also been reported.

Regarding CSF τ_{P-181} , most studies did not establish any meaningful difference between CBD patients and other Parkinsonian disorders or controls^[13-17]. A single study has reported elevated CSF τ_{P-181} levels in CBD compared to PD^[12], and another study in CBD patients compared to MSA patients^[13].

CSF $A\beta_{42}$ levels do not differ in CBD compared to other Parkinsonian disorders, according to several studies^[12,13,15,17]. However, lower $A\beta_{42}$ has been described in CBD patients compared to controls^[14,17] and PD^[16].

Regarding CSF biomarker indices, few studies have included relevant data in CBD. One study reported decreased $A\beta_{42}/\tau_T$ ratio in CBS compared to PD and controls, whereas τ_{P-181}/τ_T ratio in CBS was decreased compared to controls^[12]. According to this study, τ_{P-181}/τ_T was optimal for PSP vs. CBD discrimination (86% sensitivity and 75% specificity). Another study posited that CBD patients have elevated $\tau_T/A\beta_{42}$ and $\tau_{P-181}/A\beta_{42}$ compared to PD^[16].

Clinical and demographic data do not correlate with levels of classical CSF biomarkers in CBD, according to most relevant studies^[13-15]. A single study reported that cognitive status, as measured by MMSE, correlated with both τ_T and τ_{P-181} levels and was inversely correlated with $A\beta_{42}$ levels^[12]. Intriguingly, τ_T levels in CBD and PSP were dependent on disease severity in a single study, with maximal levels in medium-stage disease^[23].

Few studies have focused on the presence of a CSF-AD profile in cohorts of Parkinsonian disorders. According to this approach, patients are divided in those harboring a CSF-AD profile and those who do not harbor such a profile. An initial study concluded that 20% of CBS patients harbored a CSF-AD profile (as defined by $\tau_T > 400$ pg/mL, $A\beta_{42} < 400$ pg/mL and $\tau_T/A\beta_{42}$ ratio > 1)^[24]. In another cohort, 38% of CBS patients had a CSF-AD profile, based on an index which included τ_{P-181} and $A\beta_{42}$ values^[14]. Along the same lines, a third study posited that ~30% of CBS patients had a typical CSF-AD profile (elevated τ_T and τ_{P-181} combined with decreased $A\beta_{42}$)^[16]. Interestingly, when patients with a CSF-AD profile were excluded from analyses, an initially reported elevated τ_T and decreased $A\beta_{42}$ protein in CBS disappeared, and CBS patients

no longer differed from PD or controls. This implied that the elevated τ_T and decreased $A\beta_{42}$ levels in CBS might have been the result of the inclusion of AD patients who presented with a CBS phenotype. The CSF profile was implemented in another study by the same study group to investigate possible differences between AD and non-AD pathology in a CBS cohort^[25].

In conclusion, abnormalities in classical CSF biomarkers are common in CBD. The commonly reported elevation in τ_T and τ_{P-181} and decrease in $A\beta_{42}$, does not seem to be a biochemical fingerprint of CBD but rather is a result of the inclusion in analyses of patients with AD pathology and a CBS phenotype. Up to 30% of CBS patients have an AD-biomarker profile, which is in accordance with pathological studies. Thus, CSF biomarkers are particularly useful tools in the *in vivo* discrimination of corticobasal syndrome in CBS-non-AD and CBS-AD.

STUDIES IN MSA

An initial study on CSF $A\beta_{42}$ in MSA reported lower levels compared to PD, PSP and controls^[10]. A different study group also found lower $A\beta_{42}$ levels in MSA compared to controls^[26]. However, several other studies could not establish any meaningful difference in $A\beta_{42}$ levels in MSA compared to other Parkinsonian disorders or controls^[11,13,15,16,20,26,27].

Results regarding τ_T in MSA are conflicting, with most studies reporting an increase of τ_T in MSA compared to either controls^[20,28,29] or other Parkinsonian disorders^[13,20,29-31]. However, some studies have reported no difference between MSA and other Parkinsonian disorders^[16,26,27], whereas a single study has supported that MSA patients exhibit lower τ_T compared to controls^[26].

Few studies have included τ_{P-181} data in the comparison of biomarkers in MSA and other Parkinsonian disorders. These studies did not find any difference in τ_{P-181} levels in MSA compared to other study groups^[15,16,20,26]. A single study has supported lower τ_{P-181} levels in MSA compared to PD and controls^[13].

Two studies have included CSF biomarker indices in the differential diagnosis of MSA from related disorders. More specifically, a study has supported that MSA patients have significantly lower τ_{P-181}/τ_T ratios compared to PD^[20]. Another study posited that higher values of $\tau_T/A\beta_{42}$ ratio could differentiate MSA from PD with high specificity but only moderate sensitivity^[16].

Regarding disease subtypes, MSA-C and MSA-P patients did not differ in their CSF biochemical profile in all relevant studies^[11,20,28,30].

Clinical-biochemical correlation studies in MSA are sparse. A study correlated disease severity in MSA with lower CSF $A\beta_{42}$ levels^[10]. Another study supported that both τ_T and τ_{P-181} levels increased with age in MSA^[26]. A study implementing a battery of CSF biomarkers in a cohort of diverse neurodegenerative disorders could not establish any correlation between CSF biomarkers and clinical characteristics^[15]. In conclusion, data on classical CSF biomarkers are largely inconclusive.

STUDIES IN DLB

It is well documented that CSF $A\beta_{42}$ in DLB is decreased compared to controls^[14,32-36], PD^[35,37,38] and parkinson's disease dementia (PDD)^[35,37,39] according to the majority of studies on the subject. Comparison of $A\beta_{42}$ levels between AD and DLB has yielded conflicting results, with some studies reporting greater $A\beta_{42}$ levels in DLB compared to AD^[14,40,41], and other studies not reporting any significant difference between the two groups^[32-34,37]. A single study reported greater $A\beta_{42}$ values in DLB compared to controls^[42] and another study did not find any difference between the two groups^[40]. Another study did not report a difference between DLB and PDD^[34].

There is consensus among studies that CSF τ_T is increased in AD compared to DLB^[14,32,33,37,40,41]. DLB patients have been reported to harbor elevated τ_T levels compared to controls^[14,33], although several studies could not establish any difference between DLB and controls^[32,34,40]. Regarding other synucleinopathies (PD, PDD and MSA), studies have produced mixed results, reporting either an increase in τ_T in DLB compared to PD/PDD^[35,37-39] and MSA^[35] or no difference among patient groups^[34,43].

Few studies have included τ_{P-181} in analyses of DLB cohorts. According to these limited studies, τ_{P-181} is elevated in AD compared to DLB^[14,37], although one study could not establish a difference between AD and DLB^[33]. Comparison between DLB and controls regarding τ_{P-181} has produced mixed results, with some studies reporting an increase in τ_{P-181} in DLB^[14,32,41], whereas other studies could not establish any difference between groups^[34]. PD/PDD patients do not differ from DLB in their τ_{P-181} profile^[34,44].

Lower CSF $A\beta_{42}$ levels in DLB correlate with a worse outcome, according to several studies^[14,40,42,45]. Likewise, elevated τ_T levels correlate with a poorer prognosis in DLB^[37-39,41].

Several studies in DLB, which contain longitudinal data regarding CSF biomarker level alterations over time, provide conflicting results. Two studies have reported stable $A\beta_{42}$ over time in cohorts of DLB patients^[39,46]. According to these studies, CSF τ_T and τ_{P-181} levels either increase^[39] or decrease over time^[46]. Conversely, two other studies have provided a different profile, with unaltered τ_T and τ_{P-181} levels over time, combined with a decrease in $A\beta_{42}$ during the transition from prodromal to demented stage of DLB^[47,48].

A CSF-AD profile (as defined by a decrease in $A\beta_{42}$ and an increase in τ_T and/or τ_{P-181}) is common in DLB patients, presenting in 25%-50% of patients^[14,42,45,49]. This profile invariably correlates with poorer prognosis^[42,45]. According to a recent study, as many as 85% of DLB patients have decreased $A\beta_{42}$ in CSF, either isolated (45%) or in the context of a CSF-AD profile (40%)^[49].

A single study has provided insight into the relationship between pathological finding and CSF biochemical profile^[44]. In this cohort, 72% of DLB patients had senile plaques and 50% had neurofibrillary tangles. Interestingly, there was a correlation between the presence of senile plaques and CSF $A\beta_{42}$ levels. The co-occurrence of neurofibrillary tangles did not affect τ_T and τ_{P-181} levels in DLB patients.

In conclusion, there is a general agreement that $A\beta_{42}$ levels are decreased in DLB, either alone or accompanied by increased τ_T or τ_{P-181} levels. The decrease of $A\beta_{42}$ levels have been shown to correlate with faster disease progression. Decreased CSF $A\beta_{42}$ levels and increased τ_T and τ_{P-181} in DLB could be attributed to the frequent co-occurrence of AD pathology in patients with Lewy body disease. This hypothesis would explain the correlation of a CSF-AD profile with worse outcomes in DLB patients. Intriguingly, multiple lines of evidence suggest an interplay between α -synuclein and tau/amyloid beta aggregation, which may explain the frequent co-occurrence of AD pathology and Lewy body disease^[50-52].

METHODOLOGICAL CONSIDERATIONS

Most studies on classical biomarkers in atypical Parkinsonism implemented enzyme-linked immunosorbent assays (ELISAs). Advantages of ELISAs include relatively low cost, high reproducibility and high availability, since commercial ELISAs of classical CSF biomarkers are readily available. There have been efforts, however, to implement novel techniques and to target different proteins in search of new biomarkers in neurodegenerative diseases.

To this end, a study examined differences in tau isoforms in CSF, which may reflect differences in post-translational processing of tau protein. More specifically, differences in tau proteolytic products in CSF were the main study endpoint. By means of immunoprecipitation, extended (55 kDa) and truncated forms

(33 kDa) of tau protein were determined in CSF. The tau 33 kDa/55 kDa ratio was significantly lower in PSP patients, compared to patients with AD, FTD, CBD, PD and DLB and emerged as a possible biomarker for PSP^[53]. The same study group replicated these results in a larger cohort^[54]. However, these promising results were not be replicated by another study group^[55].

In another study, immuno-PCR essays were developed to measure 3R- and 4R-tau isoforms in CSF. These essays were tested in four different cohorts, which included PSP, CBD, PDD and PDD patients. The study we are referring to included four cohorts, in which the 3R and 4R-tau isoform levels varied considerably among the four cohorts (mean values in control groups ranging for both 3R- and 4R-tau from < 5 pg/mL to ~50 pg/mL). Analysis of tau isoforms in all cohorts combined indicated lower 4R-tau levels in PSP and AD. 3R-tau did not differ among study groups. The 3R-tau/ τ_T and 3R-tau/ τ_{P-181} ratios were decreased in AD compared to PDD. The 4R-tau/ τ_T was lower in PSP and AD compared to controls and the 4R-tau/ τ_{P-181} ratio was higher in PDD compared to PSP, CBD and AD^[56].

Wagshal *et al.*^[18] developed a multitude of novel ELISAs, which target different epitopes of tau proteins, in order to examine differences in the concentration of diverse tau protein fragments in CSF. These ELISAs were applied in patients with AD, PSP and controls. Interestingly, PSP patients had lower concentrations of most tau fragments compared to controls.

Another study group implemented mass spectrometry to quantify tau-specific peptides in the entire sequence of tau protein in a cohort of PSP, AD and DLB. This novel approach yielded data for 18 tau peptides. By use of these peptides, the authors determined that the 1N and 3R-tau isoforms were mostly represented in CSF. Levels of tau peptides were intra-correlated and significantly increased in AD patients. Interestingly, AD patients had relatively decreased levels of peptides in the central core region of tau protein. This region contains phosphorylation sites, which may explain this finding^[57].

Along the same lines, Cicognola *et al.*^[58] developed novel antibodies against tau fragments at amino acid 123 and 224 (tau-123 and tau-224, respectively), after having identified endogenous tau fragments ending at these amino acids. They concluded that anti-tau-224, but not anti-tau-123, was present in neurofibrillary tangles and neuropil threads of AD patients, whereas tau-224 levels were elevated in CSF of these patients and correlated with conversion from mild cognitive impairment to AD. The authors concluded that only tau-224 is neuron-specific, whereas tau-123 may represent a general non-specific marker of tau metabolism.

DISCUSSION

There has been extensive research on classical CSF biomarkers in patients with Parkinsonism, as illustrated in this review of the literature. Most studies agree that PSP patients do not exhibit a specific CSF biochemical profile. Few studies have reported decreased $A\beta_{42}$ levels. A CSF-AD profile has rarely been described in PSP patients.

An elevation in CSF τ_T levels has been systematically reported in CBS. Moreover, few studies have also documented a decrease in $A\beta_{42}$ and an increase in τ_{P-181} in these patients. Interestingly, a significant percentage (< 40%) of CBS patients can harbor a CSF-AD profile^[14,16,24]. This is in agreement with clinical-pathological studies of CBS^[59-62], where AD is a relatively common underlying pathology of CBS. A study emphasized that the frequently reported elevation in CSF τ_T and reduction in $A\beta_{42}$ in CBS might not represent an inherent biochemical characteristic of CBS and could be attributed to the inclusion of AD patients presenting with CBS^[16]. Thus, classical CSF biomarkers are particularly important in CBS cohorts, since they can indicate an underlying AD pathology.

Regarding MSA, most studies report an elevation in CSF τ_T protein, whereas a decrease in $A\beta_{42}$ levels in MSA has been rarely reported. Although an interplay between α -synuclein and $A\beta_{42}$ has been established

in vitro, alterations in CSF $A\beta_{42}$ levels in a synucleinopathy are difficult to explain^[63]. In a transgenic mouse model with expression of both β -amyloid peptides and synuclein, β -amyloid peptides promoted aggregation of α -synuclein^[50]. Likewise, pathological data from PD and DLB patients support the accumulation of tau oligomers in these brains. Moreover, oligomeric tau forms may co-exist in the same aggregates with α -synuclein, forming hybrid oligomers^[51]. Along the same lines, α -synuclein may induce specific toxic tau oligomers in cell cultures^[52].

DLB patients exhibit significantly lower CSF $A\beta_{42}$ values compared to other Parkinsonian disorders. Moreover, studies support that τ_T in DLB may be elevated. However, AD can be differentiated from DLB by means of a significant increase in τ_T and τ_{P-181} in AD compared to DLB. A significant percentage of DLB patients harbor a CSF-AD profile, which indicates the frequent co-occurrence of Lewy body and AD pathologies, especially in older patients.

CONCLUSIONS AND FUTURE DIRECTIONS

Classical CSF biomarkers in atypical Parkinsonism are important, particularly for the diagnostic work-up of CBS and DLB patients. A CSF-AD profile in a CBS patient indicates an underlying AD pathology. Conversely, in DLB patients, the presence of a CSF-AD profile indicates the co-occurrence of AD and Lewy body pathologies, which usually correlates with poorer prognosis. Although some differences have been reported in classical CSF biomarkers in both PSP and MSA, the results are inconsistent and require further research.

The disparity of results on classical biomarkers in atypical Parkinsonism can be largely attributed to the great heterogeneity of studies. This heterogeneity refers to diagnostic criteria, cohort synthesis, statistical analysis and pre-analytical and analytical factors. Standardization of these pre-analytical and analytical confounders, as established in recent recommendations, is paramount for more robust results^[64]. Moreover, large cohorts of Parkinsonian disorders with CSF biomarker data and pathology-confirmed diagnoses are lacking. These studies would reliably inform us on the interplay between CSF biomarkers and pathological findings. This disparity in results also illustrates the limitations of classical biomarkers in the differential diagnosis of patients with Parkinsonism, and emphasizes the need for novel approaches.

To this end, differences in isoforms of these known proteins (e.g., 3R- vs. 4R-tau protein), differences in proteolytic products and quantification of specific peptides of these proteins are promising new approaches. Moreover, better characterization of other proteins which may serve as biomarkers (such as α -synuclein or TDP-43) is of great importance.

DECLARATIONS

Authors' contributions

Conception of the study, drafting of the manuscript, literature review: Constantinides VC

Conception of the study, critical appraisal of manuscript: Paraskevas GP, Boufidou F

Conception of the study, critical appraisal of manuscript, literature review: Bourbouli M, Paraskevas PG

Conception of the study, critical appraisal of manuscript: Stefanis L, Kapaki E

Availability of data and materials

Not applicable.

Financial support and sponsorship

None.

Conflicts of interest

All authors declared that there are no conflicts of interest.

Ethical approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Copyright

© The Author(s) 2020.

REFERENCES

1. Spillantini MG, Schmidt ML, Lee VM, Trojanowski JQ, Jakes R, et al. Alpha-synuclein in Lewy bodies. *Nature* 1997;388:839-40.
2. Hughes AJ, Daniel SE, Kilford L, Lees AJ. Accuracy of clinical diagnosis of idiopathic Parkinson's disease: a clinico-pathological study of 100 cases. *J Neurol Neurosurg Psychiatry* 1992;55:181-4.
3. Trojanowski JQ, Revesz T; Neuropathology Working Group on MSA. Proposed neuropathological criteria for the post mortem diagnosis of multiple system atrophy. *Neuropathol Appl Neurobiol* 2007;33:615-20.
4. McKeith IG, Galasko D, Kosaka K, Perry EK, Dickson DW, et al. Consensus guidelines for the clinical and pathologic diagnosis of dementia with Lewy bodies (DLB): report of the consortium on DLB international workshop. *Neurology* 1996;47:1113-24.
5. Hauw JJ, Daniel SE, Dickson D, Horoupian DS, Jellinger K, et al. Preliminary NINDS neuropathologic criteria for Steele-Richardson-Olszewski syndrome (progressive supranuclear palsy). *Neurology* 1994;44:2015-9.
6. Dickson DW, Bergeron C, Chin SS, Duyckaerts C, Horoupian D, et al. Office of rare diseases neuropathologic criteria for corticobasal degeneration. *J Neuropathol Exp Neurol* 2002;61:935-46.
7. Goedert M, Spillantini MG, Jakes R, Rutherford D, Crowther RA. Multiple isoforms of human microtubule-associated protein tau: sequences and localization in neurofibrillary tangles of Alzheimer's disease. *Neuron* 1989;3:519-26.
8. McKhann GM, Knopman DS, Chertkow H, Hyman BT, Jack CR Jr, et al. The diagnosis of dementia due to Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimers Dement* 2011;7:263-9.
9. Jack CR Jr, Bennett DA, Blennow K, Carrillo MC, Dunn B, et al. NIA-AA Research Framework: Toward a biological definition of Alzheimer's disease. *Alzheimers Dement* 2018;14:535-62.
10. Holmberg B, Johnels B, Blennow K, Rosengren L. Cerebrospinal fluid Abeta42 is reduced in multiple system atrophy but normal in Parkinson's disease and progressive supranuclear palsy. *Mov Disord* 2003;18:186-90.
11. Verbeek MM, Abdo WF, De Jong D, Horstink MW, Kremer BP, et al. Cerebrospinal fluid Abeta42 levels in multiple system atrophy. *Mov Disord* 2004;19:238-40.
12. Aerts MB, Esselink RA, Bloem BR, Verbeek MM. Cerebrospinal fluid tau and phosphorylated tau protein are elevated in corticobasal syndrome. *Mov Disord* 2011;26:169-73.
13. Hall S, Öhrfelt A, Constantinescu R, Andreasson U, Surova Y, et al. Accuracy of a panel of 5 cerebrospinal fluid biomarkers in the differential diagnosis of patients with dementia and/or parkinsonian disorders. *Arch Neurol* 2012;69:1445-52.
14. Schoonenboom NS, Reesink FE, Verwey NA, Kester MI, Teunissen CE, et al. Cerebrospinal fluid markers for differential dementia diagnosis in a large memory clinic cohort. *Neurology* 2012;78:47-54.
15. Magdalinou NK, Paterson RW, Schott JM, Fox NC, Mummery C, et al. A panel of nine cerebrospinal fluid biomarkers may identify patients with atypical parkinsonian syndromes. *J Neurol Neurosurg Psychiatry* 2015;86:1240-7.
16. Constantinides VC, Paraskevas GP, Emmanouilidou E, Petropoulou O, Bougea A, et al. CSF biomarkers beta-amyloid, tau proteins and a-synuclein in the differential diagnosis of Parkinson-plus syndromes. *J Neurol Sci* 2017;382:91-5.
17. Noguchi M, Yoshita M, Matsumoto Y, Ono K, Iwasa K, et al. Decreased beta-amyloid peptide42 in cerebrospinal fluid of patients with progressive supranuclear palsy and corticobasal degeneration. *J Neurol Sci* 2005;237:61-5.
18. Wagshal D, Sankaranarayanan S, Guss V, Hall T, Berisha F, et al. Divergent CSF tau alterations in two common tauopathies: Alzheimer's disease and progressive supranuclear palsy. *J Neurol Neurosurg Psychiatry* 2015;86:244-50.
19. Schirizzi T, Sancesario GM, Di Lazzaro G, Scalise S, Colona VL, et al. Clinical value of CSF amyloid-beta-42 and tau proteins in Progressive Supranuclear Palsy. *J Neural Transm (Vienna)* 2018;125:1373-9.
20. Süssmuth SD, Utner I, Landwehrmeyer B, Pinkhardt EH, Brettschneider J, et al. Differential pattern of brain-specific CSF proteins tau and amyloid-beta in Parkinsonian syndromes. *Mov Disord* 2010;25:1284-8.
21. Mitani K, Furiya Y, Uchihara T, Ishii K, Yamanouchi H, et al. Increased CSF tau protein in corticobasal degeneration. *J Neurol* 1998;245:44-6.
22. Urakami K, Mori M, Wada K, Kowa H, Takeshima T, et al. A comparison of tau protein in cerebrospinal fluid between corticobasal degeneration and progressive supranuclear palsy. *Neurosci Lett* 1999;259:127-9.

23. Urakami K, Wada K, Arai H, Sasaki H, Kanai M, et al. Diagnostic significance of tau protein in cerebrospinal fluid from patients with corticobasal degeneration or progressive supranuclear palsy. *J Neurol Sci* 2001;183:95-8.
24. Borroni B, Premi E, Agosti C, Alberici A, Cerini C, et al. CSF Alzheimer's disease-like pattern in corticobasal syndrome: evidence for a distinct disorder. *J Neurol Neurosurg Psychiatry* 2011;82:834-8.
25. Constantinides VC, Paraskevas GP, Efthymiopoulou E, Stefanis L, Kapaki E. Clinical, neuropsychological and imaging characteristics of Alzheimer's disease patients presenting as corticobasal syndrome. *J Neurol Sci* 2019;398:142-7.
26. Shi M, Bradner J, Hancock AM, Chung KA, Quinn JF, et al. Cerebrospinal fluid biomarkers for Parkinson disease diagnosis and progression. *Ann Neurol* 2011;69:570-80.
27. Mollenhauer B, Locascio JJ, Schulz-Schaeffer W, Sixel-Doring F, Trenkwalder C, et al. alpha-Synuclein and tau concentrations in cerebrospinal fluid of patients presenting with parkinsonism: a cohort study. *Lancet Neurol* 2011;10:230-40.
28. Abdo WF, van de Warrenburg BP, Kremer HP, Bloem BR, Verbeek MM. CSF biomarker profiles do not differentiate between the cerebellar and parkinsonian phenotypes of multiple system atrophy. *Parkinsonism Relat Disord* 2007;13:480-2.
29. Herbert MK, Aerts MB, Beenes M, Norgren N, Esselink RA, et al. CSF neurofilament light chain but not FLT3 ligand discriminates parkinsonian disorders. *Front Neurol* 2015;6:91.
30. Abdo WF, De Jong D, Hendriks JC, Horstink MW, Kremer BP, et al. Cerebrospinal fluid analysis differentiates multiple system atrophy from Parkinson's disease. *Mov Disord* 2004;19:571-9.
31. Herbert MK, Eeftens JM, Aerts MB, Esselink RA, Bloem BR, et al. CSF levels of DJ-1 and tau distinguish MSA patients from PD patients and controls. *Parkinsonism Relat Disord* 2014;20:112-5.
32. Mollenhauer B, Cepek L, Bibl M, Wiltfang J, Schulz-Schaeffer WJ, et al. Tau protein, Abeta42 and S-100B protein in cerebrospinal fluid of patients with dementia with Lewy bodies. *Dement Geriatr Cogn Disord* 2005;19:164-70.
33. Mollenhauer B, Bibl M, Wiltfang J, Steinacker P, Ciesielczyk B, et al. Total tau protein, phosphorylated tau (181p) protein, beta-amyloid(1-42), and beta-amyloid(1-40) in cerebrospinal fluid of patients with dementia with Lewy bodies. *Clin Chem Lab Med* 2006;44:192-5.
34. Mulugeta E, Londos E, Hansson O, Ballard C, Skogseth R, et al. Cerebrospinal fluid levels of sAPPalpha and sAPPbeta in lewy body and Alzheimer's disease: clinical and neurochemical correlates. *Int J Alzheimers Dis* 2011;2011:495025.
35. Llorens F, Schmitz M, Varges D, Kruse N, Gotzmann N, et al. Cerebrospinal alpha-synuclein in alpha-synuclein aggregation disorders: tau/alpha-synuclein ratio as potential biomarker for dementia with Lewy bodies. *J Neurol* 2016;263:2271-7.
36. van Steenoven I, van der Flier WM, Scheltens P, Teunissen CE, Lemstra AW. Amyloid-beta peptides in cerebrospinal fluid of patients with dementia with Lewy bodies. *Alzheimers Res Ther* 2019;11:83.
37. Parnetti L, Tiraboschi P, Lanari A, Peducci M, Padiglioni C, et al. Cerebrospinal fluid biomarkers in Parkinson's disease with dementia and dementia with Lewy bodies. *Biol Psychiatry* 2008;64:850-5.
38. Kaerst L, Kuhlmann A, Wedekind D, Stoeck K, Lange P, et al. Using cerebrospinal fluid marker profiles in clinical diagnosis of dementia with Lewy bodies, Parkinson's disease, and Alzheimer's disease. *J Alzheimers Dis* 2014;38:63-73.
39. Gmitterova K, Gawinecka J, Llorens F, Varges D, Valkovic P, et al. Cerebrospinal fluid markers analysis in the differential diagnosis of dementia with Lewy bodies and Parkinson's disease dementia. *Eur Arch Psychiatry Clin Neurosci* 2018.
40. Gómez-Tortosa E, Gonzalo I, Fanjul S, Sainz MJ, Cantarero S, et al. Cerebrospinal fluid markers in dementia with lewy bodies compared with Alzheimer disease. *Arch Neurol* 2003;60:1218-22.
41. Boström F, Hansson O, Blennow K, Gerhardsson L, Lundh T, et al. Cerebrospinal fluid total tau is associated with shorter survival in dementia with Lewy bodies. *Dement Geriatr Cogn Disord* 2009;28:314-9.
42. Abdelnour C, van Steenoven I, Londos E, Blanc F, Auestad B, et al. Alzheimer's disease cerebrospinal fluid biomarkers predict cognitive decline in lewy body dementia. *Mov Disord* 2016;31:1203-8.
43. Mollenhauer B, Steinacker P, Bahn E, Bibl M, Brechlin P, et al. Serum heart-type fatty acid-binding protein and cerebrospinal fluid tau: marker candidates for dementia with lewy bodies. *Neurodegener Dis* 2007;4:366-75.
44. Slaets S, Le Bastard N, Theuns J, Sleegers K, Verstraeten A, et al. Amyloid pathology influences abeta1-42 cerebrospinal fluid levels in dementia with lewy bodies. *J Alzheimers Dis* 2013;35:137-46.
45. van Steenoven I, Aarsland D, Weintraub D, Londos E, Blanc F, et al. Cerebrospinal fluid Alzheimer's disease biomarkers across the spectrum of lewy body diseases: results from a large multicenter cohort. *J Alzheimers Dis* 2016;54:287-95.
46. Mollenhauer B, Bibl M, Trenkwalder C, Stiens G, Cepek L, et al. Follow-up investigations in cerebrospinal fluid of patients with dementia with Lewy bodies and Alzheimer's disease. *J Neural Transm (Vienna)* 2005;112:933-48.
47. Bousiges O, Cretin B, Lavaux T, Philippi N, Jung B, et al. Diagnostic value of cerebrospinal fluid biomarkers (Phospho-Tau181, total-Tau, Abeta42, and Abeta40) in prodromal stage of Alzheimer's disease and dementia with lewy bodies. *J Alzheimers Dis* 2016;51:1069-83.
48. Bousiges O, Bombois S, Schraen S, Wallon D, Quillard MM, et al. Cerebrospinal fluid Alzheimer biomarkers can be useful for discriminating dementia with Lewy bodies from Alzheimer's disease at the prodromal stage. *J Neurol Neurosurg Psychiatry* 2018;89:467-75.
49. Paraskevas GP, Bougea A, Constantinides VC, Bourbouli M, Petropoulou O, et al. In vivo prevalence of Alzheimer biomarkers in dementia with lewy bodies. *Dement Geriatr Cogn Disord* 2019;47:289-96.
50. Masliah E, Rockenstein E, Veinbergs I, Sagara Y, Mallory M, et al. beta-amyloid peptides enhance alpha-synuclein accumulation and neuronal deficits in a transgenic mouse model linking Alzheimer's disease and Parkinson's disease. *Proc Natl Acad Sci U S A* 2001;98:12245-50.
51. Sengupta U, Guerrero-Muñoz MJ, Castillo-Carranza DL, Lasagna-Reeves CA, Gerson JE, et al. Pathological interface between oligomeric alpha-synuclein and tau in synucleinopathies. *Biol Psychiatry* 2015;78:672-83.

52. Castillo-Carranza DL, Guerrero-Muñoz MJ, Sengupta U, Gerson JE, Kaye R. α -synuclein oligomers induce a unique toxic tau strain. *Biol Psychiatry* 2018;84:499-508.
53. Borroni B, Gardoni F, Parnetti L, Magno L, Malinverno M, et al. Pattern of Tau forms in CSF is altered in progressive supranuclear palsy. *Neurobiol Aging* 2009;30:34-40.
54. Borroni B, Malinverno M, Gardoni F, Alberici A, Parnetti L, et al. Tau forms in CSF as a reliable biomarker for progressive supranuclear palsy. *Neurology* 2008;71:1796-803.
55. Kuiperij HB, Verbeek MM. Diagnosis of progressive supranuclear palsy: can measurement of tau forms help? *Neurobiol Aging* 2012;33:204.e17-8.
56. Luk C, Compta Y, Magdalino N, Martí MJ, Hondhamuni G, et al. Development and assessment of sensitive immuno-PCR assays for the quantification of cerebrospinal fluid three- and four-repeat tau isoforms in tauopathies. *J Neurochem* 2012;123:396-405.
57. Barthélemy NR, Gabelle A, Hirtz C, Fenaille F, Sergeant N, et al. Differential mass spectrometry profiles of tau protein in the cerebrospinal fluid of patients with Alzheimer's disease, progressive supranuclear palsy, and dementia with lewy bodies. *J Alzheimers Dis* 2016;51:1033-43.
58. Cicognola C, Brinkmalm G, Wahlgren J, Portelius E, Gobom J, et al. Novel tau fragments in cerebrospinal fluid: relation to tangle pathology and cognitive decline in Alzheimer's disease. *Acta Neuropathol* 2019;137:279-96.
59. Arai H, Morikawa Y, Higuchi M, Matsui T, Clark CM, et al. Cerebrospinal fluid tau levels in neurodegenerative diseases with distinct tau-related pathology. *Biochem Biophys Res Commun* 1997;236:262-4.
60. Shelley BP, Hodges JR, Kipps CM, Xuereb JH, Bak TH. Is the pathology of corticobasal syndrome predictable in life? *Mov Disord* 2009;24:1593-9.
61. Ling H, O'Sullivan SS, Holton JL, Revesz T, Massey LA, et al. Does corticobasal degeneration exist? A clinicopathological re-evaluation. *Brain* 2010;133:2045-57.
62. Lee SE, Rabinovici GD, Mayo MC, Wilson SM, Seeley WW, et al. Clinicopathological correlations in corticobasal degeneration. *Ann Neurol* 2011;70:327-40.
63. Paik SR, Lee JH, Kim DH, Chang CS, Kim YS. Self-oligomerization of NACP, the precursor protein of the non-amyloid beta/A4 protein (A beta) component of Alzheimer's disease amyloid, observed in the presence of a C-terminal A beta fragment (residues 25-35). *FEBS Lett* 1998;421:73-6.
64. del Campo M, Mollenhauer B, Bertolotto A, Engelborghs S, Hampel H, et al. Recommendations to standardize preanalytical confounding factors in Alzheimer's and Parkinson's disease cerebrospinal fluid biomarkers: an update. *Biomark Med* 2012;6:419-30.

Review

Open Access



Assessment of cerebrospinal fluid α -synuclein as a potential biomarker in Parkinson's disease and synucleinopathies

Ioanna Chalatsa¹, Katerina Melachroinou¹, Evangelia Emmanouilidou^{1,2}, Kostas Vekrellis¹

¹Center for Basic Science, Biomedical Research Foundation Academy of Athens, Athens 11527, Greece.

²Laboratory of Biochemistry, Department of Chemistry, National and Kapodistrian University of Athens, Athens 11527, Greece.

Correspondence to: Dr. Kostas Vekrellis, Center for Basic Science, Biomedical Research Foundation Academy of Athens, 4 Soranou Ephessiou St., Athens 11527, Greece. E-mail: vekrellis@bioacademy.gr

How to cite this article: Chalatsa I, Melachroinou K, Emmanouilidou E, Vekrellis K. Assessment of cerebrospinal fluid α -synuclein as a potential biomarker in Parkinson's disease and synucleinopathies. *Neuroimmunol Neuroinflammation* 2020;7:132-40. <http://dx.doi.org/10.20517/2347-8659.2020.01>

Received: 2 Jan 2020 **First Decision:** 10 Mar 2020 **Revised:** 23 Mar 2020 **Accepted:** 7 Apr 2020 **Available online:** 16 May 2020

Science Editor: George P. Paraskevas **Copy Editor:** Jing-Wen Zhang **Production Editor:** Jing Yu

Abstract

The discovery of diagnostic and prognostic biomarkers for neurodegenerative diseases represents an unmet clinical challenge. For example, the diagnosis of Parkinson's disease (PD) relies mainly on the presence of clinical symptoms. Therefore, the identification and use of novel PD biomarkers would allow the application of disease-modifying treatments at the very early stages of neurodegeneration. The presynaptic protein, α -synuclein, has been genetically and biochemically linked with PD pathogenesis and has been considered as a potential biomarker for the diagnosis of PD and the related synucleinopathies. The vast majority of studies have assessed the measurement of α -synuclein, alone or in combination with other biomarkers in the cerebrospinal fluid (CSF), since it is the biofluid that most closely reflects the pathophysiology of the brain. The diagnostic value of the monomeric α -synuclein but also the oligomeric, the phosphorylated and the aggregated forms of the protein has been evaluated using a variety of immunoassays. The results have so far been reproducible but the assays used are still lacking the required diagnostic accuracy. Recent reports have shown that Protein misfolding cyclic amplification is a technique that has the potential to detect α -synuclein seeds in samples of CSF with high sensitivity and across different synucleinopathies. In an effort to increase the source of biomarker for PD and related synucleinopathies, α -synuclein has also been measured in neuronal exosomes, small vesicles of endosomal origin that are secreted from neurons into the CSF or the periphery. The potential diagnostic value of exosomes stems from the notion that exosomes carry a disease-specific repertoire of marker proteins. Therefore, the assessment of exosome-associated α -synuclein species may also open up new avenues for disease diagnosis in different synucleinopathies.



© The Author(s) 2020. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, sharing, adaptation, distribution and reproduction in any medium or format, for any purpose, even commercially, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.



Keywords: Cerebrospinal fluid, α -synuclein, Parkinson's disease, biomarker, exosomes, synucleinopathies

INTRODUCTION

The formation of large inclusions mostly containing protein aggregates is a common pathological hallmark in a wide spectrum of neurodegenerative disorders such as Alzheimer's Disease (AD) and Huntington Disease^[1]. Particularly three distinct neurological conditions, Parkinson's Disease (PD) including Parkinson's disease dementia, Dementia with Lewy Bodies (DLB) and Multiple System Atrophy (MSA), are characterized by the aberrant accumulation of the presynaptic protein α -synuclein. In PD and DLB, α -synuclein deposits are found either in the cytoplasm of neurons, where they are called lewy bodies (LB), or in the neuronal terminals, where they are called Lewy neurites^[2] whereas in MSA, α -synuclein deposition occurs in glial cells. α -Synuclein is also genetically linked with the development of PD since specific point mutations or multiplications (duplications, triplications) of the *SNCA* gene encoding for α -synuclein result in the familial forms of PD^[3-8]. The genetic association of α -synuclein with PD is further strengthened by all the genome-wide association studies performed so far which indicate a strong correlation of PD with variations in the *SNCA* gene^[9,10]. These biochemical and genetic linkages of α -synuclein with pathology as well as the observation that the protein is present in biological fluids or peripheral tissues led to the assumption that α -synuclein could serve as a potential candidate biomarker for PD diagnosis and also aid the differential diagnosis between the synucleinopathies^[11].

A plethora of studies have assessed the absolute quantification of α -synuclein levels as a marker of synucleinopathy with the ultimate aim to discriminate PD patients from healthy subjects or other unrelated neurological controls. In this regard, understanding the structural biology of α -synuclein is critical; the protein is highly modified at the post-translational level and has the ability to adopt different conformations depending on the surrounding milieu. From all the modifications that have been reported so far, phosphorylation is considered most closely related to PD pathology since almost 90% of α -synuclein in LB appears to be hyper-phosphorylated^[12]. In addition, the assembly into multiple-sized oligomers has been considered an early event in the pathological process of aggregate formation^[13]. As such, different forms of α -synuclein, i.e., monomeric, oligomeric and phosphorylated, have been targeted in order to increase diagnostic accuracy^[14]. The measurements have been performed in bodily fluids [cerebrospinal fluid (CSF), blood plasma or serum, saliva], isolated secreted vesicles (exosomes) and peripheral tissues (skin, olfactory or gut mucosa, salivary gland) using a variety of analytical approaches depending on the nature of the biological sample and the form of α -synuclein detected with each assay^[15].

In comparison with the other biological fluids, the assessment of α -synuclein in the CSF has provided the most consistent results in terms of analytical validation by different laboratories^[16]. CSF α -synuclein is mostly detected by means of immunoassays that use specific antibodies to target the different α -synuclein forms. Even though the absolute concentrations can vary from study to study, the results obtained so far are supported by several meta-analysis studies suggesting that CSF α -synuclein could serve as a potential marker of synucleinopathy^[17]. In this review, we aim to discuss the results from the assessment of α -synuclein in CSF and exosomes and explain the factors responsible for the variability among the different studies.

MEASUREMENT OF CSF α -SYNUCLEIN

Being primarily produced by the choroid plexus within the ventricles of the central nervous system, CSF is an established biological fluid to study neurodegenerative disorders since it is expected to mirror brain microenvironment. The quantification of total α -synuclein, as well as its oligomeric and phosphorylated forms, can be measured in CSF using different techniques, such as ELISA^[18], xMAP technology^[19],

mass spectrometry^[20], time-resolved fluorescence energy transfer^[21], electrochemiluminescence immunoassay^[22] and western blot^[23]. In addition, new biochemical assays that can detect α -synuclein aggregates have emerged, such as Protein-misfolding cyclic amplification and Real-time Quaking-induced conversion, by taking advantage the ability of α -synuclein to nucleate further aggregation^[24-26]. Using all these different methods, it is important to note that, even though there are variations in the absolute concentrations measured, the results produced for total (or monomeric) CSF α -synuclein agree on a reduction in α -synuclein levels in PD patients when compared with control subjects. When oligomeric or phosphorylated α -synuclein was assessed in the CSF, both forms were found to be increased in PD patients compared with the controls. However, it is important to note that the of ligands such as ThT may affect the actual structure of the α -synuclein species.

Even though the above findings are consistent, the diagnostic accuracy (sensitivity and specificity) remains unsatisfactory either for the detection of the monomeric or the modified forms of α -synuclein. Additionally, some studies report contradictory results; some have found similar CSF α -synuclein levels between PD patients and control subjects^[27-29], whereas others have reported increased CSF α -synuclein levels in samples from AD^[30,31], progressive supranuclear palsy or Creutzfeldt-Jacob patients compared with the control group^[32].

A number of factors could explain the observed variability in the results, as well as the differences reported in the absolute concentrations. First, the immunoassays used are based on divergent antibodies that recognize different fragments of the protein and with variable affinity. Second, the patient cohorts show great variability in terms of number, disease stage and clinical symptoms (affected mobility or dementia) present at the time of CSF collection. Third, the implementation of strict standardized guidelines concerning collection and storage protocols and allowed blood contamination have only recently started to be followed. Furthermore, common reference materials are still missing making the interpretation of results from assay to assay extremely difficult.

The quantification of CSF α -synuclein could aid the differential diagnosis in clinically overlapping neurodegenerative diseases, as suggested for PD, DLB and AD^[33-35]. However, it is unclear whether the levels of CSF α -synuclein could be correlated with the severity of disease, indicating for example a more rapid decline in motor performance or the appearance of dementia. Interestingly, recent reports have shown that simultaneous measurement of α -synuclein levels along with other proteins, such as tau, A β ₄₂ and Glucocerebrosidase 1 could be more effective in discriminating PD patients with synucleinopathies from healthy individuals or those with other neurodegenerative diseases^[30,36-38].

Other biological fluids could also serve as promising candidates for α -synuclein detection and subsequent PD diagnosis. The majority of reports studying plasma α -synuclein have exhibited increased levels in PD patients^[39-46] relative to control subjects, whereas other studies have reported similar^[27,47,48] or decreased plasma α -synuclein levels^[49] between PD patients and healthy participants. Interestingly, it was found that plasma levels of phosphorylated α -synuclein were higher in the PD samples than the controls^[44,50,51]. The results obtained from plasma have been controversial, mainly due to the fact that red blood cells are a major source of α -synuclein and the rest erythrocytes that remain in plasma^[52] can be subjected to hemolysis markedly affecting α -synuclein values^[47].

As erythrocytes are the major source of peripheral α -synuclein, a recent report has proposed erythrocytic α -synuclein as a potential PD biomarker, as it was found that the total and aggregated α -synuclein levels were significantly higher in the membrane fraction of PD patients compared to healthy controls^[53]. Saliva α -synuclein has also been considered as a prospective biomarker, as α -synuclein pathology has been found in submandibular salivary glands^[54,55] and saliva α -synuclein could be easily accessible and poorly affected

by blood contamination^[56-59]. Some studies have reported that total α -synuclein levels were reduced in the saliva of PD patients compared with control subjects, whereas oligomeric α -synuclein appeared to be elevated in the saliva of PD patients^[56,57].

MEASUREMENT OF α -SYNUCLEIN FROM NEURONAL EXOSOMES AS A POTENTIAL BIOMARKER

α -synuclein was considered to be localized mostly in the cytoplasm of neuronal cells, until several studies demonstrated its presence in human CSF, human plasma and in the conditioned medium of various cell lines^[23,60]. Many studies have shown that α -synuclein is physiologically secreted in the extracellular space, but the mechanism of α -synuclein release is still unclear. Evidence from recent studies has also suggested that extracellular α -synuclein can confer to the progression of PD^[1-3] and it has been proposed that α -synuclein secretion, either in a monomeric or oligomeric state, induces α -synuclein propagation via cell-to-cell transfer^[61]. Thus, elucidating the mechanism by which α -synuclein is secreted in the extracellular space is of great importance in understanding cellular pathways that may cause PD.

Release of α -synuclein via extracellular vesicles termed exosomes has been demonstrated by our group and others^[62,63]. Exosomes are extracellular vesicles of ~50 to 200 nm diameter and can mediate proximal and distal cellular communication through the transfer of biological molecules between cells. They originate from the inward budding of multi-vesicular bodies (MVBs) and are released to the extracellular space upon fusion of MVBs with the plasma membrane in an exocytic manner. Exosomes are released from numerous cell types including neurons and glia^[64] and in several studies have been observed to be associated with pathologic proteins including α -synuclein^[63,65]. Based on the current knowledge, exosomes are functionally active entities, with a highly versatile role, ranging from intercellular communication by delivering specific protein, lipid or RNA cargo, and removal of obsolete or misfolded proteins, as a means of cell detoxification, to deleterious shuttles that impair cell homeostasis^[66].

Some well-characterized functions of exosomes are protein secretion and intracellular uptake, immune response regulation and toxicity induction^[67]. Interestingly, Danzer *et al.*^[65] demonstrated that exosome associated α -synuclein is more potent in transmitting aggregation pathology between neurons than free-secreted α -synuclein. One study has shown that patients with PD have higher α -synuclein levels in plasma exosomes compared to healthy controls^[68], while Stuenkel *et al.*^[69] found decreased neural exosome α -synuclein levels in PD patients, consistent with the total α -synuclein levels in CSF. In addition, the quantification of CSF exosomal α -synuclein exhibited distinct differences between patients with PD and DLB. Moreover, exosomal α -synuclein levels correlated with the severity of cognitive impairment in cross-sectional samples from patients with DLB. In the same study, Stuenkel *et al.*^[69] showed that exosomes from PD and DLB patients contain pathogenic α -synuclein species which serve as seeds to induce the oligomerization of soluble α -synuclein in recipient cells. Shi *et al.*^[70] have shown that CNS-derived exosomes can efflux into blood. Importantly, they found a substantially augmented α -synuclein concentration in the plasma-isolated exosomes from PD patients compared to healthy control subjects, despite the fact that no differences were detected in plasma total α -synuclein levels. Additionally, they report a significant increase of plasma exosomal α -synuclein/total α -synuclein ratio in PD patients, negatively correlated with the disease severity, further supporting the importance of the disease-related exosomal cargos as PD biomarkers with high sensitivity and specificity. The authors concluded that plasma, CNS-derived exosomal α -synuclein can serve as a PD biomarker with high sensitivity and specificity^[70]. The same group has also shown that CNS-derived exosomal tau in plasma is significantly higher in PD patients than in controls and is correlated with CSF total tau and phosphorylated tau^[71]. Furthermore, distinct circulating exosome entities have been identified in the serum of patients with PD^[72]. A recent study demonstrated that the levels of DJ-1 and α -synuclein in plasma CSF-derived exosomes and the

ratio of plasma CSF-derived exosomal DJ-1 to total DJ-1 were significantly higher in patients with PD, compared with controls^[4]. Several factors have been shown to affect the release of α -synuclein through exosomes such as the activity of Glucocerebrosidase enzyme (GCase), ion homeostasis, such as Zn^{2+} , Ca^{2+} and Mn^{2+} , as well as neuronal activity and neurotransmitter release. The heterozygous mutations in the *GBA1* gene are considered as an important risk factor for PD. In this regard, it has been demonstrated that GCase overexpression leads to a decrease of exosome secretion while chronic pharmacological inhibition of GCase activity *in vivo* profoundly increased exosomes levels, as well as exosome-associated α -synuclein oligomers^[73]. In addition, decreased GCase activity has been demonstrated in brain samples with increased α -synuclein levels and in CSF from sporadic PD patients^[74]. More recently, a study by Cerri *et al.*^[75], showed that exosomes from PD patients contain a greater amount of α -synuclein compared to healthy subjects whereas no differences were found in plasma total α -synuclein levels. Importantly, the authors showed a significant inverse correlation between GCase activity and this ratio in PD patients.

Notably, exosomes being a snapshot of the intracellular milieu, comprise a great source of bioactive molecules, including various RNA species. In a study conducted by Gui and co-workers, where exosomal miRNAs were isolated from the CSF of PD patients, 16 and 11 exosomal miRNAs were found upregulated and down-regulated, respectively, in PD patients compared to controls^[76]. Validated hits were found to be miR-1 and miR-19b-3p, significantly reduced in PD CSF exosomes, in contrast to miR-153, miR-409-3p, miR-10a-5p, and let-7g-3p which were found to be increased^[76]. This evidence highlights the potential diagnostic value of CSF exosomal RNA in the assessment of PD.

Although the role of exosome-associated α -synuclein as a potential biomarker remains relatively controversial, there are specific parameters that should be taken into account. Firstly, with regards to the source of exosomes, including plasma, saliva, CSF, there are certain protocols for their acquisition, followed by even more meticulous procedures for exosome isolation and purification. Differences during the aforementioned protocols may account for deviations between groups. In the same context, total exosome isolation may mask differences that could be found in exosomes of specific cellular origin. For example, Tomlinson and colleagues^[72], following an unbiased proteomic approach, did not find any significant increase of α -synuclein in total exosomes isolated by ultracentrifugation from the serum of PD patients. On the contrary, Shi *et al.*^[70] showed a significant α -synuclein enrichment in neuron-specific exosomes, isolated from PD patients' plasma by immunoprecipitation of the neuronal adhesion molecule L1CAM. More importantly, the notion that α -synuclein exists and exerts its detrimental effects in different strains, leading to different aggregates that cause as many distinct synucleinopathies, i.e., PD, DLB, MSA has been cemented^[77,78]. Given the fact that exosomal cargo mirrors the state of the cell from which it originates, exosome-associated α -synuclein may reflect the dynamic nature of α -synuclein species. To this end, it is of a pivotal importance to develop techniques that could allow detection and quantification of all the different α -synuclein conformers. Overall, although in its infancy, the study of exosome-associated α -synuclein as a potential biomarker is quite promising, yet, it requires more combinatorial approaches.

CONCLUSION

Over the last 10 years there has been considerable amount of research effort placed in the evaluation of neuronal α -synuclein as a diagnostic or, even, a prognostic biomarker for PD and related synucleinopathies. The majority of studies have indicated that CSF α -synuclein could be useful for the diagnosis of synucleinopathy that could also aid the distinguishment of PD patients from patients with other neurodegenerative conditions. However, its utility as a biomarker is hampered by the lack of a universally validated assay of high diagnostic accuracy. To this end, following the strict recently established standard operating protocols for CSF collection and storage and correlating the measurement of monomeric α -synuclein with the oligomeric or phosphorylated forms would greatly improve the diagnostic value of the assessment of α -synuclein in CSF. To further ameliorate the specificity of α -synuclein measurement, recent

experimental approaches involve the assessment of α -synuclein in neural exosomes. The discoveries of pathogenic misfolded proteins such as α -synuclein in them has generated intensive research into their use as biomarkers considering that they carry proteins with disease-specific fingerprints reflecting the presence and staging of the disease. However, in order to further verify this potential we need to have a very good understanding of the actual mechanisms behind their biogenesis and release. Importantly, we need to have those tools in place that will assist us in the identification of the α -synuclein species responsible for disease generation and pathology progression. The fact that exosomal cargo mirrors the state of the cell from which it originates^[79] unravels the promising role of the plasma/CSF -derived exosomes as potential biomarkers. Proteomic profiling of exosomal proteins in PD patients with different disease stages and healthy subjects may also aid the identification of specific protein changes that occur in response to pathology progression. Finally, modulating exosome biogenesis and release may have a promising prospect in PD therapy.

DECLARATIONS

Authors' contributions

Contributed to the writing of the manuscript and in its revision: Chalatsa I, Melachroinou K, Emmanouilidou E, Vekrellis K

Availability of data and materials

Not applicable.

Financial support and sponsorship

None.

Conflicts of interest

All authors declared that there are no conflicts of interest.

Ethical approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Copyright

© The Author(s) 2020.

REFERENCES

1. Uversky VN. Neuropathology, biochemistry, and biophysics of alpha-synuclein aggregation. *J Neurochem* 2007;103:17-37.
2. Spillantini MG, Schmidt ML, Lee VM, Trojanowski JQ, Jakes R, et al. Alpha-synuclein in Lewy bodies. *Nature* 1997;388:839-40.
3. Polymeropoulos MH, Lavedan C, Leroy E, Ide SE, Dehejia A, et al. Mutation in the alpha-synuclein gene identified in families with Parkinson's disease. *Science* 1997;276:2045-7.
4. Zarranz JJ, Alegre J, Gómez-Esteban JC, Lezcano E, Ros R, et al. The new mutation, E46K, of alpha-synuclein causes Parkinson and Lewy body dementia. *Ann Neurol* 2004;55:164-73.
5. Proukakis C, Houlden H, Schapira AH. Somatic alpha-synuclein mutations in Parkinson's disease: hypothesis and preliminary data. *Mov Disord* 2013;28:705-12.
6. Lesage S, Le Ber I, Condroyer C, Broussolle E, Gabelle A, et al. C9orf72 repeat expansions are a rare genetic cause of parkinsonism. *Brain* 2013;136:385-91.
7. Chartier-Harlin MC, Kachergus J, Roumier C, Mouroux V, Douay X, et al. Alpha-synuclein locus duplication as a cause of familial Parkinson's disease. *Lancet* 2004;364:1167-9.
8. Singleton AB, Farrer MJ, Bonifati V. The genetics of Parkinson's disease: progress and therapeutic implications. *Mov Disord* 2013;28:14-23.
9. Satake W, Nakabayashi Y, Mizuta I, Hirota Y, Ito C, et al. Genome-wide association study identifies common variants at four loci as genetic risk factors for Parkinson's disease. *Nat Genet* 2009;41:1303-7.

10. Simón-Sánchez J, Schulte C, Bras JM, Sharma M, Gibbs JR, et al. Genome-wide association study reveals genetic risk underlying Parkinson's disease. *Nat Genet* 2009;41:1308-12.
11. Emamzadeh FN, Surguchov A. Parkinson's disease: biomarkers, treatment, and risk factors. *Front Neurosci* 2018;12:612.
12. Fujiwara H, Hasegawa M, Dohmae N, Kawashima A, Masliah E, et al. alpha-Synuclein is phosphorylated in synucleinopathy lesions. *Nat Cell Biol* 2002;4:160-4.
13. Paleologou KE, Kragh CL, Mann DM, Salem SA, Al-Shami R, et al. Detection of elevated levels of soluble alpha-synuclein oligomers in post-mortem brain extracts from patients with dementia with Lewy bodies. *Brain* 2009;132:1093-101.
14. Surguchov A. Parkinson's disease: assay of phosphorylated alpha-synuclein in skin biopsy for early diagnosis and association with melanoma. *Brain Sci* 2016;6.
15. Mollenhauer B, Bowman FD, Drake D, Duong J, Blennow K, et al. Antibody-based methods for the measurement of α -synuclein concentration in human cerebrospinal fluid - method comparison and round robin study. *J Neurochem* 2019;149:126-38.
16. Magdalinou N, Lees AJ, Zetterberg H. Cerebrospinal fluid biomarkers in parkinsonian conditions: an update and future directions. *J Neurol Neurosurg Psychiatry* 2014;85:1065-75.
17. Eusebi P, Hansson O, Paciotti S, Orso M, Chiasserini D, et al. Cerebrospinal fluid biomarkers for the diagnosis and prognosis of Parkinson's disease: protocol for a systematic review and individual participant data meta-analysis. *BMJ Open* 2017;7:e018177.
18. Mollenhauer B, Caspell-Garcia CJ, Coffey CS, Taylor P, Singleton A, et al. Longitudinal analyses of cerebrospinal fluid α -Synuclein in prodromal and early Parkinson's disease. *Mov Disord* 2019;34:1354-64.
19. Hong Z, Shi M, Chung KA, Quinn JF, Peskind ER, et al. DJ-1 and alpha-synuclein in human cerebrospinal fluid as biomarkers of Parkinson's disease. *Brain* 2010;133:713-26.
20. Mollenhauer B, Cullen V, Kahn I, Krastins B, Outeiro TF, et al. Direct quantification of CSF alpha-synuclein by ELISA and first cross-sectional study in patients with neurodegeneration. *Exp Neurol* 2008;213:315-25.
21. van Dijk KD, Bidinosti M, Weiss A, Raijmakers P, Berendse HW, et al. Reduced α -synuclein levels in cerebrospinal fluid in Parkinson's disease are unrelated to clinical and imaging measures of disease severity. *Eur J Neurol* 2014;21:388-94.
22. Kruse N, Mollenhauer B. Quantification of alpha-synuclein in biological fluids by electrochemiluminescence-based detection. *Methods Mol Biol* 2019;1948:59-68.
23. Borghi R, Marchese R, Negro A, Marinelli L, Forloni G, et al. Full length alpha-synuclein is present in cerebrospinal fluid from Parkinson's disease and normal subjects. *Neurosci Lett* 2000;287:65-7.
24. Saijo E, Groveman BR, Kraus A, Metrick M, Orrù CD, et al. Ultrasensitive RT-QuIC seed amplification assays for disease-associated tau, α -synuclein, and prion aggregates. *Methods Mol Biol* 2019;1873:19-37.
25. Shahnawaz M, Mukherjee A, Pritzkow S, Mendez N, Rabadia P, et al. Discriminating alpha-synuclein strains in Parkinson's disease and multiple system atrophy. *Nature* 2020;578:273-7.
26. Shahnawaz M, Tokuda T, Waragai M, Mendez N, Ishii R, et al. Development of a biochemical diagnosis of Parkinson disease by detection of α -synuclein misfolded aggregates in cerebrospinal fluid. *JAMA Neurol* 2017;74:163-72.
27. Park MJ, Cheon SM, Bae HR, Kim SH, Kim JW. Elevated levels of α -synuclein oligomer in the cerebrospinal fluid of drug-naïve patients with Parkinson's disease. *J Clin Neurol* 2011;7:215-22.
28. Ohrfelt A, Grognet P, Andreasen N, Wallin A, Vanmechelen E, et al. Cerebrospinal fluid alpha-synuclein in neurodegenerative disorders-a marker of synapse loss. *Neurosci Lett* 2009;450:332-5.
29. Aerts MB, Esselink RA, Abdo WF, Bloem BR, Verbeek MM. CSF α -synuclein does not differentiate between parkinsonian disorders. *Neurobiol Aging* 2012;33:430.e1-3.
30. García-Ayllón MS, Monge-Argilés JA, Monge-García V, Navarrete F, Cortés-Gómez MA, et al. Measurement of CSF α -synuclein improves early differential diagnosis of mild cognitive impairment due to Alzheimer's disease. *J Neurochem* 2019;150:218-30.
31. Twohig D, Rodriguez-Vieitez E, Sando SB, Berge G, Lauridsen C, et al. The relevance of cerebrospinal fluid α -synuclein levels to sporadic and familial Alzheimer's disease. *Acta Neuropathol Commun* 2018;6:130.
32. Oeckl P, Metzger F, Nagl M, von Arnim CA, Halbgebauer S, et al. Alpha-, beta-, and gamma-synuclein quantification in cerebrospinal fluid by multiple reaction monitoring reveals increased concentrations in Alzheimer's and Creutzfeldt-Jakob disease but no alteration in synucleinopathies. *Mol Cell Proteomics* 2016;15:3126-38.
33. Katayama T, Sawada J, Kikuchi-Takeguchi S, Kano K, Takahashi K, et al. Cerebrospinal fluid levels of alpha-synuclein, amyloid β , tau, phosphorylated tau, and neuron-specific enolase in patients with Parkinson's disease, dementia with Lewy bodies or other neurological disorders: their relationships with cognition and nuclear medicine imaging findings. *Neurosci Lett* 2019;715:134564.
34. Bougea A, Stefanis L, Paraskevas GP, Emmanouilidou E, Efthymiopoulou E, et al. Neuropsychiatric symptoms and α -Synuclein profile of patients with Parkinson's disease dementia, dementia with Lewy bodies and Alzheimer's disease. *J Neurol* 2018;265:2295-301.
35. Bongianini M, Ladogana A, Capaldi S, Klotz S, Baiardi S, et al. α -Synuclein RT-QuIC assay in cerebrospinal fluid of patients with dementia with Lewy bodies. *Ann Clin Transl Neurol* 2019;6:2120-6.
36. Lerche S, Wurster I, Roeben B, Zimmermann M, Riebenbauer B, et al. Parkinson's disease: glucocerebrosidase 1 mutation severity is associated with CSF alpha-synuclein profiles. *Mov Disord* 2020;35:495-9.
37. Viodé A, Epelbaum S, Benyounes I, Verny M, Dubois B, et al. Simultaneous quantification of tau and α -synuclein in cerebrospinal fluid by high-resolution mass spectrometry for differentiation of Lewy Body Dementia from Alzheimer's disease and controls. *Analyst* 2019;144:6342-51.
38. Shi M, Tang L, Toledo JB, Ghingina C, Wang H, et al. Cerebrospinal fluid α -synuclein contributes to the differential diagnosis of Alzheimer's disease. *Alzheimers Dement* 2018;14:1052-62.

39. Chen XQ, Niu JP, Peng RQ, Song YH, Xu N, et al. The early diagnosis of Parkinson's disease through combined biomarkers. *Acta Neurol Scand* 2019;140:268-73.
40. Duran R, Barrero FJ, Morales B, Luna JD, Ramirez M, et al. Plasma alpha-synuclein in patients with Parkinson's disease with and without treatment. *Mov Disord* 2010;25:489-93.
41. Lee PH, Lee G, Park HJ, Bang OY, Joo IS, et al. The plasma alpha-synuclein levels in patients with Parkinson's disease and multiple system atrophy. *J Neural Transm* 2006;113:1435-9.
42. Chang CW, Yang SY, Yang CC, Chang CW, Wu YR. Plasma and serum alpha-synuclein as a biomarker of diagnosis in patients with Parkinson's disease. *Front Neurol* 2019;10:1388.
43. Ng ASL, Tan YJ, Lu Z, Ng EYL, Ng SYE, et al. Plasma alpha-synuclein detected by single molecule array is increased in PD. *Ann Clin Transl Neurol* 2019;6:615-9.
44. Lin CH, Liu HC, Yang SY, Yang KC, Wu CC, et al. Plasma pS129-alpha-synuclein is a surrogate biofluid marker of motor severity and progression in Parkinson's disease. *J Clin Med* 2019;8.
45. Lin CH, Yang SY, Horng HE, Yang CC, Chieh JJ, et al. Plasma biomarkers differentiate Parkinson's disease from atypical parkinsonism syndromes. *Front Aging Neurosci* 2018;10:123.
46. Lin CH, Yang SY, Horng HE, Yang CC, Chieh JJ, et al. Plasma alpha-synuclein predicts cognitive decline in Parkinson's disease. *J Neurol Neurosurg Psychiatry* 2017;88:818-24.
47. Shi M, Zabetian CP, Hancock AM, Ginchina C, Hong Z, et al. Significance and confounders of peripheral DJ-1 and alpha-synuclein in Parkinson's disease. *Neurosci Lett* 2010;480:78-82.
48. Mata IF, Shi M, Agarwal P, Chung KA, Edwards KL, et al. SNCA variant associated with Parkinson disease and plasma alpha-synuclein level. *Arch Neurol* 2010;67:1350-6.
49. Li QX, Mok SS, Laughton KM, McLean CA, Cappai R, et al. Plasma alpha-synuclein is decreased in subjects with Parkinson's disease. *Exp Neurol* 2007;204:583-8.
50. Cariulo C, Martufi P, Verani M, Azzollini L, Bruni G, et al. Phospho-S129 alpha-synuclein is present in human plasma but not in cerebrospinal fluid as determined by an ultrasensitive immunoassay. *Front Neurosci* 2019;13:889.
51. Foulds PG, Diggle P, Mitchell JD, Parker A, Hasegawa M, et al. A longitudinal study on alpha-synuclein in blood plasma as a biomarker for Parkinson's disease. *Sci Rep* 2013;3:2540.
52. Barbour R, Kling K, Anderson JP, Banducci K, Cole T, et al. Red blood cells are the major source of alpha-synuclein in blood. *Neurodegener Dis* 2008;5:55-9.
53. Tian C, Liu G, Gao L, Soltys D, Pan C, et al. Erythrocytic alpha-Synuclein as a potential biomarker for Parkinson's disease. *Transl Neurodegener* 2019;8:15.
54. Adler CH, Serrano GE, Zhang N, Hinni ML, Lott DG, et al. Feasibility of repeat and bilateral submandibular gland needle biopsies in Parkinson's disease. *Parkinsonism Relat Disord* 2019;68:69-72.
55. Del Tredici K, Hawkes CH, Ghebremedhin E, Braak H. Lewy pathology in the submandibular gland of individuals with incidental Lewy body disease and sporadic Parkinson's disease. *Acta Neuropathol* 2010;119:703-13.
56. Kang W, Chen W, Yang Q, Zhang L, Zhang L, et al. Salivary total alpha-synuclein, oligomeric alpha-synuclein and SNCA variants in Parkinson's disease patients. *Sci Rep* 2016;6:28143.
57. Vivacqua G, Suppa A, Mancinelli R, Belvisi D, Fabbrini A, et al. Salivary alpha-synuclein in the diagnosis of Parkinson's disease and Progressive Supranuclear Palsy. *Park Relat Disord* 2019;63:143-8.
58. Al-Nimer MSM, Mshatat SF, Abdulla HI. Saliva alpha-synuclein and a high extinction coefficient protein: a novel approach in assessment biomarkers of Parkinson's disease. *N Am J Med Sci* 2014;6:633-7.
59. Devic I, Hwang H, Edgar JS, Izutsu K, Presland R, et al. Salivary alpha-synuclein and DJ-1: potential biomarkers for Parkinson's disease. *Brain* 2011;134:178.
60. Lee HJ, Patel S, Lee SJ. Intravesicular localization and exocytosis of alpha-synuclein and its aggregates. *J Neurosci* 2005;25:6016-24.
61. Recasens A, Dehay B. Alpha-synuclein spreading in Parkinson's disease. *Front Neuroanat* 2014;8:159.
62. Alvarez-Erviti L, Seow Y, Schapira AH, Gardiner C, Sargent IL, et al. Lysosomal dysfunction increases exosome-mediated alpha-synuclein release and transmission. *Neurobiol Dis* 2011;42:360-7.
63. Emmanouilidou E, Melachroinou K, Roumeliotis T, Garbis SD, Ntzouni M, et al. Cell-produced alpha-synuclein is secreted in a calcium-dependent manner by exosomes and impacts neuronal survival. *J Neurosci* 2010;30:6838-51.
64. Von Bartheld CS, Altick AL. Multivesicular bodies in neurons: distribution, protein content, and trafficking functions. *Prog Neurobiol* 2011;93:313-40.
65. Danzer KM, Kranich LR, Ruf WP, Cagsal-Getkin O, Winslow AR, et al. Exosomal cell-to-cell transmission of alpha synuclein oligomers. *Mol Neurodegener* 2012;7:42.
66. Bellingham SA, Guo BB, Coleman BM, Hill AF. Exosomes: vehicles for the transfer of toxic proteins associated with neurodegenerative diseases. *Front Physiol* 2012;3:124.
67. Chang C, Lang H, Geng N, Wang J, Li N, et al. Exosomes of BV-2 cells induced by alpha-synuclein: important mediator of neurodegeneration in PD. *Neurosci Lett* 2013;548:190-5.
68. Zhao ZH, Chen ZT, Zhou RL, Zhang X, Ye QY, et al. Increased DJ-1 and α -synuclein in plasma neural-derived exosomes as potential markers for Parkinson's disease. *Front Aging Neurosci* 2018;10:438.
69. Stuendl A, Kunadt M, Kruse N, Bartels C, Moebius W, et al. Induction of α -synuclein aggregate formation by CSF exosomes from patients with Parkinson's disease and dementia with Lewy bodies. *Brain* 2016;139:481-94.

70. Shi M, Liu C, Cook TJ, Bullock KM, Zhao Y, et al. Plasma exosomal α -synuclein is likely CNS-derived and increased in Parkinson's disease. *Acta Neuropathol* 2014;128:639-50.
71. Shi M, Kovac A, Korff A, Cook TJ, Ginhina C, et al. CNS tau efflux via exosomes is likely increased in Parkinson's disease but not in Alzheimer's disease. *Alzheimers Dement* 2016;12:1125-31.
72. Tomlinson PR, Zheng Y, Fischer R, Heidasch R, Gardiner C, et al. Identification of distinct circulating exosomes in Parkinson's disease. *Ann Clin Transl Neurol* 2015;2:353-61.
73. Papadopoulos VE, Nikolopoulou G, Antoniadou I, Karachaliou A, Arianoglou G, et al. Modulation of β -glucocerebrosidase increases α -synuclein secretion and exosome release in mouse models of Parkinson's disease. *Hum Mol Genet* 2018;27:1696-710.
74. Parnetti L, Paciotti S, Eusebi P, Dardis A, Zampieri S, et al. Cerebrospinal fluid β -glucocerebrosidase activity is reduced in parkinson's disease patients. *Mov Disord* 2017;32:1423-31.
75. Cerri S, Ghezzi C, Sampieri M, Siani F, Avenali M, et al. The exosomal/total α -synuclein ratio in plasma is associated with glucocerebrosidase activity and correlates with measures of disease severity in PD patients. *Front Cell Neurosci* 2018;12:125.
76. Gui YX, Liu H, Zhang LS, Lv W, Hu XY. Altered microRNA profiles in cerebrospinal fluid exosome in Parkinson disease and Alzheimer disease. *Oncotarget* 2015;6:37043-53.
77. Peelaerts W, Bousset L, Van der Perren A, Moskalyuk A, Pulizzi R, et al. alpha-Synuclein strains cause distinct synucleinopathies after local and systemic administration. *Nature* 2015;522:340-4.
78. Lee SJ, Masliah E. Neurodegeneration: aggregates feel the strain. *Nature* 2015;522:296-7.
79. De Toro J, Herschlik L, Waldner C, Mongini C. Emerging roles of exosomes in normal and pathological conditions: new insights for diagnosis and therapeutic applications. *Front Immunol.* 2015;6:203.

Review

Open Access



COVID-19 neurotropism and implications for therapy

Mahasweta Das^{1,4}, Courtney Penn^{1,4}, Taylor Martinez^{1,4}, Karthick Mayilsamy^{1,4}, Andrew McGill^{1,2,4}, Alison Wiling³, Shyam S. Mohapatra^{2,4}, Subhra Mohapatra^{1,4}

¹Department of Molecular Medicine, University of South Florida, Morsani College of Medicine, Tampa, FL 33612, USA.

²Department of Internal Medicine, University of South Florida, Morsani College of Medicine, Tampa, FL 33612, USA.

³Department of Neurosurgery and Brain Repair, Center of Excellence for aging and Brain Repair, University of South Florida, Morsani College of Medicine, Tampa, FL 33612, USA.

⁴James A Haley Veterans Hospital, Tampa, FL 33612, USA.

Correspondence to: Dr. Subhra Mohapatra, Department of Molecular Medicine, University of South Florida, Morsani College of Medicine, Tampa, FL 33612, USA. E-mail: smohapa2@usf.edu; and Dr. Shyam S. Mohapatra, Department of Internal Medicine, University of South Florida, Morsani College of Medicine, Tampa, FL 33612, USA. E-mail: smohapat@usf.edu

How to cite this article: Das M, Penn C, Martinez T, Mayilsamy K, McGill A, Wiling A, Mohapatra SS, Mohapatra S. COVID-19 neurotropism and implications for therapy. *Neuroimmunol Neuroinflammation* 2020;7: 141-9.
<http://dx.doi.org/10.20517/2347-8659.2020.36>

Received: 28 Apr 2020 **First Decision:** 25 May 2020 **Revised:** 25 May 2020 **Accepted:** 1 Jun 2020 **Available online:** 3 Jun 2020

Science Editor: Athanassios P. Kyritsis **Copy Editor:** Cai-Hong Wang **Production Editor:** Jing Yu

Abstract

The mechanism underlying the pathogenesis of severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) infection in humans is poorly understood, although the cellular receptors which facilitate the virus fusion have been identified. Although the major symptoms of the infection have been identified as acute respiratory distress, pneumonia, and fever, recently, symptoms involving nervous system dysfunctions, including encephalopathy and stroke, have been detected. Herein, we comprehensively review the evidence that SARS-CoV-2 infection involves a neurotropic mechanism including a nose-brain-lung axis suggesting implications in therapy development.

Keywords: SARS-CoV-2, nose-brain-lung axis, neurotropism

INTRODUCTION

The current global pandemic outbreak of novel severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is known for its viral tropism to the lungs, which in severe cases can lead to the fatal respiratory failure of patients who have contracted the disease. SARS-CoV-2 and SARS-CoV viruses have a 70%-80% homology and both enter the body through the same receptor, angiotensin converting enzyme 2 (ACE2)^[1].



© The Author(s) 2020. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, sharing, adaptation, distribution and reproduction in any medium or format, for any purpose, even commercially, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.



In addition to their genetic homology, the pathology that the two viruses exhibit in the clinic are highly similar^[2]. In a recent report from Wuhan, China, Mao and colleagues showed that in addition to respiratory symptoms including pneumonia and acute respiratory distress syndrome (ARDS), out of 214 hospitalized COVID-19 patients 78 (36.4%) showed neurological manifestations such as cerebrovascular diseases (5.7%), impaired consciousness (14.8%), and skeletal muscle injury (19.3%)^[3]. Six percent of COVID-19 patients have been reported to show symptoms of stroke and 15% were reported to show encephalopathy^[4]. A number of symptoms such as dizziness, headache, loss of taste and smell, impaired consciousness, seizures, and nerve pain suggest a neurological connection to this viral infection.

Previously, SARS-CoV was observed in the cerebrospinal fluid of a SARS patient showing ARDS^[5]. Neuropathy in a COVID-19 patient has been reported in which the patient had hyposmia and altered taste sensation^[6]. Establishing the link between prior coronaviruses and their neurotropism appears to be important because there is now mounting evidence for SARS-CoV-2 neuronal abnormalities emerging as a significant symptom associated with the disease^[7]. COVID-19 patients can show neurological manifestation under three situations: due to neurotropism of the virus, post-infective neurological complications or aggravated symptoms in patients with neurological co-morbidities like dementia^[8]. Moreover, the neurological complications and symptoms can be central, including headache, dizziness, altered sensorium, stroke, ataxia, encephalitis, and seizures, or peripheral, like loss of smell or taste sensation or skeletal muscle injury^[8]. Herein, we review a profound neurological basis of SARS-CoV-2 infection-induced global pandemic and how the viral infection modulates the breathing of the lungs leading to fatal ARDS, to which about 50%-80% of the severely ill patients succumb.

NEUROTROPISM OF β -CORONAVIRUSES

The family of β -coronaviruses, to which SARS and SARS-CoV-2 belong, have been extensively studied. Several of the β -coronaviruses have been discovered within the brain (especially the brain stem), including SARS-CoV, hCoV-229E, hCoV-OC43, mouse hepatitis E, and porcine hemagglutinating encephalomyelitis (HEV) coronavirus^[5,9]. Middle Eastern respiratory syndrome (MERS) virus, another notable zoonotic viral outbreak in 2008-2009, that utilizes a different host receptor, DPP4, has also been discovered in the brain stem of experimental animals even in the absence of lung infection^[10], demonstrating a strong propensity for the family of β -coronaviridae to infect the brain. While ACE2 is expressed at low levels within the brain, it has been hypothesized that the “promiscuous” spike protein of β -coronaviruses may also have an unknown receptor in the brain and peripheral nerves to which the neurotropism is owed^[10].

SARS-CoV and MERS-CoV are considered important coronaviruses with the potential of nervous system damage. SARS-CoV was found to cause neurological conditions like demyelination of nerve fibers, ischemic changes of neurons, and diseases such as encephalitis, polyneuropathy, and aortic ischemic stroke^[11]. Similarly, 1 in 5 MERS-CoV patients were reported with neurological symptoms such as paralysis, ischemic stroke, loss of consciousness and Guillain Barré syndrome^[12]. The presence of SARS-CoV has also been demonstrated in mice brain experimentally infected with SARS-CoV^[13]. Another coronavirus, the mouse hepatitis virus type 3, can disrupt the BBB by down-regulating interferon β which would allow virus and infected immune cells in^[14].

NEURAL CONTROL OF BREATHING

The major life-threatening symptoms of SARS-CoV-2 infection are respiratory failure and pneumonia. While pneumonia is largely an inflammatory condition of the lungs, breathing is extensively regulated by the brain. Breathing is regulated by numerous internal and external stimuli, which are integrated in the brain stem to produce the muscle movements necessary for inspiration (or inhalation), and expiration (or exhalation) [Figure 1A]. The rate and phases of the breathing pattern arise from respiratory centers




Figure 1. Respiratory centers of the brainstem and methods of viral entry into the nervous system. A: The brainstem contains the respiratory centers which integrate signals from the cortex through the limbic system and regulate breathing. Motor efferents from the brain stem innervate the diaphragm, airways, and abdominal muscles which create the movements necessary for inhalation and exhalation; B: olfactory neurons are susceptible to infection by viral particles in the nasal cavity (Route 1). These sensory neurons cross the cribriform plate and project through the olfactory tract into areas of the cortex and limbic system; C: alveoli in the lung are innervated by sensory neurons projecting up the spinal cord to the brainstem. The sensory neurons that innervate the lungs directly gather chemo- and mechano-signals from their dendrites in the lung and project them to pre-synapses in the brainstem. These neurons detect changes in oxygen and CO₂, mechanical stretch, and irritants that may be inhaled into the lung. Viral particles may enter these neurons and be transported anterogradely to the respiratory centers of the brainstem (Route 2). They may also enter the bloodstream and be subsequently introduced to the brain parenchyma through the blood-brain barrier (BBB) (Routes 3 and 4); D: a viral infection in the bloodstream may cross the BBB by active transport via receptors on the surface of endothelial cells (Route 3) or by passive diffusion through a leaky barrier (Route 4). Immune cells in the blood and glial cells in the brain respond to infection by releasing inflammatory cytokines, which may damage the BBB and result in greater passive diffusion of the virus. Arrows indicate the direction of signal transmission (A). Arrows indicate the direction of virus movement (B, C, D). Numerals 1-4 indicate different proposed routes of viral transmission. PNC: pneumotaxic center; APC: apneustic center; PRG: pontine respiratory group; pFL: lateral parafacial nucleus; PiCo: postinspiratory complex; PreBötz: PreBötzinger complex; VRG: ventral respiratory group; DRG: dorsal respiratory group

in the pons and medulla. This control arises from neural circuits projecting from the cerebral cortex to the brainstem^[15]. In addition to voluntary control of breathing, a variety of sensory pathways contribute to the breathing pattern^[16]. Normal breathing consists of an inexorable active inspiration phase arising from rhythmic neural activity in the preBötzinger (preBötz) complex^[17]. The preBötz is a region of the ventrolateral medulla which is active in phase with inspiration. Neurons in the preBötz project either

directly or indirectly to motor neurons which innervate the diaphragm, intercostals, and airway muscles [Figure 1A]. Rhythmic limb movement has also been shown to modulate breathing through afferent sensory pathways^[18]. Of particular interest are vagal sensory neurons expressing chemoreceptors and mechanoreceptors, which project from the lungs and airways to respiratory centers of the brainstem^[19]. Due to their inherent proximity to the primary infection site, these sensory neurons present one potential mechanism by which respiratory viruses may enter the brain^[20]. Expiration is passive at rest. However, active expiration may be brought on by increased O₂ demand, such as respiratory distress^[21]. Disinhibition or activation of neurons in the lateral parafacial nucleus (pF_L), which project to expiratory premotor neurons, causes contraction of muscles, which reduces lung volume below their resting capacity^[22]. This reduced lung volume after active exhalation increases the volume of the subsequent breath resulting in greater O₂ delivery. A third phase, post-inspiration, may occur between the inspiration and exhalation phases. Post-inspiration is a delay of lung deflation which increases gas exchange in the lung^[23]. It may originate from interneurons medial to the parafacial nucleus, referred to as the post-inspiratory complex (PiCo)^[24]. Thus, infection of any of these regions of the brain may lead to respiratory distress or even failure.

EVIDENCE OF NEUROTROPISM OF SARS-COV-2

It has been postulated that, like other β coronaviruses, SARS-CoV-2 can also infect the brain by migrating from the general circulation to the cerebral microcirculation via endothelial cells which express ACE2^[5]. In a recent case study published by Poyiadji *et al.*^[25], they described an acute hemorrhagic necrotizing encephalopathy that was directly attributed to SARS-CoV-2 infection and the concurrent cytokine storm induced upon infection^[25,26]. Beyond isolated cases of overt severe neurological pathologies, 45% of severe cases exhibited neurologic symptoms^[3], increasing the likelihood that potential complications arising from severe SARS-CoV-2 infections may have a strong neurologic component that has yet to be described and characterized. Human coronavirus OC43 has been demonstrated to be transmitted in mice both passive diffusion of released viral particles and axonal transport^[27].

Possible pathways for SARS-CoV-2 invasion of the central nervous system

The major receptor for SARS-CoV-2, ACE2, is distributed in multiple tissues of the body. It is present in epithelial cells of alveoli and small intestine arterial and venous endothelial cells and arterial smooth muscle cells^[28] neuronal, glial and endothelial cells of the brain^[5]. Low expression is observed in glomerular tubular cells. Glomerular endothelial cells, Kupffer cells, hepatocytes, spleen, thymus, lymph nodes and immune cells do not show the expression of ACE2^[29]. Similar to SARS-CoV, SARS-CoV-2 uses the spike protein S1 to attach to the host cell ACE2 receptor^[13,30], but SARS-CoV-2 binds to the receptor with 10-20 fold higher affinity^[30]. A possible explanation for this increased affinity comes from the intense work of Shang *et al.*^[31]. By determining the crystal structure of the receptor-binding domain (RBD) of the spike protein (S1), they showed that, compared to SARS-CoV, SARS-CoV-2 receptor binding motif (RBM) contains a structural change in the ACE2-binding ridge caused by a four residue motif (residues 482-485: Gly-Val-Glu-Gly) due to which the ridge becomes more compact and forms more rigid contact with the N-terminal helix of ACE2^[31]. In addition, changes in several other residues in the SARS-CoV-2 RBD caused higher stabilization of two virus-binding hotspots at the RBD-ACE2 interface causing higher affinity of SARS-CoV-2 to ACE2^[31]. A comparison between SARS-CoV-2 RBD subdomain-1 (S1) with the RBD glycoprotein of bat coronavirus (RaTG13) and S1 protein of SARS coronavirus showed strong but not identical homology in the spike proteins of all three CoVs which can possibly explain the high binding affinity of SARS-CoV-2 to human ACE2 receptor^[5]. Ou *et al.*^[32] showed that the SARS-CoV-2 spike protein is less thermostable than SARS-CoV spike protein, suggesting its higher infectivity^[32]. Upon binding with the ACE2, SARS-CoV-2 S protein is primed by serine protease TMPRSS2^[33]. SARS-CoV-2 might be able to transigrate to the brain via the general circulation, the

olfactory pathway, or the peripheral neurons of the lungs. The schematic in [Figure 1](#) shows the possible pathways of virus entry to the central nervous system (CNS).

The olfactory pathway

A proposed pathway that the virus can take to reach the brain is the olfactory pathway [[Figure 1B](#)]. Total or partial loss of smell, either by “cytokine storm” or damage of the olfactory epithelium (OE), is an early indicator of SARS-CoV-2 infection. Damage of the OE is particularly important as the cells express both ACE2 and TMPRSS2^[34]. ACE2 is highly expressed in the non-neuronal goblet/secretory cells^[35] and ciliated receptor cells of the nasal epithelium^[5,36]. Upon binding with these receptors, the virus can travel through the cribriform plate to the olfactory bulb, move trans-synaptically to the cortex, and then to the brain stem^[5]. The non-neuronal cells in the OE possibly serve as reservoir of the virus^[13,34]. The major complication causing death in COVID-19 is respiratory failure and pneumonia. The CNS control of respiration, as described above, is an important factor in maintaining proper respiratory physiology. Failure of CNS to control the rate and depth of respiration may lead to irreversible serious consequences. Brain stem respiratory centers (RC) are the major neuronal groups controlling respiration. In a recent study Moberly *et al.*^[37] showed the anatomical and functional connectivity among the RC neurons, nasal epithelium and different brain regions including the olfactory bulb and limbic system^[37]. Thus, damage to any segment of this network is expected to cause respiratory distress as well as emotional and behavioral dysfunction. Direct evidence of olfactory transmission of SARS-CoV was produced by Netland *et al.*^[13]. By immunohistochemical methods they showed the presence of SARS-CoV in several brain regions of genetically modified mice including olfactory bulb, cerebral cortex, hippocampus and brain stem following intranasal infection of the virus. Most importantly, in case of low-grade infection, the virus was observed in the brain in the absence of lung infection or pneumonia^[13]. There is a 60 h delay between virus infection and the presence of virus in the olfactory bulb^[13], and this time is probably used by the virus for replication and accumulation in the non-neuronal OE cells^[34]. The subsequent transport of the virus to other regions of the brain was relatively faster, approximately 12 h-20 h^[13]. However, a recent study conducted by Bao *et al.*^[38] showing the respiratory transmission of the virus via liquid droplets from infected to uninfected hACE2 transgenic mice did not report any neural transmission or viral presence in the brain^[38].

The respiratory neuronal pathway

Lung-adjacent peripheral neurons may offer a second pathway for SARS-CoV-2 to enter the CNS [[Figure 1C](#)]. Due to their proximity to the primary infection site, these neurons are highly susceptible to viral invasion. The previously mentioned porcine HEV has been visually identified by transmission electron microscopy to infect sensory dorsal root ganglia^[39] and appears to be trans-synaptically transferred to the brain stem via membrane-coated vesicles^[40]. HEV shares 91% homology to hCoV-OC43, a close relative to SARS-CoV-2^[41], which provides some evidence that SARS-CoV-2 may also invade via a similar pathway. The direct synaptic connections between these sensory nerves and the respiratory centers in mammals may prove to be important in the characteristic respiratory failure seen in severe COVID-19 patients.

The circulatory pathway

Upon infection, SARS-CoV-2 has been detected in the general circulation from which it can be transported to the cerebral circulation^[5]. Endothelial cells (EC) of the blood-brain barrier (BBB) express the ACE2 receptor. The virus then binds with the receptor to enter the ECs and multiply. Eventual budding enables the virus to infect the brain parenchyma, especially the neuronal cells^[5,42], which also express ACE2 receptors. The damaged capillaries cause micro-hemorrhage, which can have fatal consequences. Once in the neurons, the viruses multiply and bud off the neurons killing the cells and infecting many more cells in the brain, which leads to functional deficits [[Figure 1D](#)]. In the case of SARS-CoV infection, Netland *et al.*^[13] showed that the virus causes neuronal damage without evoking a substantial inflammatory response^[13]. In a small subset of severe influenza cases and various other viral infections, an associated cytokine storm

Table 1. Currently recruiting clinical trials involving MSCs¹

NCT number (Phases)	Interventions	Enrollment (age)	Location
NCT04313322 (Phase 1)	Wharton's Jelly MSCs	5 (18+)	Stem Cells Arabia, Amman, Jordan
NCT04336254 (Phase 1,2)	Allogeneic human dental pulp stem cells vs. saline injection (Placebo)	20 (18-65)	Renmin Hospital of Wuhan University (East Campus), Wuhan, Hubei, China
NCT04288102 (Phase 2)	MSCs vs. 1% Human serum albumin in saline	90 (18-75)	Maternal and Child Hospital of Hubei and Huoshenshan Hospital, Wuhan, Hubei, China
NCT04252118 (Phase 1)	MSCs	20 (18-70)	Beijing 302 Military Hospital of China, Beijing, China
NCT04366271 (Phase 2)	Standard of Care vs. MSC	106 (40-80)	Hospital Infantil Universitario Nino Jesus, Madrid, Spain
NCT04355728 (Phase 1/2)	UCMSCs	24 (18+)	University of Miami, USA
NCT04366063 (Phase 2/3)	MSC Therapy	60 (18-65)	Royan Institute, Tehran
NCT04339660 (Phase 1,2)	CMSCs vs. Placebo	30 (18-75)	Puren Hospital Affiliated to Wuhan University of Science and Technology, Wuhan, Hubei, China
NCT04392778 (Phase 1/2)	MSCs vs. Saline Control	30 (40-60)	Istinye University, Istanbul, Turkey
NCT04390139 (Phase 1/2)	Excel-UMC-Beta vs. Placebo	30 (18-75)	Hospital DE Bellvitge, Barcelona, Spain
NCT04371393 (Phase 3)	Remestemcel-L vs. Placebo	300 (18+)	University of Southern California, Los Angeles, CA and Ichan School of Medicine at Mount Sinai, NY, USA
NCT03042143 (Phase 1/2)	UCMSCs (CD362 Enriched) vs. Placebo	75 (16+)	Belfast Health and Social Care Trust, Royal Hospital, Northern Ireland, UK
NCT04269525 (Phase 2)	Umbilical Cord MSCs	10 (18-75)	Zhongnan Hospital of Wuhan University, Wuhan, Hubei, China
NCT04361942 (Phase 2)	MSC vs. Placebo	24 (18+)	Hospital Universitario Rio Hortega, Valladolid, Spain
NCT04333368	Umbilical cord Wharton's Jelly MSCs vs. NaCl 0.9%	60 (18+)	Hôpital Pitié-Salpêtrière and Européen Georges Pompidou, Paris, France

¹ClinicalTrials.gov; accessed 27 May 2020. MSCs: mesenchymal stem cells

and subsequent BBB breakdown has been described as a known passive method of entry for viral particles causing acute necrotizing encephalopathy in some patients^[43].

NEUROTROPISM-TARGETED COVID-19 TREATMENT APPROACH

Thus far, there have been ~300 clinical trials registered in the National Institute of Health (www.clinicaltrials.gov) on treatments for SARS-CoV-2. Most of these trials focus on antiviral drugs, without any due consideration of the neurotropism of the virus. To our knowledge there are more than 160 antiviral agents available^[44]. Screening of these compounds with the intent to identify those that lack neurodegenerative properties or those that possess neuro-regenerative effects may be useful in identifying safe and effective drugs for SARS-CoV-2. It is interesting that one small clinical study relating to SARS-CoV-2 of the effects of mesenchymal stem cells (MSCs), which are known to exert anti-inflammatory and neuro-regenerative activities in brain disorders, showed excellent efficacy, i.e., all 7 of 7 patients showed symptom reduction in 2 days after MSC transplantation^[45]. This study has provided a basis for several MSC-based either observational or phase 1-3 trials, which are currently recruiting [Table 1]. In addition, there are several other clinical trials which are registered but have not yet begun to recruit. Given the pivotal role of neuropathology of SARS-CoV-2 and neuronal control of breathing, future therapies should target neurotropism of the virus and combine anti-neuroinflammatory and neuro-regenerative features.

FUTURE THERAPEUTIC IMPLICATIONS

Several lines of evidence on coronaviruses and exploratory studies in SARS-CoV-2 have demonstrated the involvement of viral neurotropism in the brain stem respiratory center, whose components control

breathing, but it is unclear which of these three different pathways offer a potential method of invasion for the virus, and they are not necessarily mutually exclusive. The viral entry of SARS-CoV-2 may be via circulatory in one patient and via oronasal or peripheral sensory neurons in another. Each of these pathways warrants further investigation to determine the neurotropism of SARS-CoV-2, and how these pathways may be exploited to treat patients in the future.

Also, it is evident that one particular approach that is under intense investigation is the development of potential SARS-CoV-2 treatment using MSCs, which have shown neuro-regenerative properties in several brain disorders including traumatic brain injury (TBI)^[46]. One of the issues in using MSCs in inflammatory conditions is that MSCs undergo apoptosis or differentiation due to cytokines being released in those inflammatory tissues. It has been reported that pretreatment of TBI animals first with an anti-inflammatory agent followed by MSC treatments significantly increases the efficacy of MSCs^[46]. Whether a similar strategy would be more successful in COVID-19 remains to be elucidated. Given the pace and expansiveness of the COVID-19 super pandemic and the fact that the rebound infection may continue for several months to years or could be a seasonal respiratory virus infection similar to influenza and respiratory syncytial virus, it may be prudent to test these possibilities.

DECLARATIONS

Authors' contributions

Wrote the manuscript: Das M, Penn C, Martinez T, Mayilsamy K, McGill A

Made the illustration: Penn C

Reviewed the manuscript: Wiling A, Mohapatra SS, Mohapatra S

Approved the final version of the manuscript: All authors

Availability of data and materials

Not applicable.

Financial support and sponsorship

This work is partly supported by a University of South Florida Pandemic Response Research Network funding for COVID-19 and Veterans Affairs Merit Review grant (No. BX002668) to Dr. Subhra Mohapatra, and Research Career Scientist Awards to Dr. Subhra Mohapatra (No. IK6BX004212) and Dr. Shyam Mohapatra (No. IK6BX003778). Though this report is based upon work supported, in part, by the Department of Veterans Affairs, Veterans Health Administration, Office of Research and Development, the contents of this report do not represent the views of the Department of Veterans Affairs or the United States Government.

Conflicts of interest

All authors declared that there are no conflicts of interest.

Ethical approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Copyright

© The Author(s) 2020.

REFERENCES

1. Zhou P, Yang XL, Wang XG, Hu B, Zhang L, et al. A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature* 2020;579:270-3.
2. Wang Z, Yang B, Li Q, Wen L, Zhang R. Clinical features of 69 cases with coronavirus disease 2019 in Wuhan, China. *Clin Infect Dis* 2020; doi: 10.1093/cid/ciaa272.
3. Mao L, Wang M, Chen S, He Q, Chang J, et al. Neurological manifestations of hospitalized patients with COVID-19 in Wuhan, China: a retrospective case series study. *medRxiv* 2020;2020.02.22.20026500.
4. Rabin RC. Some coronavirus patients show signs of brain ailments. *N Y Times*, April 1 2020. <https://www.nytimes.com/2020/04/01/health/coronavirus-stroke-seizures-confusion.html>
5. Baig AM, Khaleeq A, Ali U, Syeda H. Evidence of the COVID-19 virus targeting the CNS: tissue distribution, host-virus interaction, and proposed neurotropic mechanisms. *ACS Chem Neurosci* 2020; doi: 10.1021/acscchemneuro.0c00122.
6. Li YC, Bai WZ, Hashikawa T. The neuroinvasive potential of SARS-CoV2 may play a role in the respiratory failure of COVID-19 patients. *J Med Virol* 2020; doi: 10.1002/jmv.25728.
7. Rodriguez-Morales AJ, Cardona-Ospina JA, Gutierrez-Ocampo E, Villamizar-Pena R, Holguin-Rivera Y, et al. Clinical, laboratory and imaging features of COVID-19: a systematic review and meta-analysis. *Travel Med Infect Dis* 2020; doi: 10.1016/j.tmaid.2020.101623:101623.
8. Lahiri D, Ardila A. COVID-19 pandemic: a neurological perspective. *Cureus* 2020;12:e7889.
9. Song Z, Xu Y, Bao L, Zhang L, Yu P, et al. From SARS to MERS, thrusting coronaviruses into the spotlight. *Viruses* 2019;11.
10. Li K, Wohlford-Lenane C, Perlman S, Zhao J, Jewell AK, et al. Middle East respiratory syndrome coronavirus causes multiple organ damage and lethal disease in mice transgenic for human dipeptidyl peptidase 4. *J Infect Dis* 2016;213:712-22.
11. Tsai LK, Hsieh ST, Chang YC. Neurological manifestations in severe acute respiratory syndrome. *Acta Neurol Taiwan* 2005;14:113-9.
12. Kim JE, Heo JH, Kim HO, Song SH, Park SS, et al. Neurological complications during treatment of Middle East respiratory syndrome. *J Clin Neurol* 2017;13:227-33.
13. Netland J, Meyerholz DK, Moore S, Cassell M, Perlman S. Severe acute respiratory syndrome coronavirus infection causes neuronal death in the absence of encephalitis in mice transgenic for human ACE2. *J Virol* 2008;82:7264-75.
14. Bleau C, Filliol A, Samson M, Lamontagne L. Brain invasion by mouse hepatitis virus depends on impairment of tight junctions and beta interferon production in brain microvascular endothelial cells. *J Virol* 2015;89:9896-908.
15. Butler JE. Drive to the human respiratory muscles. *Respir Physiol Neurobiol* 2007;159:115-26.
16. Bordoni B, Marelli F, Bordoni G. A review of analgesic and emotive breathing: a multidisciplinary approach. *J Multidiscip Healthc* 2016;9:97-102.
17. Yang CF, Feldman JL. Efferent projections of excitatory and inhibitory preBotzinger Complex neurons. *J Comp Neurol* 2018;526:1389-402.
18. Shevtsova NA, Marchenko V, Bezdudnaya T. Modulation of respiratory system by limb muscle afferents in intact and injured spinal cord. *Front Neurosci* 2019;13:289.
19. Chang RB, Storch DE, Williams EK, Umans BD, Liberles SD. Vagal sensory neuron subtypes that differentially control breathing. *Cell* 2015;161:622-33.
20. McGavern DB, Kang SS. Illuminating viral infections in the nervous system. *Nat Rev Immunol* 2011;11:318-29.
21. Saini JK, Pagliardini S. Breathing during sleep in the postnatal period of rats: the contribution of active expiration. *Sleep* 2017;40.
22. Huckstepp RT, Henderson LE, Cardoza KP, Feldman JL. Interactions between respiratory oscillators in adult rats. *Elife* 2016;5.
23. Dutschmann M, Jones SE, Subramanian HH, Stanic D, Bautista TG. The physiological significance of postinspiration in respiratory control. *Prog Brain Res* 2014;212:113-30.
24. Anderson TM, Garcia AJ 3rd, Baertsch NA, Pollak J, Bloom JC, et al. A novel excitatory network for the control of breathing. *Nature* 2016;536:76-80.
25. Poyiadji N, Shahin G, Noujaim D, Stone M, Patel S, et al. COVID-19-associated acute hemorrhagic necrotizing encephalopathy: CT and MRI features. *Radiology* 2020; doi: 10.1148/radiol.20201187:201187.
26. Das G, Mukherjee N, Ghosh S. Neurological insights of COVID-19 pandemic. *ACS Chem Neurosci* 2020; doi: 10.1021/acscchemneuro.0c00201.
27. Dube M, Le Coupanec A, Wong AHM, Rini JM, Desforges M, et al. Axonal transport enables neuron-to-neuron propagation of human coronavirus OC43. *J Virol* 2018;92.
28. Hamming I, Timens W, Bulthuis ML, Lely AT, Navis G, et al. Tissue distribution of ACE2 protein, the functional receptor for SARS coronavirus. A first step in understanding SARS pathogenesis. *J Pathol* 2004;203:631-7.
29. Cheng H, Wang Y, Wang GQ. Organ-protective effect of angiotensin-converting enzyme 2 and its effect on the prognosis of COVID-19. *J Med Virol* 2020; doi: 10.1002/jmv.25785.
30. Wrapp D, Wang N, Corbett KS, Goldsmith JA, Hsieh CL, et al. Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation. *Science* 2020;367:1260-3.
31. Shang J, Ye G, Shi K, Wan Y, Luo C, et al. Structural basis of receptor recognition by SARS-CoV-2. *Nature* 2020;581:221-4.
32. Ou X, Liu Y, Lei X, Li P, Mi D, et al. Characterization of spike glycoprotein of SARS-CoV-2 on virus entry and its immune cross-reactivity with SARS-CoV. *Nat Commun* 2020;11:1620.
33. Hoffmann M, Kleine-Weber H, Schroeder S, Kruger N, Herrler T, et al. SARS-CoV-2 cell entry depends on ACE2 and TMPRSS2 and is blocked by a clinically proven protease inhibitor. *Cell* 2020;181:271-80.e8.

34. Butowt R, Bilinska K. SARS-CoV-2: olfaction, brain infection, and the urgent need for clinical samples allowing earlier virus detection. *ACS Chem Neurosci* 2020; doi: 10.1021/acscchemneuro.0c00172.
35. Olender T, Keydar I, Pinto JM, Tatarsky P, Alkelai A, et al. The human olfactory transcriptome. *BMC Genomics* 2016;17:619.
36. Sungnak W, Huang N, Bécavin C, Berg M; HCA Lung Biological Network. SARS-CoV-2 entry genes are most highly expressed in nasal goblet and ciliated cells within human airways. *Nat Med* 2020; doi: 10.1038/s41591-020-0868-6.
37. Moberly AH, Schreck M, Bhattarai JP, Zweifel LS, Luo W, et al. Olfactory inputs modulate respiration-related rhythmic activity in the prefrontal cortex and freezing behavior. *Nat Commun* 2018;9:1528.
38. Bao L, Gao H, Deng W, Lv Q, Yu H, et al. Transmission of SARS-CoV-2 via close contact and respiratory droplets among hACE2 mice. *J Infect Dis* 2020; doi: 10.1093/infdis/jiaa281.
39. Li YC, Bai WZ, Hirano N, Hayashida T, Hashikawa T. Coronavirus infection of rat dorsal root ganglia: ultrastructural characterization of viral replication, transfer, and the early response of satellite cells. *Virus Res* 2012;163:628-35.
40. Li YC, Bai WZ, Hirano N, Hayashida T, Taniguchi T, et al. Neurotropic virus tracing suggests a membranous-coating-mediated mechanism for transsynaptic communication. *J Comp Neurol* 2013;521:203-12.
41. Gonzalez JM, Gomez-Puertas P, Cavanagh D, Gorbalenya AE, Enjuanes L. A comparative sequence analysis to revise the current taxonomy of the family Coronaviridae. *Arch Virol* 2003;148:2207-35.
42. Palasca O, Santos A, Stolte C, Gorodkin J, Jensen LJ. TISSUES 2.0: an integrative web resource on mammalian tissue expression. *Database (Oxford)* 2018;2018.
43. Rossi A. Imaging of acute disseminated encephalomyelitis. *Neuroimaging Clin N Am* 2008;18:149-61; ix.
44. De Clercq E, Li G. Approved antiviral drugs over the past 50 years. *Clin Microbiol Rev* 2016;29:695-747.
45. Leng Z, Zhu R, Hou W, Feng Y, Yang Y, et al. Transplantation of ACE2(-) mesenchymal stem cells improves the outcome of patients with COVID-19 pneumonia. *Aging Dis* 2020;11:216-28.
46. Das M, Mayilsamy K, Tang X, Han JY, Foran E, et al. Pioglitazone treatment prior to transplantation improves the efficacy of human mesenchymal stem cells after traumatic brain injury in rats. *Sci Rep* 2019;9:13646.

Review

Open Access



An association between mitochondria and microglia effector function: what do we think we know?

Gaylia Jean Harry¹, Gabrielle Childers^{1,2}, Sahana Giridharan^{1,3}, Irisyunuel Lopez Hernandez¹

¹National Toxicology Program Laboratory, National Institute of Environmental Health Sciences, Durham, NC 27709, USA.

²(Current affiliation) Gabrielle Childers, University of Alabama at Birmingham, Birmingham, AL 35294, USA.

³Sahana Giridharan, Duke University, Durham, NC 27708, USA.

Correspondence to: Gaylia Jean Harry, National Institute of Environmental Health Sciences, P.O. Box 12233, MD E1-07, Durham, NC 27709, USA. E-mail: harry@nih.gov

How to cite this article: Harry GJ, Childers G, Giridharan S, Lopez Hernandez I. An association between mitochondria and microglia effector function: what do we think we know? *Neuroimmunol Neuroinflammation* 2020;7:150-65.
<http://dx.doi.org/10.20517/2347-8659.2020.07>

Received: 14 Jan 2020 **First Decision:** 24 Feb 2020 **Revised:** 7 Apr 2020 **Accepted:** 7 Apr 2020 **Available online:** 16 Jun 2020

Science Editor: Jeffrey Bajramovic **Copy Editor:** Jing-Wen Zhang **Production Editor:** Tian Zhang

Abstract

While resident innate immune cells of the central nervous system, the microglia, represent a cell population unique in origin, microenvironment, and longevity, they assume many properties displayed by peripheral macrophages. One prominent shared property is the ability to undergo a metabolic switch towards glycolysis and away from oxidative phosphorylation (OXPHOS) upon activation by the pro-inflammatory stimuli lipopolysaccharide. This shift serves to meet specific cellular demands and allows for cell survival, similar to the Warburg effect demonstrated in cancer cells. In contrast, normal surveillance phenotype or stimulation to a non-proinflammatory phenotype relies primarily on OXPHOS and fatty acid oxidation. Thus, mitochondria appear to function as a pivotal signaling platform linking energy metabolism and macrophage polarization upon activation. These unique shifts in cell bioenergetics in response to different stimuli are essential for proper effector responses at sites of infection, inflammation, or injury. Here, we present a summary of recent developments as to how these dynamics characterized in peripheral macrophages are displayed in microglia. The new insights provided by an increased understanding of metabolic reprogramming in macrophages may allow for translation to the central nervous system and a better understanding of microglia heterogeneity, regulation, and function.

Keywords: Mitochondria bioenergetics, inflammasome, microglia, pro-inflammatory, anti-inflammatory, polarization



© The Author(s) 2020. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, sharing, adaptation, distribution and reproduction in any medium or format, for any purpose, even commercially, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.



INTRODUCTION

The function of innate immune cells such as macrophages is to recognize and respond to a novel stimulus including microbial pathogens and sterile activators. These pro- and anti-inflammatory responses are major sources of soluble molecules, cytokines, hormones, and neuropeptides. These factors provide tools to sense, process, and relay physiological signals beyond their canonical roles. Macrophages display a heterogeneous repertoire to fulfill a broad range of functions in host defense, including tissue homeostasis and repair, pathology, and development. To accomplish this, innate immune cells adopt various activation phenotypes. Precise regulation of such activation is essential for maintenance of tissue homeostasis with governance accomplished by a balance of stimulatory and inhibitory signals. Multiple lines of evidence suggest an interlinked relationship between innate immunity and the integrity and function of mitochondria serving to maintain this homeostatic balance. Metabolic pathways provide the necessary energy and serve to regulate phenotype and function. Pro-inflammatory macrophages {M[LPS(+IFN γ)]} display an enhanced glycolytic metabolism and impaired mitochondrial oxidative phosphorylation (OXPHOS). These energy shifts place mitochondria in a pivotal signaling role in macrophage response to stimuli and circumventing immune checkpoint signals^[1-3]. The link between immunological and metabolic processes associated with mitochondria, immunometabolism, may influence activation states and polarization of myeloid cells to fine-tune their functions^[4-7].

Of the specialized cells of the central nervous system (CNS), basic host defense mechanisms exist predominantly in microglia as resident macrophages. Microglia share many phenotypic characteristics with peripheral macrophages yet are unique in their origin and molecular or transcriptional profile^[8-15]. The available literature on the immunometabolism of microglia, as compared to what is known of peripheral macrophages, is limited but growing to address questions of similarities and differences^[6,16]. It may also allow for a framework to understand the various other tasks undertaken by microglia during development and chronic maintenance. Here, we present a summary of how these dynamics characterized in peripheral macrophages are displayed in microglia. While much of this work is still somewhat under a “work in progress” classification, even in the peripheral macrophage, new insights provided by an increased understanding of metabolic reprogramming foster a better understanding of macrophage and microglia regulation and function.

MICROGLIA AND IMMUNE CELLS IN THE CNS

The mechanical separation of the CNS from the circulation by the blood-brain barrier^[17,18] influences immune responses^[19,20] by excluding many peripherally derived innate and adaptive immune cells and inflammatory molecules^[21]. However, infiltrating cells significantly contribute to any neuroinflammatory response following disruption of the blood-brain barrier, as can occur with physical injury or high levels of inflammation. In such cases, blood-borne monocytes are allowed to enter the brain parenchyma and, over time, can transition and assume a brain-specific phenotype^[22-24]. Additionally, with a T cell-mediated neuroinflammation, peripheral innate immune cells enter the brain as a protective host defense against infection and facilitate repair following stroke or physical trauma^[25,26]. In such a case, interactions between microglia and T cells can be signaled via interleukin (IL)-23 and IL-1 β , leading to the production of granulocyte-macrophage colony-stimulating factor (GM-CSF) to facilitate microglia proliferation in a manner to promote an appropriate level of response to injury^[27]. Recently, the identification of innate lymphoid cells in the brain suggests an additional innate immune cell population that may act to control neuroinflammation^[28]. Thus, in such conditions, the macrophage population likely represents a combination of resident microglia and infiltrating monocytes. As a distinction between these two populations, it has been suggested that resident microglia focus on tasks related to maintaining tissue homeostasis while infiltrating cells are involved in severe inflammatory injuries^[29,30].

SENSING AND RESPONDING TO THE ENVIRONMENT

While microglia appear to be tightly adapted to the specific requirements within brain regions, they all function in a surveillance mode with mobile processes extending into the surrounding microenvironment to detect tissue changes^[31,32]. Upon sensing such changes, microglia respond to their environment via several “sosome” genes, allowing them to sense and interact with their local environment^[8,14,33]. These somesome genes include those for putative purinergic receptors, *P2ry12* and *P2ry13*, transmembrane protein 119 (*Tmem119*), G-protein coupled receptor 34 (*Gpr34*), the C-type lectin receptor, the fractalkine receptor, *Cx3cr1*, sialic acid-binding immunoglobulin-type lectin H (*Siglec-h*), and triggering receptor expressed on myeloid cells 2 (*Trem2*). Siglec proteins contribute to immune regulation by binding sialic acid residues on neurons^[34] and TREM2 contributes via recruitment of the immunoreceptor tyrosine-based activation motif-containing adapter protein, DAP-12^[35]. The final response of the cell is dictated by the overall pattern of somesome gene activation.

The microglia host-response begins with the recognition of pathogen-associated molecular patterns (PAMPs) such as bacterial, viral, and protozoal products (protein lipid, nucleic acid, and carbohydrate). This occurs via pattern recognition receptors on the plasma membrane or in the endosomal compartments^[36], or by binding phagocytic scavenger receptors^[37] and macrophage antigen complex I (MAC1, CD11b/CD18), which is a pattern recognition receptor linked to the superoxide-generating enzyme NADPH oxidase^[38]. In the absence of microorganisms, a similar but sterile inflammatory response occurs often as a result of trauma, ischemia-reperfusion injury, or chemical exposure^[39-41]. Activation in the absence of microbial compounds occurs by endogenous molecules called danger-associated molecular patterns (DAMPs)^[42]. Molecules that function as DAMPs include nucleic acids, lipids, and proteins that normally are not present to immune cells until released or unmasked during cell death due to tissue injury. In the CNS, microglia responding in various neurodegenerative diseases in the absence of pathogen have been termed disease-associated microglia (DAM). Intracellular DAMPs include high mobility group box 1 (HMGB1) and peroxiredoxin family proteins. These damage signals can activate immune cells through three major families of intracellular recognition receptors: toll-like receptors, nucleotide-binding domain leucine-rich repeat containing proteins (also known as NOD-like receptors), and Rlg1-like receptors. Receptor activation induces specific pathways and the release of cytokines that contribute to injury mitigation^[43]. Microglia are influenced by a plethora of factors including receptor agonists^[44-48] and transcription factor inducers^[49]. It is thought that microglial receptors^[50-52] can act as molecular switches to control microglial responses and that many of these actions function through alterations in calcium signals^[53]. In all cases, the immediate response upon sensing DAMPs, PAMPs, or other damaging events requires a robust increase in metabolic demand to support actions that initially are beneficial to the homeostatic balance of the nervous system.

Injury-induced inflammatory processes are dynamic and demonstrate spatial and temporal heterogeneity^[54-56]. In general, characterization of the macrophage response is based on the nature of the activating stimulus and the resulting production of factors^[57]. A conceptual framework has been proposed that suggests the nature of the activating stimulus can drive a range of activation phenotypes^[58-62], and it has been used as a basis for characterizing cellular responses^[63-70]. While phenotypic activation-state distinctions are currently under scrutiny^[67,71], it has been shown that classically activated microglia associated with inflammation can be produced upon stimulation with agonists for toll-like receptors (e.g., lipopolysaccharide, LPS) or IFN γ receptors. In contrast, different aspects of the immune response that do not involve the classical response can be observed upon stimulation by IL-4 or IL-13 with the expression of anti-inflammatory cytokines (IL-4, IL-10, IL-13, and TGF- β), arginase-1 (Arg1), CD206, and Chitinase-3-like-3 (Ym1 in rodents)^[72-76]. It is considered that the different phenotypes may be related, yet have different roles in host defense, wound healing, and resolution of inflammation^[57,60]. Differences in metabolic processes have been identified across these different activation inducers, suggesting a role for mitochondria in phenotypic outcome^[77].

STIMULUS-DRIVEN METABOLIC RE-PROGRAMMING OF MICROGLIA

In a normal “resting” cell, energy demands are addressed with the conversion of glucose to pyruvate with entry into glycolysis. Pyruvate in the cytosol can be taken up by mitochondria and enters the tricarboxylic acid (TCA) cycle where it is oxidized to generate ATP. This provides a total energy gain of approximately 36 ATP per one molecule of glucose. In contrast, with hypoxia or anoxia, the cell has the ability to divert pyruvate away from mitochondria OXPHOS, allowing for ATP generation during low oxygen conditions. In this case, one glucose molecule will generate two pyruvate molecules that will be converted to lactate by lactate dehydrogenase in the cytosol^[78,79]. While this reaction generates significantly fewer molecules of ATP, glycolysis proceeds due to the production of NAD⁺. While less efficient, a beneficial effect of a shift to glycolysis is that it can be very quickly induced to meet cellular demands in cells with high glucose capacity^[80]. The importance of this shift was initially demonstrated in cancer cells in what is known as the Warburg effect^[81,82]. In cancer, malignant cells shift their demand for biosynthetic precursors and energy change and change their metabolic profile from a relatively low rate of glycolysis and the oxidation of pyruvate by the TCA cycle. The shift in metabolic profile is characterized by a lower rate of OXPHOS, high rate of glycolysis, and elevated lactic acid production. The high glycolytic rate induced during the Warburg effect is driven by the need to meet the increased demand for production of nucleotides and amino acids. While this effect was initially identified and characterized in cancer cells, a similar ability to utilize such a metabolic switch has been demonstrated in immune cells to meet increased energy demands when responding to infection or injury^[1].

There is now evidence suggesting a role for metabolic reprogramming by mitochondria in the maintenance and establishment of innate and adaptive immune responses^[75,83-96]. Given that immune cell populations depend on unique effector functions in response to distinct stimuli that often require production and secretion of high amounts of signaling factors and antimicrobial agents, it follows that changes in mitochondria function to meet these demands are crucial for efficient response to distinct contexts^[97-101]. It was initially observed that, upon activation, macrophages increase glycolysis and decrease oxygen consumption^[102,103]. It was further demonstrated that macrophage phenotype can be shifted by reprogramming glucose metabolism^[104,105], which helps meet energy demands required for shifting cell function and survival^[106].

Under normal conditions, microglia exist in a surveillance phenotype for constant monitoring of the parenchyma^[107,108] and preferentially rely on oxidative metabolism^[90,109,110]. Upon activation by LPS^[94], amyloid- β ^[111], and iron loading^[112,113], microglia switch their reliance on OXPHOS metabolism^[69,110,114,115] towards glycolytic metabolism to maintain mitochondrial function and ensure cell survival^[91,94,95,116]. Voloboueva *et al.*^[94] showed that, upon stimulation by LPS, BV-2 microglia increased lactate production and decreased mitochondria oxygen consumption and ATP production. This shift was reported to be modulated by mitochondrial glucose-regulated protein 75/mortalin^[94]. Exposure to a combination of LPS and IFN- γ increases nitric oxide formation, glucose consumption, hexokinase activity, glucose-6-phosphate dehydrogenase activity, phosphofructokinase-1 activity, lactate dehydrogenase activity, and lactate release, suggesting potentiated glycolysis^[94]. Similar findings were reported by Orihuela *et al.*^[69]: following LPS, BV2 microglia and primary murine microglia shifted from a primary oxidative metabolic towards glycolytic metabolism with no evidence of cell death. An increase in microglial mitochondria has been observed with activation^[117,118], implicating an association with mitochondria biogenesis. Recent studies have suggested that a shift in glycolysis in microglia is accompanied by an increase in the enzyme PFKFB3, which is responsible for activation of phosphofructokinase^[119]. Additionally, this metabolic shift has been found to be regulated by the anti-inflammatory cytokine IL-10 for aerobic glycolysis inhibition and OXPHOS^[120].

In a non-classical activation state, macrophages use oxidative metabolism for functions involved in normal maintenance functions, tissue repair, and wound healing^[73,121,122]. In IL-4 stimulated macrophages [M(IL-4)],

the Krebs cycle and OXPHOS remain intact. In addition, the cells are able to utilize fatty acid oxidation and oxidative respiration for energy production^[98,123-125] and arginine metabolism is shifted to ornithine and polyamines^[126]. Work by Ferger *et al.*^[118] suggested that the stability of the electron transport chain in mitochondria plays a more substantial and critical role for the microglia response to IL-4 as compared to the response to LPS. In microglia, exposure to LPS induced a rapid and transitory decrease in the mitochondrial uncoupling protein-2 (UCP-2) levels accompanied by increased mitochondrial reactive oxygen species (mtROS) production. In UCP-2-silenced microglia, the response to LPS was exacerbated and a response to IL-4 was eliminated^[127]. An earlier study examining the translation of responses in macrophages to microglia reported reduced glucose consumption and lactate production in BV-2 cells exposed to IL-4^[109]. It was suggested that this phenotype was associated with phagocytosis of debris and the reduced need for anabolic reactions. Similar findings were reported by Orihuela *et al.*^[69] with exposure of BV-2 cells or primary murine microglia to IL-4/IL-13 in that the cells remained within an oxidative metabolic state with OCR and ECAR levels similar to non-stimulated cells. There were also elevations in mRNA levels for *Ym1*, *Il4*, *Cd163*, and *Arg1*, but no induction of *Tnfa* or *Il1*^[69]. The lack of a demonstrated metabolic shift with IL-4/IL-13 stimulation is in contrast to observations in peripheral macrophages of stimulated glucose uptake in addition to fatty acid metabolism and shift in mitochondrial biogenesis^[125].

MITOCHONDRIA AND FREE RADICAL PRODUCTION

A key feature of classically activated macrophages is their ability to produce reactive oxygen species (ROS) to facilitate killing of phagocytized bacteria^[128]. Stimulation of macrophages with LPS and IFN- γ increases inducible nitric oxide synthase (iNOS), generating nitric oxide (NO), a reactive nitrogen species that can inhibit mitochondrial respiration by nitrosylating iron-sulfur proteins in electron transport chain complexes and cytochrome c oxidase^[129,130]. It is considered that iNOS and NO-mediated inhibition of mitochondrial metabolism in macrophages is essential for the metabolic switch activated by LPS. This is not as well established in microglia, especially given that, while nitric oxide production is often linked with pro-inflammatory cytokines, such cytokines can be stimulated by sterile activators in the absence of NO. In microglia, it has been proposed that activation of the rapamycin (mTOR) pathway may actively contribute to this process^[131] as well as pro-inflammatory cytokine production and phagocytic activity^[132-134]. The resulting elevated thiamin pyrophosphate activity increased production of purines and pyrimidines, which yield nicotinamide adenine dinucleotide phosphate (NADPH) for the NADPH oxidase enzyme and ROS production^[135] implicated in the transition of microglia to a pro-inflammatory phenotype^[136-139]. It has been proposed that glycolytic ATP production may utilize the electron transport chain to compensate for this shift towards ROS production^[128]. It is known that superoxide produced by NADPH oxidase is predominantly extracellular. *In vivo*, extracellular superoxide dismutase 3 (SOD3) forms membrane permeable H₂O₂. Studies have suggested that H₂O₂, rather than SOD, serves as the primary ROS involved in mediating microglial activation and proliferation in response to pro-inflammatory stimuli^[136,140,141]. H₂O₂ has also been implicated in the increase in CD11b expression both *in vitro* and *in vivo*^[142], as well as in persistent neuroinflammation related to impaired NF- κ B p50 function^[143]. Superoxide anion is the primary ROS produced by mitochondria and mitochondria-derived H₂O₂ and, in addition to NADPH oxidase, may contribute to a pro-inflammatory phenotype of microglia such as that observed with the mitochondrial toxin, rotenone^[144]. With a response sufficient to result in ROS production, the associated intracellular damage is limited by increased generation of NADPH required for maintenance of reduced glutathione and nitric oxide production^[145,146].

GLUCOSE

In addition to the critical role that glucose plays in energy metabolism, it serves as an exclusive substrate for the hexose monophosphate shunt, which produces NADPH that is required by glutathione reductase to convert oxidized glutathione (GSSG) back to reduced glutathione (GSH). It also serves to quench ROS and

repair oxidative damage through glutathione- and thioredoxin-coupled pathways^[147]. Glucose metabolism influences microglial activation through an NADH-sensitive co-repressor termed C-terminal binding protein (CtBP). Slowed glucose flux through glycolysis reduces NADH levels and reduce NADH:NAD⁺ ratio^[148]. In both microglia and macrophage RAW264.7 cells, glucose flux regulates iNOS expression and other pro-inflammatory genes through effects on cytosolic NADH:NAD⁺ ratio and CtBP^[149].

Several glucose transporters such as GLUT1^[150], GLUT3^[151], and GLUT5^[152] are expressed in microglia. Acute fluctuation of available glucose impacts microglia activity with an elevated response to LPS upon shifting from a normal to high glucose level. Shifting from a high to normal glucose level can also induce metabolic stress^[153]. Glucose levels can influence pro-inflammatory gene transcription by several mechanisms. One such mechanism relies on the formation of advanced glycation end-products (AGE). These products consist of modified proteins and lipids as a result of non-enzymatic reactions with sugars. It is known that microglia express receptors of AGE and, upon activation, pro-inflammatory signaling pathways are stimulated^[127,154]. In peripheral macrophages, it has been reported that a shift in the cell's energy source induced by glucose deprivation results in an altered response to a pro-inflammatory stimulus^[155-157]. Multiple studies have reported an inability of microglia to respond appropriately to LPS under oxygen and glucose deprivation or with 2-DG inhibition of glucose metabolism^[109]. However, there is evidence that microglia are capable of functioning with alternative energy sources to adequately respond to an inflammatory challenge. Choi *et al.*^[158] reported an increase in mRNA and protein levels for IL-6 in microglia after 7 h of glucose and serum free medium. Upon stimulation with LPS, glucose-deprived microglia retained their normal ability to respond with elevations in nitrite, IL-1 β , and TNF α ^[159]. Primary rat microglia shifted to glucose-free medium for 1 h to LPS showed an exacerbated release of NO within 24 h and similar elevations in TNF α and IL-1 β as compared to non-glucose-deprived cells. Glucose deprivation for 24 h prior to LPS exposure increased release of IL-1 β with no deficits in NO or TNF α . The authors suggested that microglia were able to mobilize fatty acids from intracellular lipid droplets as an energy source. The majority of studies examining the effects of glucose deprivation have focused on relatively short-term exposures, within 1-24 h. While these studies demonstrated that both peripheral macrophages and microglia can shift their response to a pro-inflammatory stimulus in a selective manner, the question remains as to whether such a response would be altered when the cells were forced to a more prolonged shift in energy metabolism. When RAW 264.7 [Figure 1] or BV-2 [Figure 2] cells were maintained for three days under culture conditions to force cells to rely on galactose as an alternative energy source, the cells were able to normally respond to LPS stimulation. However, the diminished pro-inflammatory cytokine response observed when 2-DG was used in previous studies to inhibit glycolysis may have been related to the lower basal OXPHOS induced^[160]. The differences across these studies likely lie with the method of depleting glucose: removing glucose from the medium; the addition of 2-DG, which in and of itself can lower basal induction of OXPHOS^[160]; or the combination of glucose deprivation with hypoxia. In RAW cells, the morphological changes observed with LPS activation have been demonstrated to be diminished under galactose, suggesting a requirement of glucose to facilitate cell spreading^[161]. This was not clearly observed in the current study where similar LPS-induced morphological patterns were observed in the absence of pyruvate [Figure 1]. In BV-2 cells, a slight morphological shift was observed with the low level of LPS stimulation with minimal induction of nitrate and an elevation in TNF α and IL-1 protein.

GLUTAMINE

Macrophages utilize glutamine at high rates to synthesize amino acids, nucleotides, NADPH, and energy production and are dependent upon extracellular sources of the amino acid^[103]. Channeling of glutamine into the Krebs cycle is a primary route to promote succinate synthesis in macrophages. This occurs with glutamine being used for synthesis of glutamate, GABA, and succinate, bypassing the TCA cycle^[155]. This stabilizes hypoxia-inducible factor 1- α (HIF-1 α), an oxygen-sensitive transcription factor that allows the




Figure 1. Response of RAW 264.7 cells to LPS under glucose or galactose medium conditions. RAW 264.7 cells were plated in 6-well tissue culture plates (Corning, Corning, NY) and maintained in normal growth medium (NM) [DMEM (Gibco, ThermoFisher, Waltham, MA) containing 4.5 g/L glucose, 2 mM L-glutamine, sodium pyruvate, supplemented with 100 U/mL penicillin/ streptomycin (Sigma-Aldrich, Burlington, MA) and 10% fetal bovine serum (FBS #1 00-106, 0.25 endotoxin units/mL; Gemini Bio-Products, Sacramento, CA) and allowed to reach 85% confluence over 3-5 days. Cells were maintained at 37 °C, 5% CO₂/5% O₂, 90% humidity (Nu-5831 tri-gas incubator, Nuaire, Plymouth, MN). NM was changed to phenol-free NM medium or phenol-free DMEM medium containing 2 mM L-glutamine and 100 U/mL penicillin/streptomycin supplemented with (2) high glucose (25 mM; Glu), or 3] galactose (10 mM; Gla). Cells were maintained in the experimental medium for 3 days following which, under the same media conditions, cells were exposed to lipopolysaccharide (LPS; 1 µg/mL; Sigma) for up to 18 h and monitored using a live cell imaging system (IncuCyte, Sartorius) under normal incubator conditions. (A) Representative images (20x) of cell morphology under normal medium (NM), high glucose (Glu), or galactose (Gla). Scale bar = 25 microns. (B) Samples were collected at 3 h post-LPS and mRNA isolated using TRIzol® Reagent (Invitrogen, Carlsbad, CA), and 2.5 µL cDNA was used for qRT-PCR for *Tnfa*, *Il1a*, *Il1b*, and *cyclophilin A* using TaqMan™. Individual gene expression levels were normalized to cyclophilin A and presented as fold-change from vehicle controls in each medium condition. Data were analyzed by 2-way ANOVA followed by Dunnett's test for independent group mean comparisons. Data represent mean ± SEM (*n* = 3-4). *Significance level as compared to vehicle control set at *P* < 0.05

cell to adapt to a hypoxic environment^[162]. It has been suggested that HIFs function to facilitate cross-talk between inflammation and metabolism^[163]. HIF-1α can induce the expression of pro-inflammatory cytokines and has been proposed to serve a role in shifting glycolytic pathways to favor anaerobic metabolism^[114]. With classic activation, succinate regulates HIF-1α to drive a sustained production of IL-1β^[100] and the subsequent cell actions serve to maintain the macrophage survival. For the non-classical phenotype, glutamine metabolism acts at multiple levels including the generation of α-ketoglutarate and serves as a substrate for UDP-GlcNAc synthesis^[164].

NLRP3 INFLAMMASOME

One biological response to an inflammatory event that is critically dependent upon metabolic regulation is inflammasome activation. This is especially relevant for inflammasomes that require prior cell priming for full activation upon a secondary stimulus. For example, glycolytic rate can influence formation of NLRP1 and NLRP3 inflammasomes in macrophages^[165,166]. Inflammasomes are multiprotein complexes formed in the cytosol of immune and neural cells in response to pathogenic and danger signals. They consist of a cytosolic sensor belonging to the AIM2 (absent in melanoma 2), or NLR, an adaptor protein




Figure 2. Response of BV-2 cells to LPS under glucose or galactose medium conditions. BV-2 murine microglia cells were plated in 6-well tissue culture plates (Corning, Corning, NY) and maintained in normal growth medium (DMEM (Gibco, ThermoFisher, Waltham, MA) containing 4.5 g/L glucose, 2mM L-glutamine, supplemented with 100 U/mL penicillin/streptomycin (Sigma- Aldrich, Burlington, MA) and 10% fetal bovine serum (FBS #100-106, 0.25 endotoxin units/mL; Gemini Bio-Products, Sacramento, CA) and allowed to reach 85% confluence over 3-5 days. Cells were maintained at 37 °C, 5% CO₂/5% O₂, 90% humidity (Nu-5831 tri gas incubator, Nuair, Plymouth, MN). The normal growth medium was changed to phenol-free complemented DMEM medium containing 2mM L-glutamine, supplemented with 100 U/mL penicillin/ strepto supplemented with 1) high glucose (25 mM), 2) low glucose (10 mM), 3) galactose (10mM), or 4) high glucose + galactose). Cells were maintained in the experimental medium for 3 days following which, under the same media conditions, cells were exposed to lipopolysaccharide (LPS; 100 ng/mL; Sigma) monitored using a live cell imaging system (IncuCyte) under normal incubator conditions for up to 18 h. A: Representative images (20x) of cell morphology under high glucose (Hi Glu), low glucose (Lo Glu), or galactose (Gal). Scale bar = 200 microns; B: Nitrite accumulation in culture medium was measured as an indirect indicator of nitric oxide synthesis using a Greiss Reagent kit (Promega, Madison, WI) following manufacturer's instructions. Estimates were determined relative to standard curve; C: Estimates of the release of IL-1 into the medium were obtained using HEX Blue™ IL-1R cells (InvivoGen, San Diego, CA) following manufacturer's instructions. From 20 μL aliquot of the medium, after a 1.5 h incubation at 37 °C, absorbance at 620 nm was measured using a BioTeck Synergy 4 plate reader. Data were calculated relative to background control; D: TNFα protein levels were determined by Mouse TNFα ELISA MAX kit (BioLegend, San Diego, CA) with BD OptEIA Reagent Set B according to manufacturer's instructions. Absorbance was measured at 450nm with a 570nm background subtraction. Protein levels were determined based on standard curve and calculated relative to total cell number as determined using IncuCyte software. Data represent mean ± SEM (n = 5-6). Data were analyzed by 2-way ANOVA followed by Bonferroni post-hoc comparisons. *Significance level as compared to vehicle control set at P < 0.05

ASC (apoptosis-associated speck-like protein containing a CARD), and an effector caspase, primarily caspase-1. There are a variety of inflammasomes, most of which fall into the NLR domain^[167]. NLRP1 and AIM2 inflammasomes have been characterized in neurons^[168-170] and the NLRC4, NLRP2, and NLRP3 inflammasomes in astrocytes^[171,172]. Components for multiple inflammasomes are expressed in microglia^[173,174]. The NLRP3 inflammasome responds to a number of activators, including sterile activators,

such as asbestos, silica crystals, aluminum salts, and polystyrene nanoparticles^[175-178], and aberrant proteins, such as extracellular A β ^[179], thereby contributing to a broad range of common inflammatory pathologies and chronic inflammation.

The NLRP3 inflammasome responds to metabolic regulation^[180] and has been increasingly recognized as a bridge between mitochondrial damage sensing and pro-inflammatory signaling within monocytes, including microglia^[181,182]. Unlike most inflammasomes, NLRP3 typically requires a two-step activation and it is this process for which there is mounting evidence that mitochondrial damage plays a contributing role^[101]. Activation of TLR, tumor necrosis factor receptor, or interleukin-1 receptor (IL-1R) initiates an intracellular cascade of effects, including activation of NF- κ B. This upregulates NLRP3 and pro-IL-1 β within the cell and facilitates post-transcriptional changes to NLRP3 to free ubiquitinated binding sites by BRCC3 (BRCA1/2-containing complex subunit 3)^[176,183,184]. Delivery of a secondary “trigger” such as PAMPs, DAMPs, or intact pathogens to the “primed” cell causes the release of the repressed state of NLRP3. Upon release, NLRP3 activates the inflammasome forming a multiprotein complex comprised of the cytosolic sensor NLRP3, ASC, and caspase 1^[185]. Caspase 1 facilitates the cleavage of the pro-forms of IL-1 β and IL-18^[186], resulting in the release of mature protein^[187]. The release of active IL-1 family cytokines is normally related to pyroptotic cell death; however, in the absence of cell death, hyperactivity of cells and the recruitment of a process dependent on plasma membrane-localized pores can result in similar protein release^[188,189]. While inflammasome activation is an efficient producer of mature IL-1 β , inflammasome independent mechanisms exist, including cathepsin B or caspase 11 dependent pathways^[190,191], bacterial pore-forming toxins, and extracellular ATP^[177]. Thus, an upregulation of mature IL-1 β does not automatically indicate an inflammasome mechanism. In addition to the release of inflammatory factors, the physical release of ASC specks into the extracellular environment represents a stimulus for activating phagocytic cells in the immediate environment, thus contributing to a prolonged propagation of inflammation^[192] or other biological responses^[193].

Induction of mitophagy, the process by which cells clear damaged mitochondria, has been implicated in inhibition of NLRP3 signaling^[194]. Release of oxidized mitochondrial DNA (mtDNA) produced during the priming stage^[195,196] can interact with the NLRP3 receptor and induce inflammasome activation. Nakahira *et al.*^[196] reported that inhibition of mitophagy in macrophages heightened the NLRP3 inflammasome activation in parallel with uncleared mitochondrial DNA released into the cytosol. The mitochondrial cytidine/uridine monophosphate kinase-2 (CMPK2) is a nucleotide kinase required for mtDNA synthesis and production of oxidized mtDNA fragments. These fragments can act as activating ligands for the NLRP3 inflammasome complex^[197]. In addition, the release of mtROS triggered by small molecule inhibition of complex I and III has been associated with NLRP3 inflammasome activation^[198]. The association between mtROS as a trigger for NLRP3 inflammasome activation remains controversial given potential off-target effects of mtROS inhibitors. While studies have reported a role for mtROS in NLRP3 inflammasome activation, other conflicting studies have been reported. At least one study reported that mtROS inhibitors do not block the secondary activation step, but rather the initial priming step^[199]. Apart from acting as an activator of NLRP3, mitochondria can act as a docking system for inflammasome assembly. This interaction is driven by the externalization of mitochondrial lipid cardiolipin from the inner membrane to the outer membrane, which then independently interacts with caspase-1 and NLRP3^[200].

NLRP3 inflammasome activation in microglia has gained attention as a contributing mechanism in several neuroinflammatory disease pathologies including Alzheimer’s disease, amyotrophic lateral sclerosis, multiple sclerosis, and Parkinson’s disease^[201-205]. While much is similar between the biochemistries of microglia and macrophages, differences in inflammasome activation have been reported. For example, exposure of microglia cells to the antioxidant NAC did not affect LPS priming yet inhibited A β 1-42 peptide stimulation of caspase-1 dependent IL-1 β secretion^[202]. While microglia show similar expression

of inflammasome components and response to stimulus, the dependency on caspase-1 for IL-1 β secretion is only partial and a higher level of mature IL-1 β secretion is observed with longer periods of priming than in hematopoietic macrophages. Prolonged IL-1 β secretion from microglia likely occurs as a result of a deficit in negative regulation mechanisms as compared to macrophages. As an additional consideration, activation of the inflammasome in peripheral macrophages serves in a regulatory capacity in the induction of pyroptotic cell death to remove the damaging immune cell. How this translates to microglia remains in question given the long-lived nature of these cells.

CONCLUSION

The interest in metabolic functions of microglia has evolved from our knowledge of mitochondrial demands and responses of peripheral macrophages in their various effector functions. Recent findings have set the framework for an association between the metabolic status of immune cells with the characteristics of an immune response against pathogens. The majority of studies have relied on different pro-inflammatory stimuli, such as virus, GM-CSF, LPS, LPS + INF γ , or IL-4, to induce cells to examine macrophage metabolism *in vitro*. However, the resulting phenotype and metabolic profiles can differ with not all stimuli leading towards glycolysis. Conversely, the diverse non-inflammatory stimuli normally examined, i.e., IL-10, glucocorticoids, IL-13, M-CSF, and IL-4, are grouped together^[74]; however, similar to the pro-inflammatory stimuli, the phenotypic change may differ. This is not unexpected given that macrophage activation states display multiple profiles depending on the initiating stimuli^[206]. However, examination of metabolic adaptations of macrophages has demonstrated that such adaptations are critical factors regulating a variety of immune cell responses. The need to rapidly modulate cellular responses to pathogen or inflammatory signals demands a remodeling of the metabolic pathways to execute such actions. While many of the basic responses translate across peripheral macrophages and microglia, the uniqueness of microglia suggests that this may not be a complete translation across cells. Additionally, the limited range of inducing stimuli examined in microglia cells raises the question of how the cells will compare given a broader range of stimuli. Further exploration of similarities and uniqueness will contribute to our understanding of the interplay between metabolism and immune cell responses as they apply to the nervous system. It may also offer a framework from which to address issues of translation between experimental animal data to human disease conditions that involve the innate immune system^[207-209]. Understanding the mitochondrial-related characteristics of microglia will likely be critical in identifying successful therapeutic approaches to the detrimental effects of neuroinflammation or in facilitating repair.

DECLARATIONS

Acknowledgments

The authors acknowledge Drs. Christopher McPherson, Christian Lefebvre d'Hellencourt, and Negin Martin for providing reviewer comments on the manuscript.

Authors' contributions

Contributed to the conceptualization, design and interpretation of the experiments and in manuscript preparation: Childers G

Generated the original data: Lopez Hernandez I, Giridharan S

Contributed to the conceptualization of the manuscript, data interpretation, and writing of the final manuscript: Harry GJ

Availability of data and materials

Not applicable.

Financial support and sponsorship

The research was supported by NIH intramural research funding ES021164.

Conflicts of interest

All authors declared that there are no conflicts of interest.

Ethical approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Copyright

© The Author(s) 2020.

REFERENCES

- Krysko DV, Agostinis P, Krysko O, Garg AD, Bachert C, et al. Emerging role of damage-associated molecular patterns derived from mitochondria in inflammation. *Trends Immunol* 2011;32:157-64.
- Boyapati RK, Tamborska A, Dorward DA, Ho GT. Advances in the understanding of mitochondrial DNA as a pathogenic factor in inflammatory diseases. *F1000Res* 2017;6:169.
- Shimada K, Crother TR, Karlin J, Dagvadorj J, Chiba N, et al. Oxidized mitochondrial DNA activated the NLRP3 inflammasome during apoptosis. *Immunity* 2012;36:401-14.
- O'Neill LAJ, Kishton RJ, Rathmell J. A guide to immunometabolism for immunologists. *Nat Rev Immunol* 2016;16:553-65.
- Russell DG, Huang L, VanderVen BC. Immunometabolism at the interface between macrophages and pathogens. *Nat Rev Immunol* 2019;19:291-304.
- Lynch MA. Can the emerging field of immunometabolism provide insights into neuroinflammation? *Prog Neurobiol* 2020;184:101719.
- Ghosh M, Xu Y, Pearce DD. Cyclic AMP is a key regulator of M1 to M2a phenotypic conversion of microglia in the presence of Th2 cytokines. *J Neuroinflammation* 2016;13:9.
- Butovsky O, Jedrychowski MP, Moore CS, Cialic R, Lanser AJ, et al. Identification of a unique TGF-beta-dependent molecular and functional signature in microglia. *Nat Neurosci* 2014;17:131-43.
- Durafour BA, Moore CS, Zammit DA, Johnson TA, Zaguia F, et al. Comparison of polarization properties of human adult microglia and blood-derived macrophages. *Glia* 2012;60:717-27.
- Finsen B, Owens T. Innate immune responses in central nervous system inflammation. *FEBS Lett* 2011;585:3806-12.
- Ginhoux F, Greter M, Leboeuf M, Nandi S, See P, et al. Fate mapping analysis reveals that adult microglia derive from primitive macrophages. *Science* 2010;330:841-5.
- Hickman SE, Kingery ND, Ohsumi TK, Borowsky ML, Wang LC, et al. The microglial sensome revealed by direct RNA sequencing. *Nat Neurosci* 2013;16:1896-905.
- Holtman IR, Skola D, Glass CK. Transcriptional control of microglia phenotypes in health and disease. *J Clin Invest* 2017;127:3220-9.
- Ransohoff RM, Brown MA. Innate immunity in the central nervous system. *J Clin Invest* 2012;122:1164-71.
- Crotti A, Ransohoff RM. Microglial physiology and pathophysiology: insights from genome-wide transcriptional profiling. *Immunity* 2016;44:505-15.
- Borst K, Schwabenland M, Prinz M. Microglia metabolism in health and disease. *Neurochem Int* 2019;130:104331.
- Hawkins BT, Davis TP. The blood-brain barrier/neurovascular unit in health and disease. *Pharmacol Rev* 2005;57:173-85.
- Hermann DM, El-Ali A. The abluminal endothelial membrane in neurovascular remodeling in health and disease. *Sci Signal* 2012;5:re4.
- Ifergan I, Kébir H, Bernard M, Wosik K, Dodelet-Devillers A, et al. The blood-brain barrier induces differentiation of migrating monocytes into Th17-polarizing dendritic cells. *Brain* 2008;131:785-99.
- Muldoon LL, Alvarez JI, Begley DJ, Boado RJ, Del Zoppo GJ, et al. Immunologic privilege in the central nervous system and the blood-brain barrier. *Cereb Blood Flow Metab* 2013;33:13-21.
- Banerjee S, Bhat MA. Neuron-glia interactions in blood-brain barrier formation. *Annu Rev Neurosci* 2007;30:235-58.
- King IL, Dickendesher TL, Segal BM. Circulating Ly-6C⁺ myeloid precursors migrate to the CNS and play a pathogenic role during autoimmune demyelinating disease. *Blood* 2009;113:3190-7.
- Mildner A, Schmidt H, Nitsche M, Merkler D, Hanisch UK, et al. Microglia in the adult brain arise from Ly-6ChiCCR2⁺ monocytes only under defined host conditions. *Nat Neurosci* 2007;12:1544-53.
- Vallieres L, Sawchenko PE. Bone marrow-derived cells that populate the adult mouse brain preserve their hematopoietic identity. *J Neurosci* 2003;23:5197-207.
- Trifilo MJ, Montalto-Morrison C, Stiles LN, Hurst KR, Hardison JL, et al. CXC chemokine ligand 10 controls viral infection in the central nervous system: evidence for a role in innate immune response through recruitment and activation of natural killer cells. *J Virol* 2003;78:585-94.
- González H, Pacheco R. T-cell-mediated regulation of neuroinflammation involved in neurodegenerative diseases. *J Neuroinflam* 2014;11:201.
- Li JI, Gran B, Zhang GX, Ventura ES, Siglienti I, et al. Differential expression and regulation of IL-23 and IL-12 subunits and

- receptors in adult mouse microglia. *J Neurol Sci* 2003;215:95-103.
28. Romero-Suárez S, Del Rio Serrato A, Bueno RJ, Brunotte-Strecker D, Stehle C, et al. The central nervous system contains ILC1s that differ from NK cells in the response to inflammation. *Front Immunol* 2019;10:2337.
 29. Ginhoux F, Jung S. Monocytes and macrophages: developmental pathways and tissue homeostasis. *Nat Rev Immunol* 2014 14:392-404.
 30. Yamasaki R, Lu H, Butovsky O, Ohno N, Rietsch AM, et al. Differential roles of microglia and monocytes in the inflamed central nervous system. *J Exp Med* 2014;211:1533-49.
 31. Davalos D, Grutzendler J, Yang G, Kim JV, Zuo Y, et al. ATP mediates rapid microglial response to local brain injury in vivo. *Nat Neurosci* 2005;8:752-8.
 32. Nimmerjahn A, Kirchhoff F, Helmchen F. Resting microglial cells are highly dynamic surveillants of brain parenchyma in vivo. *Science* 2005;308:1314-8.
 33. Chiu IM, Morimoto ET, Goodarzi H, Liao JT, O'Keeffe S, et al. A neurodegeneration-specific gene-expression signature of acutely isolated microglia from an amyotrophic lateral sclerosis mouse model. *Cell Rep* 2013;4:385-401.
 34. Angata T, Kerr SC, Greaves DR, Varki NM, Crocker PR, et al. Cloning and characterization of human Siglec-11. A recently evolved signaling molecule that can interact with SHP-1 and SHP-2 and is expressed by tissue macrophages, including brain microglia. *J Biol Chem* 2002;277:24466-74.
 35. Takahashi K, Rochford CD, Neumann H. Clearance of apoptotic neurons without inflammation by microglial triggering receptor expressed on myeloid cells-2. *J Exp Med* 2005;201:647-57.
 36. Kawai T, Akira S. The role of pattern-recognition receptors in innate immunity: update on Toll-like receptors. *Nat Immunol* 2010;11:373-84.
 37. Wilkinson K, El Khoury J. Microglial scavenger receptors and their roles in the pathogenesis of Alzheimer's disease. *Int J Alzheimers Dis* 2012;2012:489456.
 38. Pei Z, Pang H, Qian L, Yang S, Wang T, et al. MAC1 mediates LPS-induced production of superoxide by microglia: the role of pattern recognition receptors in dopaminergic neurotoxicity. *Glia* 2007;55:1362-73.
 39. Chen GY, Nuñez G. Sterile inflammation: sensing and reacting to damage. *Nat Rev Immunol* 2010;10:826-37.
 40. Shechter R, Schwartz M. CNS sterile injury: just another wound healing? *Trends Mol Med* 2013;19:135-43.
 41. McPherson CA, Merrick BA, Harry GJ. In vivo molecular markers for pro-inflammatory cytokine M1 stage and resident microglia in trimethyltin-induced hippocampal injury. *Neurotox Res* 2014;25:45-56.
 42. Kono H, Rock KL. How dying cells alert the immune system to danger. *Nat Rev Immunol* 2008;8:279-89.
 43. Takeuchi O, Akira S. Pattern recognition receptors and inflammation. *Cell* 2010;140:805-20.
 44. Biber K, Neumann H, Inoue K, Boddeke HW. Neuronal 'On' and 'Off' signals control microglia. *Trends Neurosci* 2007;30:596-602.
 45. Hanisch UK, Kettenmann H. Microglia: active sensor and versatile effector cells in the normal and pathologic brain. *Nat Neurosci* 2007;10:1387-94.
 46. Ransohoff RM, Cardona AE. The myeloid cells of the central nervous system parenchyma. *Nature* 2010;468:253-62.
 47. Kettenmann H, Hanisch UK, Noda M, Verkhratsky A. Physiology of microglia. *Physiol Rev* 2011;91:461-553.
 48. Domercq M, Vázquez-Villoldo N, Matute C. Neurotransmitter signaling in the pathophysiology of microglia. *Front Cell Neurosci* 2013;7:49.
 49. Kierdorf K, Prinz M. Factors regulating microglia activation. *Front Cell Neurosci* 2013;7:44.
 50. Pocock JM, Kettenmann H. Neurotransmitter receptors on microglia. *Trends Neurosci* 2007;30:527-35.
 51. Garcia-Revilla J, Alonso-Bellido IM, Burguillos MA, Herrera AJ, Espinosa-Oliva AM. Reformulating pro-oxidant microglia in neurodegeneration. *J Clin Med* 2019;8:1719.
 52. Lee JK, Chung J, Kannarkat GT, Tansey MG. Critical role of regulator G-protein signaling 10 (RGS10) in modulating macrophage M1/M2 activation. *PLoS One* 2013;8:e81785.
 53. Seifert S, Pannell M, Uckert W, Färber K, Kettenmann H. Transmitter- and hormone-activated Ca(2+) responses in adult microglia/brain macrophages in situ recorded after viral transduction of a recombinant Ca(2+) sensor. *Cell Calcium* 2011;49:365-75.
 54. Ajami B, Samusik N, Wieghofer P, Ho PP, Crotti A, et al. Single-cell mass cytometry reveals distinct populations of brain myeloid cells in mouse neuroinflammation and neurodegeneration models. *Nat Neurosci* 2018;21:541-51.
 55. Pettersen JS, Fuentes-Duculan J, Suárez-Fariñas M, Pierson KC, Pitts-Kiefer A, et al. Tumor-associated macrophages in the cutaneous SCC microenvironment are heterogeneously activated. *J Invest Dermatol* 2011;131:1322-30.
 56. Ziegler-Heitbrock L, Hofer TPJ. Toward a refined definition of monocyte subsets. *Front Immunol* 2013;4:23.
 57. Mosser DM, Edwards JP. Exploring the full spectrum of macrophage activation. *Nat Rev Immunol* 2008;8:958-69.
 58. Gordon S. Alternative activation of macrophages. *Nat Rev Immunol* 2003;3:23-35.
 59. Lynch MA. The multifaceted profile of activated microglia. *Mol Neurobiol* 2009;40:139-56.
 60. Biswas SK, Mantovani A. Macrophage plasticity and interaction with lymphocyte subsets: cancer as a paradigm. *Nat Immunol* 2010;11:889-96.
 61. Sica A, Mantovani A. Macrophage plasticity and polarization: in vivo veritas. *J Clin Invest* 2012;122:787-95.
 62. Hume DA, Freeman TC. Transcriptomic analysis of mononuclear phagocyte differentiation and activation. *Immunol Rev* 2014;262:74-84.
 63. Colton CA. Heterogeneity of microglial activation in the innate immune response in the brain. *J Neuroimmune Pharmacol* 2009;4:399-418.
 64. Michelucci A, Heurtaux T, Grandbarbe L, Morga E, Heuschling P. Characterization of the microglial phenotype under specific pro-inflammatory and anti-inflammatory conditions: effects of oligomeric and fibrillar amyloid-beta. *J Neuroimmunol* 2009;210:3-12.
 65. Boche D, Perry VH, Nicoll JA. Review: activation patterns of microglia and their identification in the human brain. *Neuropathol Appl Neurobiol* 2013;39:3-18.
 66. Cherry JD, Olschowka JA, O'Banion MK. Neuroinflammation and M2 microglia: the good, the bad, and the inflamed. *J Neuroinflammation* 2014;11:98.

67. Murray PJ, Allen JE, Biswas SK, Fisher EA, Gilroy DW, et al. Macrophage activation and polarization: nomenclature and experimental guidelines. *Immunity* 2014;41:14-20.
68. Loane D, Kumar A. Microglia in the TBI Brain: the good, the bad, And the dysregulated. *Exp Neurol* 2016;275: 316-27.
69. Orihuea R, McPherson CA, Harry GJ. Microglial M1/M2 polarization and metabolic states *Br J Pharmacol* 2016;173:649-65.
70. Gerrick KY, Gerrick ER, Gupta A, Wheelan SJ, Yegnasubramanian S, et al. Transcriptional profiling identifies novel regulators of macrophage polarization. *PLoS One* 2018;13:e0208602.
71. Ransohoff RM. A polarizing question: do M1 and M2 microglia exist? *Nat Neurosci* 2016;19:987-91.
72. Stein M. Interleukin 4 potentially enhances murine macrophage mannose receptor activity: a marker of alternative immunologic macrophage activation. *J Exp Med* 1992;176:287-92.
73. Doyle AG, Herbein G, Montaner LJ, Minty AJ, Caput D, et al. Interleukin-13 alters the activation state of murine macrophages in vitro: comparison with interleukin-4 and interferon-gamma. *Eur J Immunol* 1994;24:1441-5.
74. Martinez FO, Gordon S. The M1 and M2 paradigm of macrophage activation: time for reassessment. *F1000Prime Rep* 2014;6:1-13.
75. Mills EL, O'Neill LA. Reprogramming mitochondrial metabolism in macrophages as an anti-inflammatory signal. *Eur J Immunol* 2016;46:13-21.
76. Murray PJ. Understanding and exploiting the endogenous interleukin-10/STAT3-mediated anti-inflammatory response. *Curr Opin Pharmacol* 2006;6:379-86.
77. Bordt EA, Polster BM. NADPH oxidase- and mitochondria-derived reactive oxygen species in proinflammatory microglial activation: a bipartisan affair? *Free Radic Biol Med* 2014;76:34-46.
78. Dashty M. A quick look at biochemistry: carbohydrate metabolism. *Clin Biochem* 2013;46:1339-52.
79. Liemburg-Apers DC, Schirris TJ, Russel FG, Willems PH, Koopman WJ. Mitochondrial dysfunction triggers a rapid compensatory increase in steady-state glucose flux. *Biophys J* 2015;109:1372-86.
80. Vazquez A, Liu J, Zhou Y, Oltvai ZN. Catabolic efficiency of aerobic glycolysis: the Warburg effect revisited. *BMC Syst Biol* 2010;4:58.
81. Warburg O. On respiratory impairment in cancer cells. *Science* 1956;124:269-70.
82. Vander Heiden MG, Cantley LC, Thompson CB. Understanding the Warburg effect: the metabolic requirements of cell proliferation. *Science* 2009;324:1029-33.
83. Weinberg SE, Sena LA, Chandel NS. Mitochondria in the regulation of innate and adaptive immunity. *Immunity* 2015;42:406-17.
84. Kelly B, O'Neill LA. Metabolic reprogramming in macrophages and dendritic cells in innate immunity. *Cell Res* 2015;25:771-84.
85. Mehta MM, Weinberg SE, Chandel NS. Mitochondrial control of immunity: beyond ATP. *Nat Rev Immunol* 2017;17:608-20.
86. Viola A, Munari F, Sanchez-Rodriguez R, Scolaro T, Castegna A. The metabolic signature of macrophage responses. *Front Immunol* 2019;10:1462.
87. Zuo H, Wan Y. Metabolic reprogramming in mitochondria of myeloid cells. *Cells* 2020;9:5.
88. Biswas SK, Mantovani A. Orchestration of metabolism by macrophages. *Cell Metab* 2012;15:432-7.
89. Ryan DG, O'Neill LAJ. Krebs cycle reborn in macrophage immunometabolism. *Annu Rev Immunol* 2020; Epub ahead of print. doi: 10.1146/annurev-immunol-081619-104850
90. Bernhart E, Kollroser M, Rechberger G, Reicher H, Heinemann A. Lysophosphatidic acid receptor activation affects the C13NJ microglia cell line proteome leading to alterations in glycolysis, motility, and cytoskeletal architecture. *Proteomics* 2010;10:141-58.
91. Moss DW, Bates TE. Activation of murine microglial cell lines by lipopolysaccharide and interferon-gamma causes NO-mediated decreases in mitochondrial and cellular function. *Eur J Neurosci* 2001;13:529-38.
92. Zuo H, Wan Y. Metabolic reprogramming in mitochondria of myeloid cells. *Cell* 2020;9:5.
93. Lartigue L, Faustin B. Mitochondria: Metabolic regulators of innate immune responses to pathogens and cell stress. *Int J Biochem Cell Biol* 2013;45:2052-56.
94. Voloboueva LA, Emery JF, Sun X, Giffard RG. Inflammatory response of microglial BV-2 cells includes a glycolytic shift and is modulated by mitochondrial glucose-regulated protein 75/mortalin. *FEBS Lett* 2013;587:756-62.
95. Gimeno-Bayon J, Lopez-Lopez A, Rodriguez MJ, Mahy N. Glucose pathways adaptation supports acquisition of activated microglia phenotype. *J Neurosci Res* 2014;92:723-31.
96. Chenais B, Morjani H, Drapier JC. Impact of endogenous nitric oxide on microglial cell energy metabolism and labile iron pool. *J Neurochem* 2002;81:615-23.
97. Rodríguez-Prados JC, Través PG, Cuenca J, Rico D, Aragonés J, et al. Substrate fate in activated macrophages: a comparison between innate, classic, and alternative activation. *J Immunol* 2010;185:605-14.
98. Odegaard JI, Chawla A. Alternative macrophage activation and metabolism. *Annu Rev Pathol* 2011;6:275-97.
99. Mantovani A, Biswas SK, Galdiero MR, Sica A, Locati M. Macrophage plasticity and polarization in tissue repair and remodelling. *J Pathol* 2013;229:176-85.
100. Galvan-Pena S, O'Neill LA. Metabolic reprogramming in macrophage polarization. *Front Immunol* 2014;5:420.
101. Mills EL, Kelly B, O'Neill LAJ. Mitochondria are the powerhouses of immunity. *Nat Immunol* 2017;18:488-98.
102. Newsholme P, Curi R, Gordon S, Newsholme EA. Metabolism of glucose, glutamine, long-chain fatty acids and ketone bodies by murine macrophages. *Biochem J* 1986;239:121-5.
103. Newsholme P, Gordon S, Newsholme EA. Rates of utilization and fates of glucose, glutamine, pyruvate, fatty acids and ketone bodies by mouse macrophages. *Biochem J* 1987;242:631-6.
104. Haschemi A, Kosma P, Gille L, Evans CR, Burant CF, et al. The sedoheptulose kinase CARKL directs macrophage polarization through control of glucose metabolism. *Cell Metab* 2012;15:813-26.
105. Blagih J, Jones RG. Polarizing macrophages through reprogramming of glucose metabolism. *Cell Metab* 2012;15:793-5.
106. Bonora M, Patergnani S, Rimessi A, De Marchi E, Suski JM, et al. ATP synthesis and storage. *Purinergic Signal* 2012;8:343-57.
107. Bernier LP, Bohlen CJ, York EM, Choi HB, Kamyabi A, et al. Nanoscale surveillance of the brain by microglia via cAMP-regulated

- filopodia. *Cell Reports* 2019;27:2895-908.
108. Madry C, Kyrargyri V, Arancibia-Carcamo IL, Jolivet R, Kohsaka S, et al. Microglial ramification, surveillance, and interleukin-1 β release are regulated by the two-pore domain K(+) channel THIK-1. *Neuron* 2018;97:299-312.
109. Ghosh S, Castillo E, Frias ES, Swanson RA. Bioenergetic regulation of microglia. *Glia* 2018;66:1200-12.
110. Cherry JD, Olschowka JA, O'Banion MK. Are 'resting' microglia more 'm2'? *Front Immunol* 2014;5:594.
111. Baik SH, Kang S, Lee W, Choi H, Chung S, et al. A breakdown in metabolic reprogramming causes microglia dysfunction in Alzheimer's disease. *Cell Metab* 2019;30:493-507.e6.
112. Holland R, McIntosh AL, Finucane OM, Mela V, Rubio-Araiz A, et al. Inflammatory microglia are glycolytic and iron retentive and typify the microglia in APP/PS1 mice. *Brain Behav Immun* 2017;68:183-96.
113. McIntosh A, Mela V, Harty C, Minogue AM, Costello DA, et al. Iron accumulation in microglia triggers a cascade of events that leads to altered metabolism and compromised function in APP/PS1 mice. *Brain Pathol* 2019;29:606-21.
114. Wang T, Liu H, Lian G, Zhang SY, Wang X, et al. HIF1 α -induced glycolysis metabolism is essential to the activation of inflammatory macrophages. *Mediators Inflamm* 2017;2017:1-10.
115. Nair S, Sobotka KS, Joshi P, Gressens P, Fleiss B, et al. Lipopolysaccharide-induced alteration of mitochondrial morphology induces a metabolic shift in microglia modulating the inflammatory response in vitro and in vivo. *Glia* 2019;67:1047-61.
116. Chénais B, Morjani H, Drapier JC. Impact of endogenous nitric oxide on microglial cell energy metabolism and labile iron pool. *J Neurochem* 2002;81:615-23.
117. Banati RB, Egensperger R, Maassen A, Hager G, Kreutzberg GW, et al. Mitochondria in activated microglia in vitro. *J Neurocytol* 2004;33:535-41.
118. Ferger AI, Campanelli L, Reimer V, Muth KN, Merdian I, et al. Effects of mitochondrial dysfunction on the immunological properties of microglia. *J Neuroinflamm* 2010;7:45.
119. Finucane OM, Sugrue J, Rubio-Araiz A, Guillot-Sestier MV, Lynch MA. The NLRP3 inflammasome modulates glycolysis by increasing PFKFB3 in an IL-1 β dependent manner in macrophages. *Sci Rep* 2019;9:4034.
120. Ip WKE, Hoshi N, Shouval DS, Snapper S, Medzhitov R. Anti-inflammatory effect of IL-10 mediated by metabolic reprogramming of macrophages. *Science* 2017;356:513-9.
121. Murray PJ, Wynn TA. Obstacles and opportunities for understanding macrophage polarization. *J Leukoc Biol* 2011;89:557-63.
122. Biswas SK, Chittiezath M, Shalova IN, Lim JY. Macrophage polarization and plasticity in health and disease. *Immunol Res* 2012;53:11-24.
123. Huang SC, Everts B, Ivanova Y, O'Sullivan D, Nascimento M, et al. Cell-intrinsic lysosomal lipolysis is essential for alternative activation of macrophages. *Nat Immunol* 2014;15:846-55.
124. Liu PS, Wang H, Li X, Chao T, Teav T, et al. Alpha-ketoglutarate orchestrates macrophage activation through metabolic and epigenetic reprogramming. *Nat Immunol* 2017;18:985-94.
125. Vats D, Mukundan L, Odegaard JI, Zhang L, Smith KL, et al. Oxidative metabolism and PGC-1 β attenuate macrophage-mediated inflammation. *Cell Metab* 2006;4:13-24.
126. Mills CD, Kincaid K, Alt JM, Hellman MJ, Hill AM. M-1/M-2 macrophages and the Th1/Th2 paradigm. *J Immunol* 2000;164:6166-73.
127. De Simone R, Ajmone-Cat MA, Pandolfi M, Bernardo A, De Nuccio C, et al. The mitochondrial uncoupling protein-2 is a master regulator of both M1 and M2 microglial responses. *J Neurochem* 2015;135:147-56.
128. West AP, Brodsky IE, Rahner C, Woo DK, Erdjument-Bromage H, et al. TLR signalling augments macrophage bactericidal activity through mitochondrial ROS. *Nature* 2011;472:476-80.
129. Van den Bossche J, Baardman J, Otto NA, van der Velden S, Neele AE. Mitochondrial dysfunction prevents repolarization of inflammatory macrophages. *Cell Rep* 2016;17:684-96.
130. Poderoso JJ, Helfenberger K, Poderoso C. The effect of nitric oxide on mitochondrial respiration. *Nitric Oxide* 2019;88:61-72.
131. Hu Y, Mai W, Chen L, Cao K, Zhang B, et al. mTOR-mediated metabolic reprogramming shapes distinct microglia functions in response to lipopolysaccharide and ATP. *Glia* 2020;68:1031-45.
132. Li D, Wang C, Yao Y, Chen L, Liu G, et al. mTORC1 pathway disruption ameliorates brain inflammation following stroke via a shift in microglia phenotype from M1 type to M2 type. *FASEB J* 2016;30:3388-99.
133. Ulland TK, Song WM, Huang SC, Ulrich JD, Sergushichev A, et al. TREM2 maintains microglial metabolic fitness in alzheimer's disease. *Cell* 2017;170:649-63.e13.
134. Zhao X, Liao Y, Morgan S, Mathur R, Feustel P, et al. Noninflammatory changes of microglia are sufficient to cause epilepsy. *Cell Rep* 2018;22:2080-93.
135. Bedard K, Krause KH. The NOX family of ROS-generating NADPH oxidases: physiology and pathophysiology. *Physiol Rev* 2007;87:245-313.
136. Pawate S, Shen Q, Fan F, Bhat NR. Redox regulation of glial inflammatory response to lipopolysaccharide and interferon gamma. *J Neurosci Res* 2004;77:540-51.
137. Qin L, Liu Y, Wang T, Wei SJ, Block ML, et al. NADPH oxidase mediates lipopolysaccharide-induced neurotoxicity and proinflammatory gene expression in activated microglia. *J Biol Chem* 2004;279:1415-21.
138. Loane DJ, Stoica BA, Pajoohesh-Ganji A, Byrnes KR, Faden AI. Activation of metabotropic glutamate receptor 5 modulates microglial reactivity and neurotoxicity by inhibiting NADPH oxidase. *J Biol Chem* 2009;284:15629-39.
139. Choi SH, Aid S, Kim HW, Jackson SH, Bosetti F. Inhibition of NADPH oxidase promotes alternative and anti-inflammatory microglial activation during neuroinflammation. *J Neurochem* 2012;120:292-301.
140. Eguchi H, Fujiwara N, Sakiyama H, Yoshihara D, Suzuki K. Hydrogen peroxide enhances LPS-induced nitric oxide production via the expression of interferon beta in BV-2 microglial cells. *Neurosci Lett* 2011;494:29-33.

141. Mander PK, Jekabsone A, Brown GC. Microglia proliferation is regulated by hydrogen peroxide from NADPH oxidase. *J Immunol* 2006;176:1046-52.
142. Roy A, Jana A, Yatish K, Freidt MB, Fung YK, et al. Reactive oxygen species up-regulate CD11b in microglia via nitric oxide: implications for neurodegenerative diseases. *Free Radic Biol Med* 2008;45:686-99.
143. Taetzsch T, Levesque S, McGraw C, Brookins S, Luqa R, et al. Redox Regulation of NF- κ B p50 and M1 Polarization in Microglia. *Glia* 2015;63:423-40.
144. Yuan YH, Sun JD, Wu MM, Hu JF, Peng SY, et al. Rotenone could activate microglia through NF- κ B associated pathway. *Neurochem Res* 2013;38:1553-60.
145. Kletzien RF, Harris PK, Foellmi LA. Glucose-6-phosphate dehydrogenase: a “housekeeping” enzyme subject to tissue-specific regulation by hormones, nutrients, and oxidant stress. *FASEB J* 1994;8:174-81.
146. Salvemini F, Franzé A, Iervolino A, Filosa S, Salzano S, et al. Enhanced glutathione levels and oxidoresistance mediated by increased glucose-6-phosphate dehydrogenase expression. *J Biol Chem* 1999;274:2750-7.
147. Mustacich D, Powis G. Thioredoxin reductase. *Biochem J* 2000;346:1-8.
148. Ying W. NAD⁺/NADH and NADP⁺/NADPH in cellular functions and cell death: regulation and biological consequences. *Antioxid Redox Signal* 2008;10:179-206.
149. Shen Y, Kapfhamer D, Minnella AM, Kim JE, Won SJ, et al. Bioenergetic state regulates innate inflammatory responses through the transcriptional co-repressor CtBP. *Nat Commun* 2017;8:624.
150. Wang L, Pavolu S, Du X, Bhuckory M, Xu H, et al. Glucose transporter 1 critically controls microglial activation through facilitating glycolysis. *Mol Neurodegener* 2019;14:2.
151. Kalsbeek MJ, Mulder L, Yi CX. Microglia energy metabolism in metabolic disorder. *Mol Cell Endocrinol* 2016;438:27-35.
152. Payne J, Maher F, Simpson I, Mattice L, Davies P. Glucose transporter Glut 5 expression in microglial cells. *Glia* 1997;21:327-31.
153. Hsieh C, Liu C, Lee C, Yu LE, Wang JY. Acute glucose fluctuation impacts microglial activity, leading to inflammatory activation or self-degradation. *Sci Rep* 2019;9:840.
154. Chen J, Sun Z, Jin M, Tu Y, Wang S, et al. Inhibition of AGEs/RAGE/Rho/ROCK pathway suppresses non-specific neuroinflammation by regulating BV2 microglial M1/M2 polarization through the NF- κ B pathway. *J Neuroimmunol* 2017;305:108-14.
155. Tannahill GM, Curtis AM, Adamik J, Palsson-McDermott EM, McGettrick AF, et al. Succinate is an inflammatory signal that induces IL-1 β through HIF-1 α . *Nature* 2013;496:238-42.
156. O'Neill LAJ, Pearce EJ. Immunometabolism governs dendritic cell and macrophage function. *J Exp Med* 2016;213:15-23.
157. Wang F, Zhang S, Jeon R, Vuckovic, Jiang X, et al. Interferon gamma induces reversible metabolic reprogramming of M1 macrophages to sustain cell viability and pro-inflammatory activity. *EbioMedicine* 2018;30:303-16.
158. Choi SJ, Shin IJ, Je KH, Min EK, Kim EJ, et al. Hypoxia antagonizes glucose deprivation on interleukin 6 expression in an Akt dependent, but HIF-1/2 α independent manner. *PLoS One* 2013;8:e58662.
159. Churchward MA, Tchir DR, Todd KG. Microglial function during glucose deprivation: inflammatory and neuropsychiatric implications. *Mol Neurobiol* 2018;55:1477-87.
160. Weng F, Zhang S, Vuckovic I, Ryoungchoon J, Lerman A, et al. Glycolytic stimulation is not a requirement for M2 macrophage differentiation. *Cell Metab* 2018;28:463-75.
161. Venter G, Oerlemans FTJJ, Wijers M, Willemse M, Fransen JAM, et al. Glucose controls morphodynamics of LPS-stimulated macrophages. *PLoS One* 2014;9:e96786.
162. Meiser J, Krämer L, Sapcariu SC, Battello N, Ghelfi J, et al. Pro-inflammatory macrophages sustain pyruvate oxidation through pyruvate dehydrogenase for the synthesis of itaconate and to enable cytokine expression. *J Biol Chem* 2016;291:3932-46.
163. Shay JE, Celeste-Simon M. Hypoxia-inducible factors: crosstalk between inflammation and metabolism. *Semin Cell Dev Biol* 2012;23:389-94.
164. Liu PS, Wang H, Li X, Chao T, Teav T, et al. α -ketoglutarate orchestrates macrophage activation through metabolic and epigenetic reprogramming. *Nat Immunol* 2017;8:985-94.
165. Moon JS, Hisata S, Park MA, DeNicola GM, Ryter SW, et al. mTORC1-induced HK1-dependent glycolysis regulates NLRP3 inflammasome activation. *Cell Rep* 2015;12:102-15.
166. Xie M, Yu Y, Kang R, Zhu S, Yang L, et al. PKM2-dependent glycolysis promotes NLRP3 and AIM2 inflammasome activation. *Nat Commun* 2016;7:13280.
167. Wen H, Miao EA, Ting JP. Mechanisms of NOD-like receptor-associated inflammasome activation. *Immunity* 2013;39:432-41.
168. Adamczak SE, de Rivero Vaccari JP, Dale G, Brand FJ, Nonner D, et al. Pyroptotic neuronal cell death mediated by the AIM2 inflammasome. *J Cereb Blood Flow Metab* 2014;34:621-29.
169. Kaushal V, Dye R, Pakavathkumar P. Neuronal NLRP1 inflammasome activation of Caspase-1 coordinately regulates inflammatory interleukin-1-beta production and axonal degeneration-associated Caspase-6 activation. *Cell Death Differ* 2015;22:1676-86.
170. Yogarajah T, Ong KC, Perera D, Wong KT. AIM2 inflammasome-mediated pyroptosis in enterovirus A71-infected neuronal cells restricts viral replication. *Sci Rep* 2017;7:5845.
171. Minkiewicz J, de Rivero Vaccari JP, Keane RW. Human astrocytes express a novel NLRP2 inflammasome. *Glia* 2013;87:3655-67.
172. Freeman L, Guo H, David CN, Brickey WJ, Jha S, et al. NLR members NLRC4 and NLRP3 mediate sterile inflammasome activation in microglia and astrocytes. *J Exp Med* 2017;214:1351-70.
173. de Rivero Vaccari JP, Dietrich WD, Keane RW. Activation and regulation of cellular inflammasomes: gaps in our knowledge for central nervous system injury. *J Cereb Blood Flow Metab* 2014;34:369-75.
174. Burm SM, Zuiderwijk-Sick EA, 't Jong AE, van der Putten C, Veth J, et al. Inflammasome-induced IL-1 β secretion in microglia is characterized by delayed kinetics and is only partially dependent on inflammatory caspases. *J Neurosci* 2015;35:678-87.
175. Hornung V, Bauernfeind F, Halle A, Samstad EO, Kono H, et al. Silica crystals and aluminum salts activate the NALP3 inflammasome

- through phagosomal destabilization. *Nat Immunol* 2008;9:847-56.
176. Franchi L, Eigenbrod T, Muñoz-Planillo R, Núñez G. The inflammasome: a caspase-1-activation platform that regulates immune responses and disease pathogenesis. *Nat Immunol* 2009;10:241-7.
177. Hanamsagar R, Torres W, Kielian T. Inflammasome activation and IL-1 β /IL-18 processing are influenced by distinct pathways in microglia. *J Neurochem* 2011;119:736-48.
178. Lunov O, Syrovets T, Loos C, Nienhaus GU, Mailander V, et al. Amino-Functionalized polystyrene nanoparticles activate the NLRP3 inflammasome in human macrophages. *ACS Nano* 2011;5:9648-57.
179. Schnaars M, Beckert H, Halle A. Assessing β -amyloid-induced NLRP3 inflammasome activation in primary microglia. *Methods Mol Biol* 2013;1040:1-8.
180. Hughes MM, O'Neill LAJ. Metabolic regulation of NLRP3. *Immunol Rev* 2018;281:88-98.
181. Heid ME, Keyel PA, Kamga C, Shiva S, Watkins SC, et al. Mitochondrial reactive oxygen species induces NLRP3-dependent lysosomal damage and inflammasome activation. *J Immunol* 2013;191:5230-8.
182. Gustin A, Kirchmeyer M, Koncina E, Felten P, Losciuto S, et al. NLRP3 inflammasome is expressed and functional in mouse brain microglia but not in astrocytes. *PLoS One* 2015;19;10:e0130624.
183. Afonina IS, Zhong Z, Karin M, Beyaert R. Limiting inflammation - the negative regulation of NF- κ B and the NLRP3 inflammasome. *Nat Immunol* 2017;18:861-9.
184. Elliott EI, Sutterwala FS. Initiation and perpetuation of NLRP3 inflammasome activation and assembly. *Immunol Rev* 2015;265:35-52.
185. Broz P, Dixit VM. Inflammasomes: mechanism of assembly, regulation and signalling. *Nat Rev Immunol* 2016;16:407-20.
186. Zambetti LP, Laudisi F, Licandro G, Ricciardi-Castagnoli P, Mortellaro A. The rhapsody of NLRPs: master players of inflammation...and a lot more. *Immunol Res* 2012;53:78-90.
187. Dinarello CA. Immunological and inflammatory functions of the interleukin-1 family. *Annu Rev Immunol* 2009;27:519-50.
188. Evavold CL, Ruan J, Tan Y, Xia S, Wu H, et al. The pore-forming protein gasdermin D regulates interleukin-1 secretion from living macrophages. *Immunity* 2018;48:35-44.e6.
189. Heilig R, Dick MS, Sborgi L, Meunier E, Hiller S, et al. The Gasdermin-D pore acts as a conduit for IL-1 β secretion in mice. *Eur J Immunol* 2018;48:584-92.
190. Lee HM, Kang J, Lee SJ, Jo EK. Microglial activation of the NLRP3 inflammasome by the priming signals derived from macrophages infected with mycobacteria. *Glia* 2013;61:441-52.
191. Viganò E, Mortellaro A. Caspase-11: the driving factor for noncanonical inflammasomes. *Eur J Immunol* 2013;43:2240-5.
192. Franklin BS, Bossaller L, De Nardo D, Ratter JM, Stutz A, et al. The adaptor ASC has extracellular and 'prionoid' activities that propagate inflammation. *Nat Immunol* 2014;15:727-37.
193. Venegas C, Kumar S, Franklin BS, Dierkes T, Brinkschulte R, et al. Microglia-derived ASC specks cross-seed amyloid- β in Alzheimer's disease. *Nature* 2017;552:355-61.
194. Gurung P, Lukens JR, Kanneganti TD. Mitochondria: diversity in the regulation of NLRP3 inflammasome. *Trends Mol Med* 2015;3:193-201.
195. Shimada K, Crother TR, Karlin J, Dagvadorj J, Chiba N, et al. Oxidized mitochondrial DNA activates the NLRP3 inflammasome during apoptosis. *Immunity* 2012;36:401-14.
196. Nakahira, K, Haspel JA, Rathinam VA, Lee SJ, Dolinay T, et al. Autophagy proteins regulate innate immune responses by inhibiting the release of mitochondrial DNA mediated by the NALP3 inflammasome. *Nat Immunol* 2011;12:222-30.
197. Zhong Z, Liang S, Sanchez-Lopez E, He F, Shalapour S, et al. New mitochondrial DNA synthesis enables NLRP3 inflammasome activation. *Nature* 2018;560:198-203.
198. Zhou R, Yazdi AS, Menu P, Tschopp J. A role for mitochondria in NLRP3 inflammasome activation. *Nature* 2011;469:221-5.
199. Bauernfeind F, Bartok E, Rieger A, Franchi L, Núñez G, et al. Cutting edge: reactive oxygen species inhibitors block priming, but not activation, of the NLRP3 inflammasome. *J Immunol* 2011;187:613-7.
200. Elliott EI, Miller A, Banoth B, Iyer SS, Stotland A, et al. Cutting edge: mitochondrial assembly of the NLRP3 inflammasome complex is initiated at priming. *J Immunol* 2018;200:3047-52.
201. Codolo G, Plotegher N, Pozzobon T, Brucale M, Tessari I, et al. Triggering of inflammasome by aggregated α -synuclein, an inflammatory response in synucleinopathies. *PLoS One* 2013;8:e55375.
202. Parajuli B, Sonobe Y, Horiuchi H, Takeuchi H, Mizuno T, et al. Oligomeric amyloid beta induces IL-1 β processing via production of ROS: implication in Alzheimer's disease. *Cell Death Dis* 2013;4:e975.
203. Bellezza I, Grottelli S, Costanzi E, Scarpelli P, Pigna E, et al. Peroxynitrite activates the NLRP3 inflammasome cascade in SOD1(G93A) mouse model of amyotrophic lateral sclerosis. *Mol Neurobiol* 2018;55:2350-61.
204. Venegas C, Heneka MT. Inflammasome-mediated innate immunity in Alzheimer's disease. *FASEB J* 2019;33:13075-84.
205. Deora V, Lee JD, Albornoz EA, McAlary L, Jagaraj CJ, et al. The microglial NLRP3 inflammasome is activated by amyotrophic lateral sclerosis proteins. *Glia* 2020;68:407-21.
206. Xue J, Schmidt SV, Sander J, Draffehn A, Krebs W, et al. Transcriptome-based network analysis reveals a spectrum model of human macrophage activation. *Immunity* 2014;40:274-88.
207. Healy LM, Yaqubi M, Ludwin S, Antel JP. Species differences in immune-mediated CNS tissue injury and repair: a (neuro) inflammatory topic. *Glia* 2020;68:811-29.
208. Stoger JL, Gijbels MJ, van der Velden S, Manca M, van der Loos CM, et al. Distribution of macrophage polarization markers in human atherosclerosis. *Atherosclerosis* 2012;225:461-8.
209. Van den Bossche J, O'Neill LA, Menon D. Macrophage immunometabolism: where are we (going)? *Trends Immunol* 2017;38:395-406.

Original Article

Open Access



Phf15 - a novel transcriptional repressor regulating inflammation in a mouse microglial cell line

Sandra E. Muroy¹, Greg A. Timblin¹, Marcela K. Preininger¹, Paulina Cedillo¹, Kaoru Saijo^{1,2}

¹Department of Molecular and Cell Biology, University of California, Berkeley, CA 94720, USA.

²Helen Wills Neuroscience Institute, University of California, Berkeley, CA 94720, USA.

Correspondence to: Prof. Kaoru Saijo, Department of Molecular and Cell Biology, University of California, Berkeley, 439 Life Sciences Addition, Berkeley, CA 94720, USA. E-mail: ksaijo@berkeley.edu

How to cite this article: Muroy SE, Timblin GA, Preininger MK, Cedillo P, Saijo K. *Phf15* - a novel transcriptional repressor regulating inflammation in mouse microglia cell line. *Neuroimmunol Neuroinflammation* 2020;7:166-82. <http://dx.doi.org/10.20517/2347-8659.2020.16>

Received: 11 Feb 2020 **Accepted:** 14 Feb 2020 **Available online:** 17 Apr 2020

Science Editor: Andis Klegeris **Copy Editor:** Jing-Wen Zhang **Production Editor:** Tian Zhang

Abstract

Aim: Excessive microglial inflammation has emerged as a key player in mediating the effects of aging and neurodegeneration on brain dysfunction. Thus, there is great interest in discovering transcriptional repressors that can control this process. We aimed to examine whether *Phf15* - one of the top differentially expressed genes in microglia during aging in humans - could regulate transcription of proinflammatory mediators in microglia.

Methods: Real-time quantitative PCR was used to assess *Phf15* mRNA expression in mouse brain during aging. Loss-of-function [short hairpin RNA (shRNA) -mediated knockdown (KD) and CRISPR/Cas9-mediated knockout (KO) of *Phf15*] and gain-of-function [retroviral overexpression (OE) of murine *Phf15* cDNA] studies in a murine microglial cell line (SIM-A9) followed by immune activation with lipopolysaccharide were used to determine the effect of *Phf15* on proinflammatory factor (*Tnfα*, *IL-1β*, and *Nos2*) mRNA expression. RNA sequencing was used to determine global transcriptional changes after *Phf15* knockout under basal conditions and after lipopolysaccharide stimulation.

Results: *Phf15* expression increases in mouse brain during aging, similar to humans. KD, KO, and OE studies determined that *Phf15* represses mRNA expression levels of proinflammatory mediators such as *Tnfα*, *IL-1β*, and *Nos2*. Global transcriptional changes after *Phf15* KO showed that *Phf15* specifically represses genes related to the antiviral (type I interferon) response and cytokine production in microglia.

Conclusion: We provide the first evidence that *Phf15* is an important transcriptional repressor of microglial inflammation, regulating the antiviral response and proinflammatory cytokine production. Importantly, *Phf15* regulates both basal and signal-dependent activation and controls the magnitude and duration of the microglial inflammatory response.

Keywords: *Phf15*, microglia, transcriptional repression, neuroinflammation



© The Author(s) 2020. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, sharing, adaptation, distribution and reproduction in any medium or format, for any purpose, even commercially, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.



INTRODUCTION

Microglia are the resident myeloid-lineage cells of the brain. They actively provide homeostatic surveillance of the brain parenchyma, playing critical roles during development, maintenance, and repair throughout the life of an organism. As innate immune cells, however, microglia are also capable of mounting a full inflammatory response to environmental challenge in order to clear threats and restore homeostasis^[1-6]. Microglia express pattern recognition receptors including Toll-like receptors to sense changes in their environment, such as infection by pathogens or endogenous danger signals. They can then respond by releasing proinflammatory mediators such as tumor necrosis factor alpha (TNF α), interleukin 1 beta (IL-1 β), IL-6, reactive oxygen species, and reactive nitrogen species including nitric oxide (NO) to protect against threats^[1,5,7].

Although beneficial when their production is tightly controlled, deregulated or sustained microglial production of inflammatory mediators can lead to collateral damage of surrounding neurons and other cells^[5,7,8]. Thus, the transition to an activated state, as well as timely resolution of the inflammatory response, must be tightly regulated. Increasing evidence suggests that, during aging, microglia lose homeostatic function and acquire a proinflammatory phenotype that exacerbates aging-related brain dysfunction^[9]. Indeed, aberrant microglia activation has been found in many types of age-related neurodegenerative conditions, for example Parkinson's disease (PD) and Alzheimer's disease (AD), which are marked by inflammatory processes involving glia, and microglia in particular^[9-11].

Since excessive production of proinflammatory mediators is neurotoxic^[8,12-14], various molecular mechanisms exist to regulate transcriptional repression of inflammatory gene expression. For example, basal state repression, that is, before the arrival of an activating signal, is generally carried out via recruitment of co-repressor complexes that prevent initiation of inflammatory gene transcription. After stimulation by an activating signal, additional mechanisms can maintain quiescence by restraining active transcription. Finally, numerous mechanisms mediate the timely resolution of the inflammatory response at the transcriptional level, including transrepression mechanisms that can remove transcription factors from inflammatory gene promoters^[8,15-18].

Studies have also highlighted an important role for chromatin modifications in the transcriptional control of inflammatory gene expression^[19,20]. A recent study by Soreq *et al.*^[21], which compared transcriptional profiles of different brain cell types and regions throughout healthy human aging, found microglial gene expression profiles as being one of the most predictive markers of biological age in the brain. The same study identified a relatively unknown gene, PHD finger protein 15 (*PHF15*), among the top 25 differentially expressed genes in microglia during aging. Work in embryonic stem cells, as well as sequence and structural similarity to other members of the PHF family, indicate that PHF15 is a putative chromatin-mediated gene regulator^[22].

Given that aging skews microglia towards a proinflammatory phenotype, and that *PHF15* was found to be highly upregulated during non-pathological aging, we sought to determine whether *Phf15* might regulate microglial inflammatory function. We found that *Phf15* strongly represses proinflammatory gene expression, regulating both basal and signal-dependent activation and modulating the magnitude and duration of the mouse microglial inflammatory response. Importantly, *Phf15* seems to regulate proinflammatory and interferon type I (IFN-I)-dependent gene expression. Increased IFN-I tone and proinflammatory cytokine expression are both hallmarks of the aging brain^[23-26]. Our findings suggest that *Phf15* is an important novel repressor of microglial inflammatory function that might work to counteract age-induced inflammation in the healthy, aging brain.

METHODS

Animals

Adult male C57Bl6/J mice were purchased from The Jackson Laboratory and maintained on a 12-h/12-h light/dark cycle (lights on at 07:00) with *ad libitum* access to food and water and aged for ~2.5, ~14, or ~20 months. All animal care and procedures were approved by the University of California, Berkeley Animal Care and Use Committee.

shRNA-mediated knockdown of *Phf15* in murine microglial cells

pGIPZ lentiviral mouse *Phf15* shRNA constructs or a control scrambled shRNA were purchased from Dharmacon (Lafayette, CO). Lentivirus was packaged via co-transfection of each pGIPZ shRNA with pCMV-VSV-G (Addgene plasmid #8454)^[27] and pCMV-dR8.2 (Addgene plasmid #8455)^[27] into HEK 293T cells using Lipofectamine 3000 reagent (Life Technologies, Carlsbad, CA) according to the manufacturer's instructions. Viral supernatant was harvested after 48 h and incubated with SIM-A9 murine microglial cells in SIM-A9 complete medium [DMEM/F12 (Life Technologies, Carlsbad, CA), 10% fetal bovine serum (FBS; GE Healthcare Life Sciences, Chicago, IL), 5% horse serum (HS; GE Healthcare Life Sciences, Chicago, IL), and 1% Penicillin-Streptomycin (Life Technologies, Carlsbad, CA)]. After 48 h, GFP+ cells were sorted by Fluorescence-activated cell sorting (FACS) on an Aria Fusion (BD Biosciences, San Jose, CA; UC Berkeley Cancer Research Laboratory), expanded, and subcultured for immune stimulation experiments. Percent knockdown (KD) was determined via real-time quantitative PCR (RT-qPCR).

Overexpression of *Phf15* in murine microglial cells

A *Phf15* overexpression (OE) vector was constructed by cloning the full length *Phf15* cDNA (*Mus musculus* PHD finger protein 15, mRNA cDNA clone MGC:143877 IMAGE:40094330) obtained from Dharmacon (Lafayette, CO) into a pMYs-IRES-GFP retroviral vector (Cell Biolabs Inc., San Diego, CA). Viruses expressing the full length *Phf15* cDNA or empty vector control were co-transfected with pCL-10 A1 (Addgene plasmid #15805)^[28] in HEK 293T cells using Lipofectamine 3000 (Life Technologies, Carlsbad, CA) reagent according to the manufacturer's instructions. SIM-A9 cells were incubated with virus for 24 h and then sorted via FACS on an Aria Fusion, expanded, and subcultured for immune stimulation experiments. Fold OE was verified via RT-qPCR.

Generation of *Phf15* knockout microglia

Phf15 knockout (KO) SIM-A9 cells were generated using the Alt-R CRISPR-Cas9-mediated gene editing system (guide RNA sequence ACTACATCCTGGCGGACCCGTGG) from IDT (Coralville, IA) using CRISPRMAX Lipofectamine reagent (IDT) as per the manufacturer's instructions. ATTO 550+ cells were single-cell sorted on an Aria Fusion. Clones were screened for *Phf15* deletion using PCR (primers Forward: agcacacttgtaaccctct and Reverse: gaccaatgtctgtgtgttcg) followed by restriction digest with BtgI (New England Biolabs, Ipswich, MA). Percent decrease in *Phf15* mRNA transcript expression was determined via RT-qPCR. Primer sequences are listed in [Supplementary Table 1](#).

Immune stimulation

For all immune stimulation time course experiments, cells (KD, KO, and OE, and respective controls) were subcultured in 24-well plates at a density of 0.05×10^6 cells/well (in triplicate) and stimulated with lipopolysaccharide (LPS; final concentration of 100 ng/mL; Sigma Aldrich, St. Louis, MO), CpG oligodeoxynucleotide (CpG ODN; final concentration of 2.5 μ mol/L; InvivoGen, San Diego, CA), or polyinosinic:polycytidylic acid [Poly(I:C); final concentration of 25 μ mol/L; Sigma Aldrich, St. Louis, MO] for 1, 6, 12, or 24 h. No stimulation controls received an equivalent volume of sterile 1 \times PBS (Invitrogen, Carlsbad, CA).

RNA extraction

Mice were sacrificed according to the approved protocol. Brains were quickly isolated and frontal cortical areas were dissected, flash frozen, and stored at -80 °C. RNA was extracted using a bead homogenizer (30 s, setting “5”; Bead Mill, VWR) in Trizol reagent (ThermoFisher, Waltham, MA). Total RNA was extracted using the Direct-zol RNA miniprep kit (Zymo Research, Irvine, CA) according to the manufacturer's instructions. For cell lines, after immune stimulation, media was aspirated and wells were washed 2 × with ice-cold 1 × PBS (Invitrogen, Carlsbad, CA). RNA was extracted using the Direct-zol RNA miniprep kit (Zymo Research, Irvine, CA).

RT-qPCR

cDNA was reversed transcribed from total RNA using the SuperScriptTM III First-Strand Synthesis System kit (ThermoFisher, Waltham, MA) following the manufacturer's instructions. RT-qPCR was run using SYBR green (Roche, Pleasanton, CA) on a QuantStudio 6 (ThermoFisher, Waltham, MA) real-time PCR machine. All RT-qPCR primers were specific to the desired template, spanned exon-exon junctions, and captured all transcript variants for the specific gene under study. Ct values were normalized to the housekeeping gene hypoxanthine phosphoribosyltransferase (*Hprt*). Primer sequences used in this study are listed in [Supplementary Table 1](#).

RNA-seq library preparation and analysis

RNA was extracted from a total of $n = 3$ replicates per condition (*Phf15* KO or control) and was used to prepare libraries for RNA sequencing using the mRNA HyperPrep Kit according to the manufacturer's instructions (KAPA Biosystems, Wilmington, MA). Libraries were quality control checked via Qubit (ThermoFisher, Waltham, MA) and via RT-qPCR with a next generation sequencing library quantification kit (Zymo Research, Irvine, CA). RNA sequencing (one lane) was performed on a HiSeq4000 sequencing system (Illumina Inc., San Diego, CA; UC Berkeley Genomics Sequencing Laboratory). Sequencing reads were aligned to the *Mus musculus* reference genome assembly GRCm38 (mm10) using Spliced Transcripts Alignment to a Reference (STAR) aligner^[29]. Count data were analyzed with Hypergeometric Optimization of Motif EnRichment (HOMER) software for next-generation sequencing analysis (<http://homer.ucsd.edu/homer/ngs/index.html>), which uses the R/Bioconductor package DESeq2^[30] to perform differential gene expression analysis. To adjust for multiple comparisons, DESeq2 uses the Benjamini-Hochberg procedure to control the false discovery rate and returned false discovery rate adjusted P values and \log_2 -fold expression changes between *Phf15* KO and control conditions for each gene. Genes were filtered by adjusted P value (adjusted $P < 0.01$ for upregulated genes or 0.05 for downregulated genes) and \log_2 -fold change in expression (greater than 1.5 \log_2 -fold change for upregulated genes and less than -1.5 for downregulated genes). Too few downregulated genes (< 200) passed the more stringent adjusted $P < 0.01$ cutoff for robust downstream biological function analysis, thus the adjusted P value threshold was lowered to $P_{adj} < 0.05$. Results were visualized using the R package EnhancedVolcano^[31]. Lists of upregulated and downregulated genes were input into Metascape^[32], a gene annotation and analysis tool, to determine enriched biological themes within the gene lists.

Motif enrichment

Transcription factor binding site (“motif”) enrichment was analyzed using HOMER (<http://homer.ucsd.edu/homer/ngs/index.html>).

Statistical analysis

Relative mRNA expression of *Phf15* in mouse frontal cortical areas was analyzed using ordinary one-way ANOVA with *post hoc* Tukey's multiple comparisons to compare expression levels across age. Percent KD and time course experiments measuring expression levels of inflammatory markers [*Tnfα*, nitric oxide synthase, inducible (*Nos2*), and *IL-1β*] between control and *Phf15* shRNAs sh*Phf15*-1 and sh*Phf15*-2 after




Figure 1. *Phf15* expression increases in aged mouse frontal cortical areas. *Phf15* mRNA expression was significantly elevated in frontal cortical areas of old (~20-month-old; red bar) mice compared to young (~2.5-month-old; black bar) mice. Data are mean ± SEM ($n = 4$ young, $n = 5$ middle aged, $n = 12$ old). One-way ANOVA with Tukey's post hoc comparisons between age groups: ** $P < 0.01$

immune stimulation [with LPS, CpG-ODN, or Poly(I:C)] were analyzed via ordinary one-way ANOVA with Dunnett's multiple comparisons between each shRNA versus control shRNA within timepoint. Fold OE or percent reduction for *Phf15* OE and KO cell lines, respectively, were analyzed using unpaired *t*-tests (OE or KO vs. respective control). Time course experiments for *Phf15* OE and KO cell lines were analyzed using unpaired *t*-tests with Holm-Sidak correction for multiple comparisons. RNA-seq data were analyzed as described above. $P < 0.05$ was considered significant in all experiments.

RESULTS

Aging increases *Phf15* expression in mouse brain

To investigate whether *Phf15* increases in mouse brains similar to humans^[21], we measured *Phf15* mRNA expression in mouse frontal cortical brain areas across age. We were interested in frontal cortical regions because of their involvement in mediating various aspects of cognitive function and because they are selectively affected in several aging-related neurodegenerative conditions, e.g., PD, AD, and frontotemporal dementia^[33,34].

We found that compared to young (~2.5-month-old) mice, old (~20-month-old) mice had significantly elevated *Phf15* mRNA levels in frontal cortical areas [Figure 1]. Middle-aged (~14-month-old) mice showed a trend towards increased *Phf15* mRNA expression that did not reach statistical significance. Our data suggest that *Phf15* expression increases in mouse frontal cortical regions upon normal aging, similar to what was previously reported in humans^[21].

Knockdown of *Phf15* increases the magnitude of the microglial inflammatory response

To determine whether *Phf15* regulates microglial inflammatory function, we performed loss-of-function studies via shRNA-mediated KD in a murine microglial cell line, SIM-A9, followed by immune activation with LPS, a component of Gram-negative bacterial cell walls and TLR4 agonist. SIM-A9 cells are an established murine microglial cell line and their original characterization has been previously published^[35]. In short, SIM-A9 cells are a spontaneously transformed microglial cell line isolated from cultured primary glia from postnatal murine cerebral cortices. They display key characteristics of cultured primary microglia; for example, they express the microglia/macrophage-specific proteins cluster of differentiation 68 (CD68) and ionized calcium-binding adapter molecule 1 (IBA1) and they are responsive to immune

stimulation with LPS, triggering nuclear factor kappa-light-chain-enhancer of activated B cells (NF- κ B) inflammatory signaling cascades, increase of protein levels of the proinflammatory factors NOS2 (the enzyme that catalyzes the production of NO) and cyclooxygenase 2b (COX2), and secretion of TNF α . Importantly, they are also responsive to the anti-inflammatory effects of interleukin 4 and increasing levels of Arginase 1, demonstrating that SIM-A9 cells can switch between pro- and anti-inflammatory states. Finally, they also exhibit phagocytic uptake of bacterial particles and fluorescently-labeled amyloid- β (A β).

Additionally, we chose LPS as the immune stimulant because: (1) intraperitoneal and/or intracranial administration of LPS in mice led to increased microglial activation, neuroinflammation, neuronal loss including loss of dopaminergic neurons in the substantia nigra in a mouse model of PD^[8], and cognitive and neurological deficits^[36]; (2) aged individuals show increased systemic levels of LPS in the bloodstream^[37], which are associated with increased inflammation and microglial activation^[38]; and (3) in humans, TLR4 activation is linked to age-related pathologies such as PD and AD^[39-41]; thus, LPS serves as a relevant aging-related physiological immune stimulant.

KD of *Phf15* resulted in a significant reduction in *Phf15* mRNA transcript levels of 52% and 60% for cell lines sh*Phf15*-1 and sh*Phf15*-2, respectively [Figure 2A], as well as significantly increased mRNA expression of *Tnf α* , a proinflammatory cytokine, after KD with sh*Phf15*-2 at 0, 1, 6, and 12 h after LPS stimulation [Figure 2B].

Similarly, mRNA levels of *Nos2* were significantly elevated at 1, 6, and 12 h post stimulation for sh*Phf15*-2 and 0, 6, and 12 h for sh*Phf15*-1 [Figure 2D]. Overall, our experiments show that ~50%-60% KD, the equivalent of a “heterozygous” condition, results in increased expression of proinflammatory mediators over a 12-h time course that resolves and falls below control levels by 24 h after immune stimulation. Importantly, microglial inflammatory function was elevated in the absence of immune stimulation (0 h time point, Figure 2B and D, and no stimulation condition, Figure 2C and E), suggesting a loss of repressive mechanisms that inhibit basal state inflammatory gene transcription.

We repeated the immune activation time course experiments in *Phf15* KD cells using two separate immune stimulants specific to two distinct Toll-like receptors to test the pathway specificity of the inflammatory response: CpG ODN, a synthetic bacterial and viral DNA mimic that targets TLR9, and Poly(I:C), a synthetic viral dsRNA mimic that targets TLR3. While TLR4 uses both the myeloid differentiation primary response 88 (MyD88) and TIR-domain-containing adapter-inducing interferon- β (TRIF) downstream adapters to transduce its inflammatory cascade, TLR9 and TLR3 utilize MyD88 and TRIF, respectively [Supplementary Figure 1]^[42,43].

Immune stimulation with CpG ODN and Poly(I:C) both yielded similar results to those obtained with LPS stimulation [Supplementary Figures 2 and 3, respectively], denoting no adapter selectivity and confirming that *Phf15* antagonizes inflammatory gene expression downstream of both the MyD88 and TRIF signaling pathways.

Genetic deletion of *Phf15* increases the magnitude and prolongs the duration of the microglial inflammatory response

Since our KD strategy resulted in ~50% reduction in *Phf15* mRNA expression, we next performed CRISPR/Cas9-mediated genetic deletion of *Phf15* in SIM-A9 microglial cells followed by immune activation with LPS. KO of *Phf15* [Figure 3A] resulted in significantly increased LPS-induced expression of *Tnf α* [Figure 3B], *IL-1 β* [Figure 3D], and *Nos2*, albeit to a lesser extent [Figure 3F], over a 24-h time course. Importantly, mRNA levels of both *Tnf α* and *IL-1 β* remained elevated at 24 h compared to control cells, denoting a prolonged inflammatory response and failure to return to steady-state. mRNA expression of *Nos2* showed a significant




Figure 2. Knockdown of *Phf15* increases the magnitude of the microglial inflammatory response. (A) Knockdown efficiency for anti-*Phf15* shRNAs sh*Phf15*-1 (blue bar, 52% knockdown) and sh*Phf15*-2 (red bar, 60% knockdown). Data are mean \pm SEM ($n = 3$ per condition). One-way ANOVA with Dunnett's multiple comparisons between sh*Phf15*-1 or sh*Phf15*-2 and shCtrl cells: $^{**}P < 0.01$. Twenty-four-hour time course experiments showing relative mRNA expression levels of *Tnfα* (B) and *Nos2* (D) after LPS stimulation of shRNAs sh*Phf15*-1 and sh*Phf15*-2 compared to shCtrl (control scrambled shRNA). No stimulation (0-h time point) is shown for *Tnfα* (C) and *Nos2* (E). Data are mean \pm SEM ($n = 3$ per condition). One-way ANOVA with Dunnett's multiple comparisons for sh*Phf15*-1 or sh*Phf15*-2 relative to shCtrl cells for individual timepoints: $^{*}P < 0.05$, $^{**}P < 0.01$, $^{***}P < 0.001$, $^{****}P < 0.0001$. LPS: lipopolysaccharide; *Tnfα*: tumor necrosis factor alpha; *Nos2*: nitric oxide synthase, inducible

upregulation over 12 h (0-, 1-, and 12-h timepoints) but had returned to control levels by 24 h [Figure 3F]. Notably, basal expression of all three genes was significantly elevated, with a 4-fold increase in *Tnfα*, 14-fold increase in *IL-1β*, and 32-fold increase in *Nos2* when comparing KO to control cells [Figure 3C, E and G].

Time course experiments after stimulation of TLR9 with CpG-ODN [Supplementary Figure 4] and TLR3 with Poly(I:C) [Supplementary Figure 5] in *Phf15* KO cells again yielded similar results to LPS stimulation in *Phf15* KO microglial cells, denoting no difference in downstream adapter selectivity and confirming our prior KD results.

Overall, KO of *Phf15* resulted in a more severe phenotype compared to our KD results, increasing the magnitude and prolonging the duration of the microglial inflammatory response. Taken together, our KD and KO results indicate that *Phf15* functions to restrict microglial inflammatory output, regulating the magnitude and duration, as well as basal inhibition of the inflammatory response.




Figure 3. Knockout of *Phf15* increases the magnitude and duration of inflammatory gene expression. (A) Percent reduction in *Phf15* transcript expression in *Phf15* knockout SIM-A9 microglia (*Phf15* KO, red bar) compared to control (Ctrl, open bar). Data are mean \pm SEM ($n = 3$ per condition). Unpaired t -test between *Phf15* KO and control cells. Twenty-four-hour time course experiments showing relative mRNA expressions levels of *Tnfα* (B), *IL-1β* (D), and *Nos2* (F) after LPS stimulation. No stimulation (0-h time point or baseline) expressions of *Tnfα* (C), *IL-1β* (E), and *Nos2* (G) are also shown. Data are mean \pm SEM ($n = 3$ per condition). Unpaired t -tests with Holm-Sidak correction for multiple comparisons between *Phf15* KO and control cells within timepoint: * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$. LPS: lipopolysaccharide; KO: knockout; *Tnfα*: tumor necrosis factor alpha; inducible; *IL-1β*: interleukin 1 beta; *Nos2*: nitric oxide synthase, inducible

Overexpression of *Phf15* in microglia results in a dampened inflammatory response

To further test the role of *Phf15* as a repressor of proinflammatory genes, we carried out gain-of-function studies of *Phf15* in SIM-A9 cells. OE via retroviral delivery of the full-length murine *Phf15* cDNA [Figure 4A] resulted in significantly decreased expression of *Nos2* at 0 and 24 h [Figure 4F]. mRNA expression levels of *Tnfα* [Figure 4B], *IL-1β* [Figure 4D], and *Nos2* [Figure 4F] were decreased at 6 h post LPS stimulation at an adjusted $P = 0.0501$, which did not reach statistical significance. Notably, basal levels (no stimulation) of *Nos2* were also significantly decreased [Figure 4E, G].

Time course experiments following stimulation with CpG-ODN [Supplementary Figure 6] and Poly(I:C) [Supplementary Figure 7] showed a similar but stronger repressive phenotype compared to LPS




Figure 4. *Phf15* overexpression decreases the microglial inflammatory response. (A) Fold OE of *Phf15* in SIM-A9 microglia (red bar) versus control cells (Ctrl, open bar). Data are mean \pm SEM ($n = 3$ per condition). Unpaired t -test between *Phf15* OE and control cells. Twenty-four-hour time course experiments showing relative mRNA expression levels of *Tnfα* (B), *IL-1β* (D), and *Nos2* (F) after LPS stimulation. Baseline (0-h time point, no stimulation) expressions of *Tnfα* (C), *IL-1β* (E), and *Nos2* (G) are displayed separately from time course experiments. Data are mean \pm SEM ($n = 3$ per condition). Unpaired t -tests with Holm-Sidak correction for multiple comparisons between *Phf15* OE and control cells within timepoint: * $P < 0.05$, ** $P < 0.01$. OE: overexpression; LPS: lipopolysaccharide; *Tnfα*: tumor necrosis factor alpha; inducible; *IL-1β*: interleukin 1 beta; *Nos2*: nitric oxide synthase, inducible

stimulation, with decreased expression levels of *Tnfα* [Supplementary Figures 6A, B and Figures 7A, B], *IL-1β* [Supplementary Figures 6C, D and Figures 7C, D], and *Nos2* [Supplementary Figures 6E, F and Figures 7E, F], over the time course, as well as under basal (no stimulation) conditions, confirming our previous results. Taken together, our OE results show a dampened microglial inflammatory response, revealing a reciprocal response phenotype compared to our KD and KO experiments. Collectively, these results confirm that *Phf15* functions to repress both basal and stimulus-dependent inflammatory gene expressions in microglia.

Loss of *Phf15* affects global expression of genes involved in antiviral responses and regulation of inflammatory processes

To examine global transcriptional changes as a result of *Phf15* deletion in microglia, we carried out RNA-sequencing (RNA-seq) on *Phf15* KO SIM-A9 cells under no stimulation conditions and 6 h post LPS stimulation. We chose to examine the no stimulation condition (0-h time point) based on our KD and KO time course results showing that baseline is one of the most consistently and strongly deregulated time points. Importantly, elevated or “leaky” proinflammatory mediator expression at baseline might result in chronic inflammation leading to neurodegeneration. Similarly, 6 h after LPS stimulation corresponded to the peak of the transcriptional inflammatory response, with large increases in magnitude for both *IL-1 β* and *Nos2*.

Differential gene expression analysis revealed that 466 genes with \log_2 -fold change > 1.5 and $P_{adj} < 0.01$ were upregulated and 309 genes with \log_2 -fold change < -1.5 and $P_{adj} < 0.05$ were downregulated [Figure 5A]. Biological theme enrichment analysis using Metascape^[32] on the upregulated genes revealed that the most enriched biological process categories under basal conditions were “response to virus” and “cytokine production” [Figure 5B and C]. Under the “response to virus” category, there was significant upregulation of various interferon-stimulated genes (ISGs), for example *Isg15*, interferon induced protein with tetratricopeptide repeats 1 (*Ifit1*), *Ifit3*, interferon regulatory factor 7 (*Irf7*), 2'-5'-oligoadenylate synthetase 2 (*Oas2*), and Oas-like 2 (*Oasl2*) [Figure 5C]. The downregulated genes showed more variability in the types of pathways affected, largely involving growth, differentiation, and glial cell migration processes [Figure 5A and Supplementary Figure 8A].

Motif analysis for transcription factor binding sites enriched in the promoters of the upregulated genes at baseline revealed consensus motifs for interferon regulatory factors (IRFs), i.e., the interferon (IFN) stimulated response element (ISRE) and motifs for IRF3 and IRF8 specifically in the top 5 best matches. Activator protein 1 (AP-1) and NF- κ B p65 subunit (NF- κ B-p65) motifs were also enriched. Both can regulate expression of canonical proinflammatory cytokines such as *TNF α* and *IL-1 β* ^[44,45] [Figure 5D]. Motif enrichment for the set of downregulated genes revealed motifs for twist-related protein 2 (Twist2) and Class A basic helix-loop-helix protein 15 (BHLHA15). Twist2 has been shown to mediate cytokine downregulation after chronic nucleotide-binding oligomerization domain-containing protein 2 (NOD2, a bacterial peptidoglycan sensor) stimulation^[46]. BHLHA15 has been shown to induce and maintain secretory architecture in cells specialized for secretion^[47] [Supplementary Figure 8B].

Differential gene expression analysis after 6 h of LPS stimulation in KO versus control cells revealed 576 upregulated genes (\log_2 -fold change > 1.5 and $P_{adj} < 0.01$) and 322 downregulated genes (\log_2 -fold change < -1.5 and $P_{adj} < 0.05$) [Figure 6A]. Interestingly, by 6 h after LPS administration, some of the most enriched biological process categories in KO cells were related to “cytokine secretion” and “immunoregulatory interaction” [Figure 6B, C], denoting a strong increase in magnitude of expression of genes involved in regulating the secretion of proinflammatory mediators. The downregulated genes at 6 h after LPS stimulation in KO cells relative to control again displayed more variability, but did show decreases in biological process categories related to “regulation of defense response” and “cytokine production”, indicating negative regulation of these processes in *Phf15* KO cells compared to control [Supplementary Figure 9A].

Motif enrichment analysis for transcription factor binding sites enriched in the promoters of upregulated genes at the 6-h time point revealed consensus sequences for AP-1, a key regulator of microglia reactivity in inflammation^[48] [Figure 6D]. Motif enrichment for the set of downregulated genes revealed motifs for IRFs (the ISRE) and motifs for IRF1 and IRF3 specifically [Supplementary Figure 9B], supporting the observation that there is a negative “regulation of defense response” by 6 h post stimulation. It is interesting to note that a functional transition from cytokine production to cytokine secretion seems to occur in the 6-h period after LPS activation.




Figure 5. Loss of *Phf15* affects the expression of genes involved in viral response and regulation of inflammatory processes in the absence of immune stimulation. A: Volcano plot representing the RNA-seq results. Orange dots represent differentially expressed genes in *Phf15* knockout microglia compared to control (upregulated genes at a cutoff of \log_2 -fold change > 1.5 and P_{adj} < 0.01; downregulated genes at a cutoff of \log_2 -fold change < -1.5 and P_{adj} < 0.05); B: GO analysis for significantly upregulated genes showing biological process categories related to “response to virus” and “inflammatory response”; C: Upregulated genes associated with “response to virus” and “inflammatory response” in the no stimulation (baseline) condition. Relative FPKM values were obtained by normalizing FPKM values of *Phf15* knockout SIM-A9 microglia to control FPKM values for each gene ($n = 3$ per condition). Statistics are by DESeq2: # P < 0.0001; D: Top 5 enriched transcription factor binding motifs for the set of upregulated genes in the no stimulation (baseline) condition. KO: knockout; FPKM: Fragments per kilobase million; GO: Gene ontology




Figure 6. Knockout of *Phf15* affects the expression of genes involved in inflammatory factor secretion and immunoregulatory processes after LPS stimulation. A: volcano plot representing the RNA-seq results. Orange dots represent differentially expressed genes in *Phf15* knockout microglia 6 h after LPS administration compared to control (upregulated genes at a cutoff of \log_2 -fold change > 1.5 and $P_{adj} < 0.01$; downregulated genes at a cutoff of \log_2 -fold change < -1.5 and $P_{adj} < 0.05$); B: GO analysis for upregulated genes shows biological process categories associated with “cytokine secretion” and “immunoregulatory interaction”; C: Upregulated genes associated with “cytokine secretion” and “immunoregulatory interaction” biological process categories 6 h post LPS stimulation. Relative FPKM values were obtained by normalizing FPKM values of *Phf15* knockout SIM-A9 microglia to control FPKM values for each gene ($n = 3$ wells per condition). Statistics are by DESeq2: ** $P < 0.01$, \$ $P < 0.001$, # $P < 0.0001$; D: Transcription factor binding motifs for the set of upregulated genes 6 h after LPS stimulation are enriched for activator protein 1 (AP-1). LPS: lipopolysaccharide; KO: knockout; FPKM: Fragments per kilobase million; GO: Gene ontology

Taken together, our RNA-seq results confirm that *Phf15* is a repressor of microglial inflammatory gene expression, regulating the antiviral responses - specifically, IFN-I-dependent responses - as well as processes related to proinflammatory cytokine production and release.

DISCUSSION

Our results show that *Phf15* inhibits microglial expression of proinflammatory mediators under basal and signal-dependent activation, regulating both the magnitude and duration of the inflammatory response. Genetic deletion of *Phf15* in a microglial cell line followed by stimulation with LPS led to an exaggerated proinflammatory response with increased production of *Tnfα*, *IL-1β*, and *Nos2* over a time course of 24 h. Importantly, levels of proinflammatory factors remained elevated at 24 h, demonstrating a sustained and prolonged response. Consistent with our LPS stimulation of TLR4 results, similar results were obtained after TLR9 and TLR3 activation, confirming that *Phf15* is a general negative regulator and controls both the MyD88 and TRIF downstream signal transduction pathways [Supplementary Figure 1]. Overexpression of *Phf15* showed a dampened microglial inflammatory response, highlighting a reciprocal response phenotype that further supports our loss-of-function results.

Prolonged inflammation can damage surrounding healthy tissue, eventually resulting in neuronal degeneration and loss, and negatively affecting brain function. For example, levels of *TNFα* are seen to rapidly rise in experimental models of PD and are highly toxic to dopaminergic neurons^[13,14,49]. Similarly, high levels of *TNFα* are a hallmark of PD in humans^[50-52]. Additionally, both *TNFα* and *IL-1β* are involved in maintaining proper synaptic plasticity at physiological levels^[53,54] and overproduction of these cytokines can result in neuronal death via excitotoxicity and cognitive dysfunction^[55,56].

Our studies further demonstrate that *Phf15* can regulate both basal and signal-dependent microglial inflammatory gene expression. KD and KO of *Phf15* in microglial cell lines resulted in significantly increased levels of proinflammatory cytokine gene expression - without stimulation and after immune activation - while OE had the reverse effect. The inflammatory response is a tightly controlled process in immune cells in order to protect against unintended damage to healthy tissue. Even in aged microglia, where production and secretion of proinflammatory mediators is generally increased, this process is dependent upon treatment with immune stimulants^[9,57,58]. Increased proinflammatory cytokine gene expression without stimulation denotes constitutive or “leaky” expression of inflammatory mediators, simulating a state of low-grade but constant activation. Similarly, hyperresponsiveness to immune stimuli combined with a lack of resolution of the inflammatory response can lead to a state of chronic inflammation. All three can trigger pathological chronic inflammation in the brain, which is detrimental to brain function.

Importantly, distinct molecular mechanisms regulate transcriptional control of different phases (“modules”) of the inflammatory response and it is noteworthy that *Phf15* might be involved in regulating several of these. Basal inflammatory function, for example, is generally regulated by co-repressors such as nuclear receptor co-repressor (NCOR), silencing mediator of retinoid and thyroid receptors (SMRT), and RE1 silencing transcription factor (REST) co-repressor 1 (RCOR1 or CoREST) that block poised promoters from active transcription, preventing “leaky” expression of primary response genes (e.g., *TNFα*, Type I IFNs, *IL-1β*, etc.) (for review, see^[17]). Significantly increased inflammatory gene transcription under baseline conditions, as observed in our *Phf15* KD and KO experiments, suggests a loss of this repressive mechanism.

After stimulation by an activating signal, additional mechanisms can maintain quiescence by restraining active transcription. For example, nuclear receptors such as peroxisome proliferator-activated receptor-γ, glucocorticoid receptor, and liver X receptors can inhibit the signal-activated exchange of co-repressors for co-activators at poised promoters, inhibiting the initiation of transcription^[15,17]. Lastly, several mechanisms regulate resolution of inflammation at the transcriptional level, including transrepression

mechanisms that can remove transcription factors such as NF- κ B from inflammatory gene promoters, effectively blocking expression of secondary response genes, that is, genes which require chromatin-modification as well as protein synthesis for their induction (e.g., *Nos2* and ISGs)^[8,16,17]. Timely resolution of an inflammatory response is crucial in order to limit cellular and tissue damage caused by prolonged or chronic inflammation. Our results suggest that *Phf15* may be involved in regulating all three of the abovementioned mechanisms.

However, how might *Phf15* be involved in regulating transcriptional repression of the inflammatory response in microglia? PHF15 was first described in embryonic stem cells as an E3 ligase that directly targets Lysine-specific demethylase 1 (LSD1, *Kdm1a*) - a key demethylase of histone 3 lysine 4 - for degradation^[22]. LSD1 has been identified as a member of the CoREST co-repressor complex^[59,60], which is required for transcriptional repression of inflammation in microglia^[8]. We therefore initially hypothesized that increased levels of *Phf15* upon aging might lead to decreased levels of LSD1 and increased microglial inflammatory output. Our results, however, demonstrate that *Phf15* itself inhibits microglial inflammatory function; thus, its purported mechanism for inhibition is likely not via degradation of LSD1.

Interestingly, the global transcriptional changes caused by *Phf15* deletion are highly similar to previously reported age-associated transcriptional changes in microglia^[9,61,62]. In particular, a study by Deczkowska et al.^[63], found “immune system process” and specifically “response to virus” among the most highly upregulated biological categories for differentially expressed genes in microglia of young (2-month old) versus aged (22-month old) mice, consistent with our results in *Phf15* KO microglia. Notably, a study by Hammond et al.^[62], which used single-cell RNAseq to look at microglia profiles throughout the mouse lifespan, found subpopulations in aged (P540) mouse brains which were largely: (1) inflammatory, that is, they upregulated *IL-1 β* , *Tnf α* , and other cytokines; or (2) IFN-I-responsive, upregulating *Irf7* and ISGs, particularly *Ifit3*, *Isg15*, *Oasl2*, interferon induced transmembrane protein 3 (*Ifitm3*), and receptor transporter protein 4 (*Rtp4*), compared to younger adult (P100) brains. Similarly, a recent study from the Tabula Muris Consortium^[64], which produced a single-cell transcriptomic atlas of 23 tissues and organs across the *Mus musculus* life span, confirmed that microglia in the aged (P540 and P720) brain are enriched for IFN-I-responsive genes and upregulate a similar set of genes including *Ifit3*, *Irf7*, *Isg15*, *Oasl2*, *Ifitm3*, and *Rtp4*. The genes upregulated by the interferon-responsive microglia clusters in both studies are highly similar to those upregulated in our *Phf15* KO cells under basal conditions [see Figure 5A and C]. Because ISGs can modulate inflammation^[23], it is possible that interferon-responsive microglia could play a role in contributing to the inflammatory signature found in the aged brain. Interestingly, among the set of downregulated genes in *Phf15* KO cells at baseline and 6 h after LPS stimulation is myocyte enhancer factor 2C (*Mef2C*). MEF2C is an important checkpoint inhibitor that restrains microglial activation in response to proinflammatory insults and is lost in brain aging via IFN-I mediated downregulation^[63,65]. Thus, an increase in *Phf15* expression in microglia during healthy aging could putatively work to counteract not only microglial activation but increased IFN-I in the aged brain as well.

Notably, a recent study by Readhead et al.^[66] found that several virus species are commonly present in the aged human brain. Among them, human herpesvirus 6A and 7 (HHV-6A and HHV-7) were highly upregulated in the brain of AD patients and were found to modulate host genes associated with AD risk, for example amyloid precursor protein (*APP*) processing. APP is the precursor molecule whose proteolysis forms A β and formation of A β plaques has long been thought of as the driving force behind Alzheimer’s disease^[67]. A β has more recently been found to have antimicrobial properties^[68], conferring increased resistance against infection from both bacteria and viruses^[69]. *App* is among the significantly upregulated genes under basal conditions in our *Phf15* KO cells (log₂-fold change = 1.492 and *P adj* < 0.0001; see Figure 5A). Upregulation of *App* due to loss of *Phf15* in mouse microglia is thus consistent with our data showing *Phf15* regulation of the antiviral microglial response.

Altogether, our results show that *Phf15* is a novel repressor of microglial inflammatory gene expression, regulating both the magnitude and time-to-resolution of the inflammatory response. Importantly, *Phf15* also serves to repress baseline inflammatory output in the absence of immune activation. Putatively, increases in *Phf15* during healthy aging could help counteract brain inflammation and protect brain health.

Future studies will determine the mechanism of action of *Phf15*. For example, the identity of its binding partner proteins and its genome-wide binding sites and associated histone marks will be elucidated to determine the specific gene regulatory regions it interacts with (e.g., active enhancers or promoters). Additionally, studies in *Phf15* KO mice will determine whether loss of *Phf15*-mediated repression of proinflammatory factors is sufficient to induce cognitive decline or exacerbate LPS-induced neurotoxicity of dopaminergic neurons in the substantia nigra.

DECLARATIONS

Acknowledgments

We thank Prof. Ellen Robey for helpful comments on the manuscript and Wendy Yan for technical assistance.

Authors' contributions

Designed and performed experiments, analyzed data, and wrote the manuscript: Muroy SE

Performed experiments and analyzed data: Timblin GA, Preininger MK

Performed experiments: Cedillo P

Designed experiments and wrote the manuscript: Saijo K

Availability of data and materials

Not applicable.

Financial support and sponsorship

This work was supported by the Berkeley Fellowship to S.E.M., ADA Postdoctoral fellowship to G.A.T., NSF GRFP to M.K.P., and R01HD092093 and Pew Scholarship to K.S.

Conflicts of interest

All authors declared that there are no conflicts of interest.

Ethical approval and consent to participate

All procedures were approved by the Animal Care and Use Committee of the University of California, Berkeley (Animal Use Protocol AUP-2017-02-9539).

Consent for publication

Not applicable.

Copyright

© The Author(s) 2020.

REFERENCES

1. Kettenmann H, Hanisch UK, Noda M, Verkhratsky A. Physiology of microglia. *Physiol Rev* 2011;91:461-553.
2. Nimmerjahn A, Kirchhoff F, Helmchen F. Resting microglial cells are highly dynamic surveillants of brain parenchyma in vivo. *Science* 2005;308:1314-8.
3. Parkhurst CN, Gan WB. Microglia dynamics and function in the CNS. *Curr Opin Neurobiol* 2010;20:595-600.
4. Ransohoff RM, Perry VH. Microglial physiology: unique stimuli, specialized responses. *Ann Rev Immunol* 2009;27:119-45.
5. Saijo K, Glass CK. Microglial cell origin and phenotypes in health and disease. *Nat Rev Immunol* 2011;11:775-87.

6. Wake H, Moorhouse AJ, Miyamoto A, Nabekura J. Microglia: actively surveying and shaping neuronal circuit structure and function. *Trends Neurosci* 2013;36:209-17.
7. Hanisch UK, Kettenmann H. Microglia: active sensor and versatile effector cells in the normal and pathologic brain. *Nat Neurosci* 2007;10:1387-94.
8. Saijo K, Winner B, Carson CT, Collier JG, Boyer L, et al. A Nurr1/CoREST pathway in microglia and astrocytes protects dopaminergic neurons from inflammation-induced death. *Cell* 2009;137:47-59.
9. Mosher KI, Wyss-Coray T. Microglial dysfunction in brain aging and Alzheimer's disease. *Biochem Pharmacol* 2014;88:594-604.
10. Hickman S, Izzy S, Sen P, Morsett L, El Khoury J. Microglia in neurodegeneration. *Nat Neurosci* 2018;21:1359-69.
11. Salter MW, Stevens B. Microglia emerge as central players in brain disease. *Nat Med* 2017;23:1018-27.
12. Glass CK, Saijo K, Winner B, Marchetto MC, Gage FH. Mechanisms underlying inflammation in neurodegeneration. *Cell* 2010;140:918-34.
13. McCoy MK. Blocking soluble tumor necrosis factor signaling with dominant-negative tumor necrosis factor inhibitor attenuates loss of dopaminergic neurons in models of Parkinson's disease. *J Neurosci* 2006;26:9365-75.
14. Sriram K, Matheson JM, Benkovic SA, Miller DB, Luster MI, et al. Mice deficient in TNF receptors are protected against dopaminergic neurotoxicity: implications for Parkinson's disease. *FASEB J* 2002;16:1474-6.
15. Glass CK, Ogawa S. Combinatorial roles of nuclear receptors in inflammation and immunity. *Nat Rev Immunol* 2005;6:44-55.
16. Glass CK, Saijo K. Nuclear receptor transrepression pathways that regulate inflammation in macrophages and T cells. *Nat Rev Immunol* 2010;10:365-76.
17. Medzhitov R, Horng T. Transcriptional control of the inflammatory response. *Nat Rev Immunol* 2009;9:692-703.
18. Smale ST, Natoli G. Transcriptional control of inflammatory responses. *Cold Spring Harb Perspect Biol* 2014;6:a016261.
19. Bernstein BE, Meissner A, Lander ES. The mammalian epigenome. *Cell* 2007;128:669-81.
20. Smale ST, Tarakhovskiy A, Natoli G. Chromatin contributions to the regulation of innate immunity. *Ann Rev Immunol* 2014;32:489-511.
21. Soreq L, Rose J, Soreq E, Hardy J, Trabzuni D, et al. Major shifts in glial regional identity are a transcriptional hallmark of human brain aging. *Cell Rep* 2017;18:557-70.
22. Han X, Gui B, Xiong C, Zhao L, Liang J, et al. Destabilizing LSD1 by Jade-2 promotes neurogenesis: an antibraking system in neural development. *Mol Cell* 2014;55:482-94.
23. Baruch K, Deczkowska A, David E, Castellano JM, Miller O, et al. Aging. Aging-induced type I interferon response at the choroid plexus negatively affects brain function. *Science* 2014;346:89-93.
24. Gabuzda D, Yankner BA. Physiology: inflammation links ageing to the brain. *Nature* 2013;497:197-8.
25. Godbout JP, Chen J, Abraham J, Richwine AF, Berg BM, et al. Exaggerated neuroinflammation and sickness behavior in aged mice following activation of the peripheral innate immune system. *FASEB J* 2005;19:1329-31.
26. Sparkman NL, Johnson RW. Neuroinflammation associated with aging sensitizes the brain to the effects of infection or stress. *Neuroimmunomodulation* 2008;15:323-30.
27. Stewart SA. Lentivirus-delivered stable gene silencing by RNAi in primary cells. *RNA* 2003;9:493-501.
28. Sena-Esteves M, Tebbets JC, Steffens S, Crombleholme T, Flake AW. Optimized large-scale production of high titer lentivirus vector pseudotypes. *J Virol Methods* 2004;122:131-9.
29. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, et al. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* 2012;29:15-21.
30. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* 2014;15:31-21.
31. Blighe K, Rana S, Lewis M. *enhancedVolcano*: Publication-ready volcano plots with enhanced colouring and labeling n.d. Available from: <https://doi.org/10.18129/B9.bioc.enhancedVolcano> [Last accessed on 6 Mar 2020]
32. Zhou Y, Zhou B, Pache L, Chang M, Khodabakhshi AH, et al. Metascape provides a biologist-oriented resource for the analysis of systems-level datasets. *Nat Commun* 2019;10:1523.
33. Craik FIM, Salthouse TA. *The handbook of aging and cognition*. 1st ed. Psychology Press; 2011.
34. Simen AA, Bordner KA, Martin MP, Moy LA, Barry LC. Cognitive dysfunction with aging and the role of inflammation. *Ther Adv Chronic Dis* 2011;2:175-95.
35. Nagamoto-Combs K, Kulas J, Combs CK. A novel cell line from spontaneously immortalized murine microglia. *J Neurosci Methods* 2014;233:187-98.
36. Zhao J, Bi W, Xiao S, Lan X, Cheng X, et al. Neuroinflammation induced by lipopolysaccharide causes cognitive impairment in mice. *Sci Rep* 2019;9:5790.
37. Glaros TG, Chang S, Gilliam EA, Maitra U, Deng H, et al. Causes and consequences of low grade endotoxemia and inflammatory diseases. *Front Biosci (Schol Ed)* 2013;5:754-65.
38. Sandiego CM, Gallezot JD, Pittman B, Nabulsi N, Lim K, et al. Imaging robust microglial activation after lipopolysaccharide administration in humans with PET. *Proc Natl Acad Sci U S A* 2015;112:12468-73.
39. Brown GC. The endotoxin hypothesis of neurodegeneration. *J Neuroinflammation* 2019;16:180.
40. Chen YC, Yip PK, Huang YL, Sun Y, Wen LL, et al. Sequence variants of toll like receptor 4 and late-onset Alzheimer's disease. *PLoS One* 2012;7:e50771.
41. Perez-Pardo P, Dodiya HB, Engen PA, Forsyth CB, Huschens AM, et al. Role of TLR4 in the gut-brain axis in Parkinson's disease: a translational study from men to mice. *Gut* 2019;68:829-43.
42. Akira S, Takeda K. Toll-like receptor signalling. *Nat Rev Immunol* 2004;4:499-511.
43. Takeda K, Akira S. TLR signaling pathways. *Semin Immunol* 2004;16:3-9.
44. Chen LF, Greene WC. Shaping the nuclear action of NF- κ B. *Nat Rev Mol Cell Biol* 2004;5:392-401.

45. McDonald PP. Transcriptional regulation in neutrophils: teaching old cells new tricks. *Adv Immunol* 2004;82:1-48.
46. Zheng S, Hedl M, Abraham C. Twist1 and Twist2 contribute to cytokine downregulation following chronic NOD2 stimulation of human macrophages through the coordinated regulation of transcriptional repressors and activators. *J Immunol* 2015;195:217-26.
47. Lo HYG, Jin RU, Sibbel G, Liu D, Karki A, et al. A single transcription factor is sufficient to induce and maintain secretory cell architecture. *Genes Dev* 2017;31:154-71.
48. Kaminska B, Mota M, Pizzi M. Signal transduction and epigenetic mechanisms in the control of microglia activation during neuroinflammation. *Biochim Biophys Acta* 2016;1862:339-51.
49. Nagatsu T, Sawada M. Inflammatory process in Parkinson's disease: role for cytokines. *Curr Pharm Des* 2005;11:999-1016.
50. Boka G, Anglade P, Wallach D, Javoy-Agid F, Agid Y, et al. Immunocytochemical analysis of tumor necrosis factor and its receptors in Parkinson's disease. *Neurosci Lett* 1994;172:151-4.
51. Mogi M, Harada M, Riederer P, Narabayashi H, Fujita K, et al. Tumor necrosis factor-alpha (TNF-alpha) increases both in the brain and in the cerebrospinal fluid from parkinsonian patients. *Neurosci Lett* 1994;165:208-10.
52. Qin XY, Zhang SP, Cao C, Loh YP, Cheng Y. Aberrations in peripheral inflammatory cytokine levels in Parkinson disease. *JAMA Neurol* 2016;73:1316-9.
53. Rizzo FR, Musella A, De Vito F, Fresegna D, Bullitta S, et al. Tumor necrosis factor and interleukin-1 β modulate synaptic plasticity during neuroinflammation. *Neural Plast* 2018;2018:1-12.
54. Stellwagen D, Malenka RC. Synaptic scaling mediated by glial TNF- α . *Nature* 2006;440:1054-9.
55. Takeuchi H, Jin S, Wang J, Zhang G, Kawanokuchi J, et al. Tumor necrosis factor-alpha induces neurotoxicity via glutamate release from hemichannels of activated microglia in an autocrine manner. *J Biol Chem* 2006;281:21362-8.
56. Ye L, Huang Y, Zhao L, Li Y, Sun L, et al. IL-1 β and TNF- α induce neurotoxicity through glutamate production: a potential role for neuronal glutaminase. *J Neurochem* 2013;125:897-908.
57. Denver P, McClean P. Distinguishing normal brain aging from the development of Alzheimer's disease: inflammation, insulin signaling and cognition. *Neural Regen Res* 2018;13:1719-12.
58. Njie eMalick G, Boelen E, Stassen FR, Steinbusch HWM, Borchelt DR, et al. Ex vivo cultures of microglia from young and aged rodent brain reveal age-related changes in microglial function. *Neurobiol Aging* 2012;33:195.e1-12.
59. Foster CT, Dovey OM, Lezina L, Luo JL, Gant TW, et al. Lysine-specific demethylase 1 regulates the embryonic transcriptome and CoREST stability. *Mol Cell Biol* 2010;30:4851-63.
60. Lee MG, Wynder C, Cooch N, Shiekhhattar R. An essential role for CoREST in nucleosomal histone 3 lysine 4 demethylation. *Nature* 2005;437:432-5.
61. Grabert K, Michoel T, Karavolos MH, Clohisey S, Baillie JK, et al. Microglial brain region-dependent diversity and selective regional sensitivities to aging. *Nat Neurosci* 2016;19:504-16.
62. Hammond TR, Dufort C, Dissing-Olesen L, Giera S, Young A, et al. Single-cell RNA sequencing of microglia throughout the mouse lifespan and in the injured brain reveals complex cell-state changes. *Immunity* 2019;50:253-71.e6.
63. Deczkowska A, Matcovitch-Natan O, Tsitsou-Kampeli A, Ben-Hamo S, Dvir-Szternfeld R, et al. Mef2C restrains microglial inflammatory response and is lost in brain ageing in an IFN-I-dependent manner. *Nat Commun* 2017;8:717.
64. The Tabula Muris Consortium, Pisco AO, McGeever A, Schaum N, Karkanias J, Neff NF, et al. A single cell transcriptomic atlas characterizes aging tissues in the mouse. *bioRxiv*; 2019. Available from: <https://doi.org/10.1101/661728> [Last accessed on 6 Mar 2020]
65. Deczkowska A, Amit I, Schwartz M. Microglial immune checkpoint mechanisms. *Nat Neurosci* 2018;21:779-86.
66. Readhead B, Haure-Mirande JV, Funk CC, Richards MA, Shannon P, et al. Multiscale analysis of independent Alzheimer's cohorts finds disruption of molecular, genetic, and clinical networks by human herpesvirus. *Neuron* 2018;99:64-82.e7.
67. Makin S. The amyloid hypothesis on trial. *Nature* 2018;559:S4-7.
68. Vijaya Kumar DK, Choi SH, Washicosky KJ, Eimer WA, Tucker S, et al. Amyloid- β peptide protects against microbial infection in mouse and worm models of Alzheimer's disease. *Sci Transl Med* 2016;8:340ra72.
69. Gosztyla ML, Brothers HM, Robinson SR. Alzheimer's amyloid- β is an antimicrobial peptide: a review of the evidence. *J Alzheimers Dis* 2018;62:1495-506.

AUTHOR INSTRUCTIONS

1. Submission Overview

Before you decide to publish with us, please read the following items carefully and make sure that you are well aware of Editorial Policies and the following requirements.

1.1 Topic Suitability

The topic of the manuscript must fit the scope of the journal. Please refer to Aims and Scope for more information.

1.2 Open Access and Copyright

The journal adopts Gold Open Access publishing model since its establishment and has been distributing contents under Attribution 4.0 International License since October 2017, whereas Attribution-NonCommercial-ShareAlike 3.0 Unported had been adopted by then. Please make sure that you are well aware of these policies.

1.3 Publication Fees

Authors are required to pay Article Processing Charges of 360 US Dollars after the manuscript is officially accepted. For more details, please refer to Article Processing Charges.

1.4 Language Editing

All submissions are required to be presented clearly and cohesively in good English. Authors whose first language is not English are advised to have their manuscripts checked or edited by a native English speaker before submission to ensure the high quality of expression. A well-organized manuscript in good English would make the peer review even the whole editorial handling more smooth and efficient.

If needed, authors are recommended to consider the language editing services provided by Charlesworth to ensure that the manuscript is written in correct scientific English before submission. Authors who publish with OAE journals enjoy a special discount for the services of Charlesworth via the following two ways.

Submit your manuscripts directly at <http://www.charlesworthauthorservices.com/~OAE>;

Open the link <http://www.charlesworthauthorservices.com/>, and enter Promotion Code “OAE” when you submit.

1.5 Work Funded by the National Institutes of Health

If an accepted manuscript was funded by National Institutes of Health (NIH), the author may inform editors of the NIH funding number. The editors are able to deposit the paper to the NIH Manuscript Submission System on behalf of the author.

2. Submission Preparation

2.1 Cover Letter

A cover letter is required to be submitted accompanying each manuscript. It should be concise and explain why the study is significant, why it fits the scope of the journal, and why it would be attractive to readers, etc.

Here is a guideline of a cover letter for authors' consideration:

In the first paragraph: include the title and type (e.g., Original Article, Review, Case Report, etc.) of the manuscript, a brief on the background of the study, the question the author sought out to answer and why;

In the second paragraph: concisely explain what was done, the main findings and why they are significant;

In the third paragraph: indicate why the manuscript fits the Aims and Scope of the journal, and why it would be attractive to readers;

In the fourth paragraph: confirm that the manuscript has not been published elsewhere and not under consideration of any other journal. All authors have approved the manuscript and agreed on its submission to the journal. Journal's specific requirements have been met if any.

If the manuscript is contributed to a special issue, please also mention it in the cover letter.

If the manuscript was presented partly or entirely in a conference, the author should clearly state the background information of the event, including the conference name, time and place in the cover letter.

2.2 Types of Manuscripts

There is no restriction on the length of manuscripts, number of figures, tables and references, provided that the manuscript is concise and comprehensive. The journal publishes Original Article, Review, Meta-Analysis, Case Report, Commentary, etc. For more details about paper type, please refer to the following table.

Manuscript Type	Definition	Abstract	Keywords	Main Text Structure
-----------------	------------	----------	----------	---------------------

Original Article	An Original Article describes detailed results from novel research. All findings are extensively discussed.	Structured abstract including Aim, Methods, Results and Conclusion. No more than 250 words.	3-8 keywords	The main content should include four sections: Introduction, Methods, Results and Discussion.
Review	A Review paper summarizes the literature on previous studies. It usually does not present any new information on a subject.	Unstructured abstract. No more than 250 words.	3-8 keywords	The main text may consist of several sections with unfixed section titles. We suggest that the author includes an "Introduction" section at the beginning, several sections with unfixed titles in the middle part, and a "Conclusion" section in the end.
Case Report	A Case Report details symptoms, signs, diagnosis, treatment, and follows up an individual patient. The goal of a Case Report is to make other researchers aware of the possibility that a specific phenomenon might occur.	Unstructured abstract. No more than 150 words.	3-8 keywords	The main text consists of three sections with fixed section titles: Introduction, Case Report, and Discussion.
Meta-Analysis	A Meta-Analysis is a statistical analysis combining the results of multiple scientific studies. It is often an overview of clinical trials.	Structured abstract including Aim, Methods, Results and Conclusion. No more than 250 words.	3-8 keywords	The main content should include four sections: Introduction, Methods, Results and Discussion.
Systematic Review	A Systematic Review collects and critically analyzes multiple research studies, using methods selected before one or more research questions are formulated, and then finding and analyzing related studies and answering those questions in a structured methodology.	Structured abstract including Aim, Methods, Results and Conclusion. No more than 250 words.	3-8 keywords	The main content should include four sections: Introduction, Methods, Results and Discussion.
Technical Note	A Technical Note is a short article giving a brief description of a specific development, technique or procedure, or it may describe a modification of an existing technique, procedure or device applied in research.	Unstructured abstract. No more than 250 words.	3-8 keywords	/
Commentary	A Commentary is to provide comments on a newly published article or an alternative viewpoint on a certain topic.	Unstructured abstract. No more than 250 words.	3-8 keywords	/
Editorial	An Editorial is a short article describing news about the journal or opinions of senior editors or the publisher.	None required	None required	/
Letter to Editor	A Letter to Editor is usually an open post-publication review of a paper from its readers, often critical of some aspect of a published paper. Controversial papers often attract numerous Letters to Editor	Unstructured abstract (optional). No more than 250 words.	3-8 keywords (optional)	/
Opinion	An Opinion usually presents personal thoughts, beliefs, or feelings on a topic.	Unstructured abstract (optional). No more than 250 words.	3-8 keywords	/
Perspective	A Perspective provides personal points of view on the state-of-the-art of a specific area of knowledge and its future prospects. Links to areas of intense current research focus can also be made. The emphasis should be on a personal assessment rather than a comprehensive, critical review. However, comments should be put into the context of existing literature. Perspectives are usually invited by the Editors.	Unstructured abstract. No more than 150 words.	3-8 keywords	/

2.3 Manuscript Structure

2.3.1 Front Matter

2.3.1.1 Title

The title of the manuscript should be concise, specific and relevant, with no more than 16 words if possible. When gene or

protein names are included, the abbreviated name rather than full name should be used.

2.3.1.2 Authors and Affiliations

Authors' full names should be listed. The initials of middle names can be provided. Institutional addresses and email addresses for all authors should be listed. At least one author should be designated as corresponding author. In addition, corresponding authors are suggested to provide their Open Researcher and Contributor ID upon submission. Please note that any change to authorship is not allowed after manuscript acceptance.

2.3.1.3 Abstract

The abstract should be a single paragraph with word limitation and specific structure requirements (for more details please refer to Types of Manuscripts). It usually describes the main objective(s) of the study, explains how the study was done, including any model organisms used, without methodological detail, and summarizes the most important results and their significance. The abstract must be an objective representation of the study: it is not allowed to contain results which are not presented and substantiated in the manuscript, or exaggerate the main conclusions. Citations should not be included in the abstract.

2.3.1.4 Keywords

Three to eight keywords should be provided, which are specific to the article, yet reasonably common within the subject discipline.

2.3.2 Main Text

Manuscripts of different types are structured with different sections of content. Please refer to Types of Manuscripts to make sure which sections should be included in the manuscripts.

2.3.2.1 Introduction

The introduction should contain background that puts the manuscript into context, allow readers to understand why the study is important, include a brief review of key literature, and conclude with a brief statement of the overall aim of the work and a comment about whether that aim was achieved. Relevant controversies or disagreements in the field should be introduced as well.

2.3.2.2 Methods

Methods should contain sufficient details to allow others to fully replicate the study. New methods and protocols should be described in detail while well-established methods can be briefly described or appropriately cited. Experimental participants selected, the drugs and chemicals used, the statistical methods taken, and the computer software used should be identified precisely. Statistical terms, abbreviations, and all symbols used should be defined clearly. Protocol documents for clinical trials, observational studies, and other non-laboratory investigations may be uploaded as supplementary materials.

2.3.2.3 Results

This section contains the findings of the study. Results of statistical analysis should also be included either as text or as tables or figures if appropriate. Authors should emphasize and summarize only the most important observations. Data on all primary and secondary outcomes identified in the section Methods should also be provided. Extra or supplementary materials and technical details can be placed in supplementary documents.

2.3.2.4 Discussion

This section should discuss the implications of the findings in context of existing research and highlight limitations of the study. Future research directions may also be mentioned.

2.3.2.5 Conclusion

It should state clearly the main conclusions and include the explanation of their relevance or importance to the field.

2.3.3 Back Matter

2.3.3.1 Acknowledgments

Anyone who contributed towards the article but does not meet the criteria for authorship, including those who provided professional writing services or materials, should be acknowledged. Authors should obtain permission to acknowledge from all those mentioned in the Acknowledgments section. This section is not added if the author does not have anyone to acknowledge.

2.3.3.2 Authors' Contributions

Each author is expected to have made substantial contributions to the conception or design of the work, or the acquisition, analysis, or interpretation of data, or the creation of new software used in the work, or have drafted the work or substantively revised it.

Please use Surname and Initial of Forename to refer to an author's contribution. For example: made substantial contributions

to conception and design of the study and performed data analysis and interpretation: Salas H, Castaneda WV; performed data acquisition, as well as provided administrative, technical, and material support: Castillo N, Young V. If an article is single-authored, please include “The author contributed solely to the article.” in this section.

2.3.3.3 Availability of Data and Materials

In order to maintain the integrity, transparency and reproducibility of research records, authors should include this section in their manuscripts, detailing where the data supporting their findings can be found. Data can be deposited into data repositories or published as supplementary information in the journal. Authors who cannot share their data should state that the data will not be shared and explain it. If a manuscript does not involve such issue, please state “Not applicable.” in this section.

2.3.3.4 Financial Support and Sponsorship

All sources of funding for the study reported should be declared. The role of the funding body in the experiment design, collection, analysis and interpretation of data, and writing of the manuscript should be declared. Any relevant grant numbers and the link of funder’s website should be provided if any. If the study is not involved with this issue, state “None.” in this section.

2.3.3.5 Conflicts of Interest

Authors must declare any potential conflicts of interest that may be perceived as inappropriately influencing the representation or interpretation of reported research results. If there are no conflicts of interest, please state “All authors declared that there are no conflicts of interest.” in this section. Some authors may be bound by confidentiality agreements. In such cases, in place of itemized disclosures, we will require authors to state “All authors declare that they are bound by confidentiality agreements that prevent them from disclosing their conflicts of interest in this work.” If authors are unsure whether conflicts of interest exist, please refer to the “Conflicts of Interest” of OAE Editorial Policies for a full explanation.

2.3.3.6 Ethical Approval and Consent to Participate

Research involving human subjects, human material or human data must be performed in accordance with the Declaration of Helsinki and approved by an appropriate ethics committee. An informed consent to participate in the study should also be obtained from participants, or their parents or legal guardians for children under 16. A statement detailing the name of the ethics committee (including the reference number where appropriate) and the informed consent obtained must appear in the manuscripts reporting such research.

Studies involving animals and cell lines must include a statement on ethical approval. More information is available at Editorial Policies.

If the manuscript does not involve such issue, please state “Not applicable.” in this section.

2.3.3.7 Consent for Publication

Manuscripts containing individual details, images or videos, must obtain consent for publication from that person, or in the case of children, their parents or legal guardians. If the person has died, consent for publication must be obtained from the next of kin of the participant. Manuscripts must include a statement that a written informed consent for publication was obtained. Authors do not have to submit such content accompanying the manuscript. However, these documents must be available if requested. If the manuscript does not involve this issue, state “Not applicable.” in this section.

2.3.3.8 Copyright

Authors retain copyright of their works through a Creative Commons Attribution 4.0 International License that clearly states how readers can copy, distribute, and use their attributed research, free of charge. A declaration “© The Author(s) 2020” will be added to each article. Authors are required to sign License to Publish before formal publication.

2.3.3.9 References

References should be numbered in order of appearance at the end of manuscripts. In the text, reference numbers should be placed in square brackets and the corresponding references are cited thereafter. Only the first five authors’ names are required to be listed in the references, other authors’ names should be omitted and replaced with “et al.”. Abbreviations of the journals should be provided on the basis of Index Medicus. Information from manuscripts accepted but not published should be cited in the text as “Unpublished material” with written permission from the source.

References should be described as follows, depending on the types of works:

Types	Examples
Journal articles by individual authors	Weaver DL, Ashikaga T, Krag DN, Skelly JM, Anderson SJ, et al. Effect of occult metastases on survival in node-negative breast cancer. <i>N Engl J Med</i> 2011;364:412-21. [PMID: 21247310 DOI: 10.1056/NEJMoa1008108]
Organization as author	Diabetes Prevention Program Research Group. Hypertension, insulin, and proinsulin in participants with impaired glucose tolerance. <i>Hypertension</i> 2002;40:679-86. [PMID: 12411462]

Both personal authors and organization as author	Vallancien G, Emberton M, Harving N, van Moorselaar RJ; Alf-One Study Group. Sexual dysfunction in 1,274 European men suffering from lower urinary tract symptoms. <i>J Urol</i> 2003;169:2257-61. [PMID: 12771764 DOI: 10.1097/01.ju.0000067940.76090.73]
Journal articles not in English	Zhang X, Xiong H, Ji TY, Zhang YH, Wang Y. Case report of anti-N-methyl-D-aspartate receptor encephalitis in child. <i>J Appl Clin Pediatr</i> 2012;27:1903-7. (in Chinese)
Journal articles ahead of print	Odibo AO. Falling stillbirth and neonatal mortality rates in twin gestation: not a reason for complacency. <i>BJOG</i> 2018; Epub ahead of print [PMID: 30461178 DOI: 10.1111/1471-0528.15541]
Books	Sherlock S, Dooley J. Diseases of the liver and billiary system. 9th ed. Oxford: Blackwell Sci Pub; 1993. pp. 258-96.
Book chapters	Meltzer PS, Kallioniemi A, Trent JM. Chromosome alterations in human solid tumors. In: Vogelstein B, Kinzler KW, editors. The genetic basis of human cancer. New York: McGraw-Hill; 2002. pp. 93-113.
Online resource	FDA News Release. FDA approval brings first gene therapy to the United States. Available from: https://www.fda.gov/NewsEvents/Newsroom/PressAnnouncements/ucm574058.htm . [Last accessed on 30 Oct 2017]
Conference proceedings	Harnden P, Joffe JK, Jones WG, editors. Germ cell tumours V. Proceedings of the 5th Germ Cell Tumour Conference; 2001 Sep 13-15; Leeds, UK. New York: Springer; 2002.
Conference paper	Christensen S, Oppacher F. An analysis of Koza's computational effort statistic for genetic programming. In: Foster JA, Lutton E, Miller J, Ryan C, Tettamanzi AG, editors. Genetic programming. EuroGP 2002: Proceedings of the 5th European Conference on Genetic Programming; 2002 Apr 3-5; Kinsdale, Ireland. Berlin: Springer; 2002. pp. 182-91.
Unpublished material	Tian D, Araki H, Stahl E, Bergelson J, Kreitman M. Signature of balancing selection in Arabidopsis. <i>Proc Natl Acad Sci U S A</i> . Forthcoming 2002.

For other types of references, please refer to U.S. National Library of Medicine.

The journal also recommends that authors prepare references with a bibliography software package, such as EndNote to avoid typing mistakes and duplicated references.

2.3.3.10 Supplementary Materials

Additional data and information can be uploaded as Supplementary Material to accompany the manuscripts. The supplementary materials will also be available to the referees as part of the peer-review process. Any file format is acceptable, such as data sheet (word, excel, csv, cdx, fasta, pdf or zip files), presentation (powerpoint, pdf or zip files), image (cdx, eps, jpeg, pdf, png or tiff), table (word, excel, csv or pdf), audio (mp3, wav or wma) or video (avi, divx, flv, mov, mp4, mpeg, mpg or wmv). All information should be clearly presented. Supplementary materials should be cited in the main text in numeric order (e.g., Supplementary Figure 1, Supplementary Figure 2, Supplementary Table 1, Supplementary Table 2, *etc.*). The style of supplementary figures or tables complies with the same requirements on figures or tables in main text. Videos and audios should be prepared in English, and limited to a size of 500 MB or a duration of 3 minutes.

2.4 Manuscript Format

2.4.1 File Format

Manuscript files can be in DOC and DOCX formats and should not be locked or protected.

2.4.2 Length

There are no restrictions on paper length, number of figures, or amount of supporting documents. Authors are encouraged to present and discuss their findings concisely.

2.4.3 Language

Manuscripts must be written in English.

2.4.4 Multimedia Files

The journal supports manuscripts with multimedia files. The requirements are listed as follows:

Videos or audio files are only acceptable in English. The presentation and introduction should be easy to understand. The frames should be clear, and the speech speed should be moderate.

A brief overview of the video or audio files should be given in the manuscript text.

The video or audio files should be limited to a duration of 3 min and a size of up to 500 MB.

Please use professional software to produce high-quality video files, to facilitate acceptance and publication along with the submitted article. Upload the videos in mp4, wmv, or rm format (preferably mp4) and audio files in mp3 or wav format.

2.4.5 Figures

Figures should be cited in numeric order (e.g., Figure 1, Figure 2) and placed after the paragraph where it is first cited;

Figures can be submitted in format of tiff, psd, AI or jpeg, with resolution of 300-600 dpi;

Figure caption is placed under the Figure;

Diagrams with describing words (including, flow chart, coordinate diagram, bar chart, line chart, and scatter diagram, etc.) should be editable in word, excel or powerpoint format. Non-English information should be avoided; Labels, numbers, letters, arrows, and symbols in figure should be clear, of uniform size, and contrast with the background; Symbols, arrows, numbers, or letters used to identify parts of the illustrations must be identified and explained in the legend;

Internal scale (magnification) should be explained and the staining method in photomicrographs should be identified;

All non-standard abbreviations should be explained in the legend;

Permission for use of copyrighted materials from other sources, including re-published, adapted, modified, or partial figures and images from the internet, must be obtained. It is authors' responsibility to acquire the licenses, to follow any citation instruction requested by third-party rights holders, and cover any supplementary charges.

2.4.6 Tables

Tables should be cited in numeric order and placed after the paragraph where it is first cited;

The table caption should be placed above the table and labeled sequentially (e.g., Table 1, Table 2);

Tables should be provided in editable form like DOC or DOCX format (picture is not allowed);

Abbreviations and symbols used in table should be explained in footnote;

Explanatory matter should also be placed in footnotes;

Permission for use of copyrighted materials from other sources, including re-published, adapted, modified, or partial tables from the internet, must be obtained. It is authors' responsibility to acquire the licenses, to follow any citation instruction requested by third-party rights holders, and cover any supplementary charges.

2.4.7 Abbreviations

Abbreviations should be defined upon first appearance in the abstract, main text, and in figure or table captions and used consistently thereafter. Non-standard abbreviations are not allowed unless they appear at least three times in the text. Commonly-used abbreviations, such as DNA, RNA, ATP, *etc.*, can be used directly without definition. Abbreviations in titles and keywords should be avoided, except for the ones which are widely used.

2.4.8 Italics

General italic words like *vs.*, *et al.*, *etc.*, *in vivo*, *in vitro*; *t* test, *F* test, *U* test; related coefficient as *r*, sample number as *n*, and probability as *P*; names of genes; names of bacteria and biology species in Latin.

2.4.9 Units

SI Units should be used. Imperial, US customary and other units should be converted to SI units whenever possible. There is a space between the number and the unit (i.e., 23 mL). Hour, minute, second should be written as h, min, s.

2.4.10 Numbers

Numbers appearing at the beginning of sentences should be expressed in English. When there are two or more numbers in a paragraph, they should be expressed as Arabic numerals; when there is only one number in a paragraph, number < 10 should be expressed in English and number > 10 should be expressed as Arabic numerals. 12345678 should be written as 12,345,678.

2.4.11 Equations

Equations should be editable and not appear in a picture format. Authors are advised to use either the Microsoft Equation Editor or the MathType for display and inline equations.

2.5 Submission Link

Submit an article via <https://oaemesas.com/nn>.



Neuroimmunology and Neuroinflammation (NN)

Los Angeles Office
245 E Main Street ste122, Alhambra,
CA 91801, USA
Tel: +1 323 9987086
E-mail: nn_editor001@nnjournal.net
Website: www.nnjournal.net

