

Research Article

Open Access



Enantioselective synthesis of 6-(Indole-2-yl)-3,4-dihydropyran-2-one skeletons by *N*-Heterocyclic carbene-catalyzed asymmetric [3 + 3] cycloaddition of α -bromocinnamaldehyde

Gao He^{1,#}, Xiaoyu Chen^{1,#}, Siqi Xia¹, Guofu Zhong^{2,*}, Limin Yang^{1,*} 

¹College of Material, Chemistry and Chemical Engineering, Hangzhou Normal University, Hangzhou 311121, Zhejiang, China.

²Department of Chemistry, Eastern Institute for Advanced Study, Ningbo 315200, Zhejiang, China.

#Authors contributed equally.

*Correspondence to: Prof. Guofu Zhong, Department of Chemistry, Eastern Institute of Technology, No.568 Tongxin Road, Ningbo 315200, Zhejiang, China. E-mail: guofuzhong@eias.ac.cn; Prof. Limin Yang, College of Material, Chemistry and Chemical Engineering, Hangzhou Normal University, No.2318 Yuhangtang Road, Hangzhou 311121, Zhejiang, China. E-mail: myang@hznu.edu.cn;

How to cite this article: He G, Chen X, Xia S, Zhong G, Yang L. Enantioselective synthesis of 6-(Indole-2-yl)-3,4-dihydropyran-2-one skeletons by *N*-Heterocyclic carbene-catalyzed asymmetric [3 + 3] cycloaddition of α -bromocinnamaldehyde. *Chem Synth* 2023;3:35. <https://dx.doi.org/10.20517/cs.2023.14>

Received: 02 Mar 2023 **First Decision:** 28 Mar 2023 **Revised:** 29 Apr 2023 **Accepted:** 16 May 2023 **Published:** 21 Jul 2023

Academic Editors: Bao-Lian Su, Feng Shi **Copy Editor:** Dan Zhang **Production Editor:** Dan Zhang

Abstract

The enantioselective construction of chiral 6-(indole-2-yl)-3,4-dihydropyran-2-one skeleton was demonstrated by the formal [3 + 3] cycloaddition reaction of α -bromocinnamaldehyde with β -ketoester indole catalyzed by chiral *N*-heterocyclic carbene (NHC). The reaction proceeds smoothly via a vinyl acyl azolium intermediate (electron-poor enone) generated from NHC-aldehyde adducts, providing 6-(indole-2-yl)-3,4-dihydropyran-2-one derivatives in good yields with excellent enantioselectivities (up to 98% ee).

Keywords: Organocatalysis, enantioselective synthesis, *N*-heterocyclic carbene (NHC), acyl azolium intermediate, 6-(indole-2-yl)-3, 4-dihydropyran-2-one skeleton

INTRODUCTION

Indole skeletons are widely present in many important biomolecules and possess a heterocyclic structure with diverse properties. Their structural characteristics can offer unique features, such as modifying electron



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, sharing, adaptation, distribution and reproduction in any medium or format, for any purpose, even commercially, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.



cloud density, adjusting steric hindrance, and realizing post-functionalization. In recent years, asymmetric catalytic reactions based on indole skeletons have emerged as a new research field^[1-9].

Indole-2-yl-pyrans are the basic structural framework for numerous biologically active natural products and drugs. Two novel alkaloids isolated from plant roots have an indole-C-glucoside core^[10]. Indole-C-glucopyranoside **A** showed cytotoxic activity against human myeloid leukemia HL60 cells (IC₅₀ = 1.3 mM) and human hepatocellular carcinoma HepG2 cells. Isomer **B** showed cytotoxic activity against both HL60 cells and human myeloid leukemia Mata cells (IC₅₀) [Figure 1].

Given the importance of the indole skeleton, the selective functionalization of this skeleton catalyzed by *N*-heterocyclic carbene (NHC) has been gaining increasing interest in recent years. Armido Studer and other chemists reported intra- and intermolecular cycloaddition by NHC catalysis to construct spirocyclic indole skeletons by introducing formyl benzyl groups, -CHO, -OH, alkenyl groups, and other functional groups into the indole structure^[11-19]. Balanna *et al.* reported the NHC-catalyzed construction of the indole skeleton by introducing -CHO, -CH₃, -NO₂, and other groups into the indole structure^[20-23]. Liu *et al.* also reported the NHC-catalyzed construction of polycyclic substrates by designing the indole skeleton^[24-27]. Du *et al.* studied the construction of an axially chiral indole skeleton^[28]. In addition, Chi *et al.* and Gong *et al.* developed acyclic aldol reactions of the indole skeleton to form quaternary stereogenic centers by post-aldol stereochemistry control^[29,30].

Although the NHC-catalyzed synthesis of indole derivatives is widely studied, the application of NHC catalysis for the synthesis of indole-pyran skeletons has received limited attention. In 2013, Chi^[31] *et al.* developed the direct β -carbon functionalization of saturated aldehydes through oxidative NHC catalysis, leading to dihydropyranone with high enantioselectivity [Scheme 1A]. Subsequently, Sundén *et al.* used O₂ instead of high molecular weight stoichiometric oxidants and introduced a system of electron transfer mediators (ETMs) to realize β -carbon functionalization of unsaturated aldehydes [Scheme 1B]^[32]. Although preliminary achievements have been made in the construction of the dihydropyranone skeleton, the avoidance of any oxidants and additives remains a challenge. Our research group has reported the construction of spiroindole skeleton compounds catalyzed by NHC^[12]. Our continuous interest in NHC-catalyzed cycloaddition inspired us to construct chiral 6-(indole-2-yl)-3,4-dihydropyran-2-one derivatives by NHC-catalyzed [3 + 3] cycloaddition reactions between β -ketoester indole and α -bromocinnamaldehydes [Scheme 1C].

EXPERIMENTAL

To an oven-dried 10 mL vial, β -ketoester indole **1** (0.1 mmol, 1.0 equiv), β -bromo- α,β -unsaturated aldehyde **2** (0.1 mmol, 1.2 equiv), cat. **A** (7.4 mg, 0.02 mmol, 0.2 equiv), NaHCO₃ (12.6 mg, 0.15 mmol, 1.5 equiv) were added, followed by 1.5 mL of THF. The mixture was stirred overnight at room temperature. Once the reaction was completed (monitored by TLC), the desired product **3** was purified by silica gel column chromatography with EA/PE (1:10) as an eluent.

RESULTS AND DISCUSSION

In view of the fact that some enantioenriched indole and pyran derivatives are pharmaceutically attractive compounds, we chose β -ketoester indole **1a** and α -bromocinnamaldehyde **2a** as model reaction substrates to optimize the reaction conditions [Figure 2]. After evaluating triazolium pre-catalysts A-C, we found that pre-catalyst A gave the product **3a** in 67% yield and 45% *ee*, while pre-catalyst C, bearing C₆F₅ substituents, did not provide the desired product. Installing a NO₂ group to the indane moiety of pre-catalyst A (to get pre-catalyst B) led to a small improvement on the reaction yield but with a decrease in enantioselectivity






Figure 1. Representative natural products and bioactive compounds containing the indole-2-yl-pyrans skeleton.


A) direct β -activation of saturated aldehydes to synthesize dihydropyranones (Chi's work)



B) asymmetric aerobic oxidation assisted by electron transfer mediators (Sundén's work)



C) enantioselective synthesis of 6-(indole-2-yl)-3,4-dihydropyran-2-one skeletons (this work)



Scheme 1. Profile of the construction of NHC-catalyzed construction of dihydropyranone.

(entry 2, 72% yield, 35% *ee*). Various inorganic bases could be used in this transformation (entries 4-5), and NaHCO_3 was found as the most efficient one that could give **3a** with good yield and excellent enantioselectivity (entry 5, 75% yield, 94% *ee*). Screening different reaction solvents did not show further improvements in either yields or enantioselectivities (entries 6-7). When 5 Å MS was used as the additive, the yield and the enantioselectivity increased slightly higher (entry 8, 78% yield, 97% *ee*). Further





Figure 2. Optimization of the reaction conditions^a. ^aReaction conditions: pre-catalyst (20 mol%), base (1.5 equiv), **1a** (0.1 mmol), and **2a** (0.12 mmol) in solvent (1.5 mL). ^bIsolated yield. ^cDetermined by chiral HPLC analysis. ^d5Å molecular sieves (10 mg) were used as an additive at room temperature. ^epre-catalyst (10 mol%). ^fThe temperature was 0 °C. ^gThe temperature was 35 °C. ^hTHF = 3 mL. ⁱTHF = 1 mL. Mes = 2,4,6-trimethylphenyl.

optimization studies revealed that high enantioselectivity with low reaction yield (entry 9, 68% yield, 97% *ee*) was delivered when the reaction was performed with 10 mol% pre-catalyst A. Then, the reaction temperature was investigated. When the reaction proceeded at 0 °C, the yield was dramatically decreased to 8%, and the enantioselectivity was slightly decreased to 92% [Figure 2] (entry 10). Higher reaction temperature did not result in better yield or enantioselectivity [Figure 2] (entry 11). While the changing of reactant concentration led to lower yields (entries 12-13). The best reaction condition was established with a 20 mol% pre-catalyst, 1.5 equiv NaHCO₃, and 1.0:1.2 reactant ratio at room temperature.

After establishing the optimized reaction conditions, the scope of the reaction was examined. First, we studied variations of the β-ketoester indoles [Scheme 2]. Substrates with electron-releasing and electron-withdrawing groups on the benzene ring of the indole group of 1,3-dicarbonyl compound (R²) underwent a cycloaddition reaction, affording the compounds in moderate to good yields (61%-80% yield) and excellent enantioselectivities (**3a–3j**, 92%-98% *ee*). It is worth mentioning that the enantioselectivity of the reaction can reach 98% when the C5 position of the indol-phenyls is NO₂. Subsequently, we investigated different N-protecting groups on the indole skeleton, such as Me, PMB, or tert-butyl benzyl. To our great delight, β-



Scheme 2. Substrates scope. Reaction conditions: pre-cat. **A** (20 mol%), NaHCO_3 (1.5 equiv), **1** (0.1 mmol), and **2** (0.12 mmol) in THF (1.5 mL).

ketoester indoles with different N-protecting groups also worked well in this process under the current reaction condition, and the desired products were generated in good to excellent yields and optical purities (**3k-3m**, 97%-98% ee). However, the N-electron withdrawing protecting group (such as Boc) showed lower reaction activity and enantioselectivities. Furthermore, the enantioselectivity of the reaction has a slight decrease without an N-protecting group (**3n**, 87% ee). This may be due to the steric resistance effect of the N-protecting group. Encouraged by these results, the generality of α -bromocinnamaldehyde **2** was further investigated with β -ketoester indole **1a** [Scheme 2]. Substituents with different electronic properties at the orthoposition (**3o**, 65% yield, 95% ee) and the paraposition (**3p-3r**, 93-97% ee) of the phenyl group were well tolerated. Para-F on the phenyl ring led to some decrease in reaction yield (**3p**, 50% yield). Replacement of the phenyl substituent with 2-furanyl had little effect on the reaction outcomes (**3s**, 68% yield, 83% ee). In addition, trace desired products were obtained by employing alkyl-substituted aliphatic aldehydes.




Figure 3. X-ray single crystal data for compound 3t.


To demonstrate the practical nature of the present catalytic asymmetric strategy, a gram-scale preparation of compound 3a was performed [Scheme 3]. Under the standard reaction conditions, amplifying the model reaction to 1.00 mmol straightforwardly gave optically pure 6-(indole-2-yl)-3,4-dihydropyran-2-one 3a without the loss of enantioselectivity in modest yield (56% yield and 96% *ee*).

The absolute configuration of the cycloadduct 3t was determined by X-ray crystallographic analysis as an (*S*)-configuration [Figure 3]. The absolute configuration of all other products was assigned accordingly (CCDC 2243010, please see the SI for details).


The plausible catalytic cyclic for this [3 + 3] cycloaddition pathway is shown in Scheme 4. The cycloaddition reaction begins with the addition of NHC to α -bromocinnamaldehyde 2a, forming a Breslow intermediate, which undergoes tautomerization and debromination to generate form α , β -unsaturated acylazolium III. The Michael addition of enolate generated by β -ketoester indole 1a to intermediate III led to intermediate IV. Intermediate IV undergoes tautomerization and intramolecular esterification to give the desired product 3a and regenerate the NHC catalyst.

CONCLUSIONS

In conclusion, we have developed a mild NHC-catalyzed efficient [3 + 3] cycloaddition reaction between α -bromocinnamaldehyde and β -ketoester indole, avoiding the use of high molecular weight stoichiometric oxidants. This reaction resulted in the highly enantioselective construction of 6-(indole-2-yl)-3,4-dihydropyran-2-one skeleton in good yields and exhibiting excellent enantioselectivities.



Scheme 3. A gram scale reaction.



Scheme 4. Proposed catalytic pathway and transition state.

DECLARATIONS

Authors' contributions

Designing the experiments, writing the manuscript, and being responsible for the whole work: Yang L
 Performing the experiments: He G

Manuscript writing and picture drawing, [Supplementary Materials](#): Chen X

Synthesizing the substrates and data review: Xia S

Co-directing this project: Zhong G

Directing this project and revising the manuscript: Yang L

Availability of data and materials

Not applicable.

Financial support and sponsorship

We gratefully acknowledge the National Natural Science Foundation of China (Grant 22271071); the Natural Science Foundation of Zhejiang Province (Grant LY20B020010); and the Foundation of Shanghai Key Laboratory for Molecular Engineering of Chiral Drugs for financial support (Grant SMECD202300X).

Conflicts of interest

All authors declared that there are no conflicts of interest.

Ethical approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Copyright

© The Author(s) 2023.

REFERENCES

1. Norton RS, Wells RJ. A series of chiral polybrominated biindoles from the marine blue-green alga rivularia firma. application of carbon-13 NMR spin-lattice relaxation data and carbon-13-proton coupling constants to structure elucidation. *J Am Chem Soc* 1982;104:3628-35. [DOI](#)
2. Jiang F, Chen KW, Wu P, Zhang YC, Jiao Y, Shi F. A strategy for synthesizing axially chiral naphthyl-indoles: catalytic asymmetric addition reactions of racemic substrates. *Angew Chem Int Ed Engl* 2019;58:15104-10. [DOI](#)
3. Wang C, Li T, Liu S, et al. Axially chiral aryl-alkene-indole framework: a nascent member of the atropisomeric family and its catalytic asymmetric construction. *Chin J Chem* 2020;38:543-52. [DOI](#)
4. Chen KW, Chen ZH, Yang S, Wu SF, Zhang YC, Shi F. Organocatalytic atroposelective synthesis of N-N Axially chiral indoles and pyrroles by de novo ring formation. *Angew Chem Int Ed Engl* 2022;61:e202116829. [DOI](#)
5. Yang Y, Xu Y, Yue Y, et al. Investigate natural product indolmycin and the synthetically improved analogue toward antimycobacterial agents. *ACS Chem Biol* 2022;17:39-53. [DOI](#)
6. Humphrey GR, Kuethe JT. Practical methodologies for the synthesis of indoles. *Chem Rev* 2006;106:2875-911. [DOI](#) [PubMed](#)
7. Bandini M, Eichholzer A. Catalytic functionalization of indoles in a new dimension. *Angew Chem Int Ed Engl* 2009;48:9608-44. [DOI](#)
8. Kochanowska-Karamyan AJ, Hamann MT. Marine indole alkaloids: potential new drug leads for the control of depression and anxiety. *Chem Rev* 2010;110:4489-97. [DOI](#) [PubMed](#) [PMC](#)
9. Sun C, Tian W, Lin Z, Qu X. Biosynthesis of pyrroloindoline-containing natural products. *Nat Prod Rep* 2022;39:1721-65. [DOI](#) [PubMed](#)
10. Yepremyan A, Minehan TG. Total synthesis of indole-3-acetonitrile-4-methoxy-2-C-β-D-glucopyranoside. Proposal for structural revision of the natural product. *Org Biomol Chem* 2012;10:5194-6. [DOI](#) [PubMed](#) [PMC](#)
11. Bera S, Daniliuc CG, Studer A. Oxidative N-heterocyclic carbene catalyzed dearomatization of indoles to spirocyclic indolenines with a quaternary carbon stereocenter. *Angew Chem Int Ed Engl* 2017;56:7402-6. [DOI](#) [PubMed](#)
12. Xie D, Yang L, Lin Y, et al. Rapid access to spirocyclic oxindoles: application of asymmetric N-heterocyclic carbene-catalyzed [3 + 3] cycloaddition of imines to oxindole-derived enals. *Org Lett* 2015;17:2318-21. [DOI](#)
13. Zhu SY, Zhang H, Ma QW, Liu D, Hui XP. Oxidative NHC catalysis: direct activation of β sp³ carbons of saturated acid chlorides. *Chem Commun* 2019;55:298-301. [DOI](#)
14. Xu J, Zhang W, Liu Y, et al. Formal [3 + 3] annulation of isatin-derived 2-bromoaldehydes with 1,3-dicarbonyl compounds enabled by Lewis acid/N-heterocyclic carbene cooperative catalysis. *RSC Adv* 2016;6:18601-6. [DOI](#)

15. Przydacz A, Topolska A, Skrzyńska A, Albrecht L. NHC-catalyzed 1,4-elimination in the dearomative activation of 3-furaldehydes towards (4 + 2)-cycloadditions. *Adv Synth Catal* 2022;364:1434-9. DOI
16. Zhang ZJ, Wen YH, Song J, Gong LZ. Kinetic resolution of aziridines enabled by N-heterocyclic carbene/copper cooperative catalysis: carbene dose-controlled chemo-switchability. *Angew Chem Int Ed Engl* 2021;60:3268-76. DOI PubMed
17. Rong X, Yao H, Xia W, Du Y, Zhou Y, Liu H. Enantioselective assembly of spirolactones through NHC-catalyzed remote γ -carbon addition of enals with isatins. *ACS Comb Sci* 2016;18:220-4. DOI PubMed
18. Yang X, Sun J, Huang X, Jin Z. Asymmetric synthesis of structurally sophisticated spirocyclic pyrano[2,3-c]pyrazole derivatives bearing a chiral quaternary carbon center. *Org Lett* 2022;24:5474-9. DOI
19. Breuers CBJ, Daniliuc CG, Studer A. Oxidative N-heterocyclic carbene-catalyzed intramolecular friedel-crafts alkylation of indoles for the synthesis of spirocyclic indolenines. *Org Lett* 2022;24:5314-8. DOI PubMed
20. Balanna K, Madica K, Mukherjee S, et al. N-heterocyclic carbene-catalyzed formal [6 + 2] annulation reaction via cross-conjugated Aza-trienolate intermediates. *Chemistry* 2020;26:818-22. DOI
21. Chen X, Yang S, Song BA, Chi YR. Functionalization of benzylic C(sp³)-H bonds of heteroaryl aldehydes through N-heterocyclic carbene organocatalysis. *Angew Chem Int Ed Engl* 2013;52:11134-7. DOI
22. Wu X, Zhou L, Maiti R, Mou C, Pan L, Chi YR. Sulfinate and carbene co-catalyzed rauhut-currier reaction for enantioselective access to azepino[1,2-a]indoles. *Angew Chem Int Ed Engl* 2019;58:477-81. DOI
23. Huang H, Li Q, Liu Y, et al. Dearomative [4 + 2] annulations between 3-nitroindoles and enals through oxidative N-heterocyclic carbene catalysis. *Org Chem Front* 2020;7:3862-7. DOI
24. Liu Y, Luo G, Yang X, et al. Carbene-catalyzed enantioselective aromatic N-nucleophilic addition of heteroarenes to ketones. *Angew Chem Int Ed Engl* 2020;59:442-8. DOI
25. Li W, Yuan H, Liu Z, Zhang Z, Cheng Y, Li P. NHC-catalyzed enantioselective [4 + 3] cycloaddition of *ortho*-hydroxyphenyl substituted *para*-quinone methides with isatin-derived enals. *Adv Synth Catal* 2018;360:2460-4. DOI
26. Mukherjee S, Shee S, Poisson T, Besset T, Biju AT. Enantioselective N-heterocyclic carbene-catalyzed cascade reaction for the synthesis of pyrroloquinolines via N-H functionalization of indoles. *Org Lett* 2018;20:6998-7002. DOI PubMed
27. Xu J, Yuan S, Miao M. N-Heterocyclic carbene catalyzed [4 + 2] annulation reactions with in situ generated heterocyclic ortho-quinodimethanes. *Org Lett* 2016;18:3822-5. DOI PubMed
28. Ma R, Wang X, Zhang Q, et al. Atroposelective synthesis of axially chiral 4-aryl α -carbolines via n-heterocyclic carbene catalysis. *Org Lett* 2021;23:4267-72. DOI
29. Yang X, Majhi PK, Chai H, et al. Carbene-catalyzed enantioselective aldol reaction: post-aldol stereochemistry control and formation of quaternary stereogenic centers. *Angew Chem Int Ed Engl* 2021;60:159-65. DOI
30. Fan T, Song J, Gong LZ. Asymmetric redox allylic alkylation to access 3,3'-disubstituted oxindoles enabled by Ni/NHC cooperative catalysis. *Angew Chem Int Ed Engl* 2022;61:e202201678. DOI PubMed
31. Mo J, Shen L, Chi YR. Direct β -activation of saturated aldehydes to formal Michael acceptors through oxidative NHC catalysis. *Angew Chem Int Ed Engl* 2013;52:8588-91. DOI PubMed
32. Axelsson A, Hammarvid E, Ta L, Sundén H. Asymmetric aerobic oxidative NHC-catalysed synthesis of dihydropyranones utilising a system of electron transfer mediators. *Chem Commun* 2016;52:11571-4. DOI PubMed