

Review

Open Access



# Emerging trends in robotic breast surgery in the era of artificial intelligence

Tingting Li, Chen Li

Department of Breast Surgery, Southwest Hospital, The First Affiliated Hospital of Army Medical University, Key Laboratory of Chongqing Health Commission for Minimally Invasive and Precise Diagnosis and Treatment of Breast Cancer, Chongqing 400038, China.

**Correspondence to:** Prof. Li Chen, Department of Breast Surgery, Southwest Hospital, The First Affiliated Hospital of Army Medical University, Key Laboratory of Chongqing Health Commission for Minimally Invasive and Precise Diagnosis and Treatment of Breast Cancer, No. 30 GaoTanYan Street, ShaPingBa District, Chongqing 400038, China. E-mail: chenli@tmmu.edu.cn

**How to cite this article:** Li T, Li C. Emerging trends in robotic breast surgery in the era of artificial intelligence. *Plast Aesthet Res.* 2025;12:5. <https://dx.doi.org/10.20517/2347-9264.2024.143>

**Received:** 8 Nov 2024 **First Decision:** 20 Jan 2025 **Revised:** 13 Feb 2025 **Accepted:** 4 Mar 2025 **Published:** 11 Mar 2025

**Academic Editor:** Raymund E. Horch **Copy Editor:** Ting-Ting Hu **Production Editor:** Ting-Ting Hu

## Abstract

The advent of artificial intelligence (AI) heralds a new era in the field of robotic surgery. This article discusses recent trends in the integration of AI technology with robotic surgical procedures, highlighting the latest advancements in robotic breast surgery. The application of AI in robotic surgery ranges from preoperative planning to intraoperative assistance. Machine learning algorithms are now utilized to analyze medical imaging data, enabling surgeons to devise detailed surgical plans tailored to the unique characteristics of each patient's tumor. This approach leads to more precise tumor excision and better preservation of healthy tissue. Robotic systems equipped with advanced visualization and sensor technologies can provide real-time feedback during surgery and training. Additionally, AI algorithms can predict the occurrence of postoperative complications, allowing for early intervention. With the ongoing development of AI and robotic technologies, significant progress has been made in robotic automation. The future of robotic breast surgery holds the promise of even greater accuracy, and the quality of life for breast cancer patients may be significantly improved.

**Keywords:** Artificial intelligence, robotic breast surgery, convolutional neural network, automation, deep learning



© The Author(s) 2025. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, sharing, adaptation, distribution and reproduction in any medium or format, for any purpose, even commercially, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.



## INTRODUCTION

Minimally invasive breast surgery is a popular surgical treatment method at present, which enables improved cosmetic outcomes<sup>[1,2]</sup>. Although endoscopic breast surgery is safe and technically feasible, it remains hindered by limited degrees of freedom, insufficient anatomical angles, and interference between instruments at a single axillary access point<sup>[3]</sup>. The robotic-assisted surgical system has obvious advantages in breast surgery due to the 3D visualization, enhanced flexibility, and reduction of intraoperative tremor, and the safety and feasibility of its adoption are currently under investigation in the surgical field<sup>[4,5]</sup>. However, robotic surgery also has disadvantages such as high training requirements for medical staff, lack of tactile feedback, long surgical preparation times, and technological limitations in complex surgeries<sup>[6]</sup>. The integration of artificial intelligence (AI) with robotic platforms could refine robotic surgery further. AI can combine computational power with human intelligence, thereby enabling deep learning<sup>[7]</sup>. AI technology can provide personalized treatment recommendations based on the individual health data of patients, and it can also serve as a platform to deliver remote care through telemedicine technology<sup>[8-11]</sup>. The data analysis and deep learning functions of AI are essential to achieve semi-automation and automation of robotic surgery, including preoperative planning and precision surgical planning, as well as real-time intraoperative decision support. The combination of AI and robotic surgery is constantly pushing surgical procedures in a more accurate and efficient direction and it could also lead to further advances in robotic breast surgery. This review highlights more recent technological advancements in robotic breast surgery in the era of AI.

## ROBOT-ASSISTED SURGERY

The concept of using robots to operate remotely was first proposed by the US Army and the National Aeronautics and Space Administration (NASA)<sup>[12]</sup>. In the 1990s, the first master-slave system was developed, which consisted of a robot with remote manipulators controlled by a surgeon at a surgical workstation. A widely used platform is the da Vinci robot, a master-slave device developed by Intuitive Surgical Inc<sup>[13]</sup>. The da Vinci device in human abdominal surgeries was approved by the Food and Drug Administration (FDA) in 2000, and its approval for breast surgery still requires more evaluation of oncological safety<sup>[14]</sup>. The technical advantages of robotic surgery, such as 3D vision, stable and magnified images, physiological tremor filtration, and motion scaling, were able to overcome many limitations of laparoscopic surgery<sup>[15,16]</sup>. Robotic surgery has been rapidly adopted and now plays a significant role in gastrointestinal, hepatobiliary, pancreatic, urological, and gynecological procedures<sup>[17-24]</sup>. In 2015, the Da Vinci robot surgical system was first reported to be used in breast surgery<sup>[25]</sup>. Other robotic surgical systems developed in recent years include the MUSA Surgical System and the Symani Surgical System, which have more operational flexibility due to the high degree of freedom in their wrist tools, primarily benefiting the field of microsurgery<sup>[26,27]</sup>. With the development of fifth-generation (5G) networks and AI, robotic surgery may offer new frontiers, such as enabling remote surgeries and the potential for the future to herald an era of autonomous robot-delivered surgeries<sup>[28,29]</sup>.

## ROBOTIC BREAST SURGERY

### Robotic nipple-sparing mastectomy

Both nipple-sparing mastectomy (NSM) and skin-sparing mastectomy (SSM) are considered standard modalities of performing a mastectomy<sup>[30-33]</sup>. A devastating complication associated with NSM is nipple-areolar complex (NAC) necrosis, and the rate of NAC necrosis ranges from 7.4% to 11% after NSM<sup>[34,35]</sup>. Intraoperative frozen section of subareolar tissue is commonly used to confirm negative margins, with atypical or positive frozen results leading to SSM, occurring in 2.5%-12% of cases<sup>[36,37]</sup>. NSM with immediate reconstruction is associated with enhanced patient satisfaction<sup>[38]</sup>. Endoscopic NSM has been proven feasible and is associated with high patient satisfaction<sup>[39]</sup>. The robot-assisted nipple-sparing mastectomy (RNSM)

was first described by Toesca *et al.* in 2015, and they performed NSM through a single axillary incision, followed by immediate implant reconstruction<sup>[25]</sup>. In Toesca's initial series of 29 cases with RNSM, only two cases required conversion to open surgery due to technical problems. The total surgery duration did not exceed 3 h, and there were no cases of hematoma, skin or NAC injury/necrosis, or infection<sup>[40]</sup>. To achieve a safe and efficient RNSM, it is essential to provide training in robotic surgery not only to the surgical staff but also to the entire team. A meta-analysis shows that compared to traditional NSM, RNSM does not show a significant difference in the incidence of postoperative complications<sup>[41]</sup>. Although the RNSM takes longer and is more costly, the clinical outcomes of RNSM were comparable to those of traditional NSM, and RNSM showed favorable satisfaction with esthetic results<sup>[42]</sup>. RNSM is a safe option in terms of complications, particularly with improved wound healing, and the advantage of the minimal access group in RNSM is the higher satisfaction than conventional NSM related to wound esthetics<sup>[43]</sup>. Although there is a paucity of data to make definitive conclusions, one study suggested that compared to endoscopic nipple-sparing mastectomy (ENSM), RNSM offers higher satisfaction with wound esthetic, less bleeding, and a shorter learning curve<sup>[44]</sup>. The cost of RNSM is significantly higher than ENSM, which is one of the limitations of RNSM at present, and more research is needed to evaluate its cost-effectiveness<sup>[43,44]</sup>. Relevant comparative studies are shown in [Table 1](#). The inherent limitations of the multi-arm surgical platform restrict the ergonomic use of the mechanical arms and the mobility of the robotic camera. In 2018, the FDA approved the da Vinci SP<sup>®</sup> surgical system for urological surgery, marking another significant step forward in the evolution of robotic surgery<sup>[45]</sup>. The SP<sup>®</sup> surgical system, designed specifically for single-port (SP) procedures, utilizes a single 25 mm multi-channel port (which includes a 3D robotic camera and three instrument slots), offering more flexible operation and superior camera control. The SP<sup>®</sup> surgical system is equipped with a fully hinged elbow joint that enables complex operations in confined spaces with greater precision and adaptability. In 2018, two surgeons with experience in robotic mastectomy conducted a preclinical cadaver study using the SP surgical system in Berlin, Germany, and then applied the system to patients<sup>[46]</sup>. Park *et al.* performed the first clinical case using the SP<sup>®</sup> surgical system for RNSM with minimal incision<sup>[46]</sup>. Go *et al.* assessed the safety of the SP<sup>®</sup> surgical system for RNSM, reporting a nipple necrosis rate of 2.5% (one case), with no cases requiring conversion to open mastectomy<sup>[47]</sup>. A prospective study on the da Vinci SP<sup>®</sup> surgical system for robotic nipple-sparing mastectomy (SPrNSM) found that the overall postoperative complication rate was 7.5%, which included one case of postoperative expander infection and one case of skin necrosis<sup>[48]</sup>. The study also demonstrated that SPrNSM had higher rates of intact NAC and skin sensation than open NSM. Currently, there is a lack of data on the oncological safety of the SP<sup>®</sup> surgical system for RNSM, as well as data comparing it with multi-arm RNSM. Moreover, whether it is single-arm or multi-arm RNSM, there is a lack of long-term oncological safety data with big data, which requires multicenter collaboration for research. Of course, technological advances in the Da Vinci robotic surgical system have made our breast surgery incisions smaller, and the operation more flexible, which may be associated with better quality of life and improved esthetic, potentially offering benefits to patients.

### Robotic latissimus dorsi breast reconstruction

The latissimus dorsi flap has been used for breast reconstruction since the 1970s<sup>[49]</sup>. Traditional latissimus dorsi muscle flap harvest techniques require an incision on the back that ranges from 15 to 45 cm, in addition to an axillary incision to transfer the pedicle<sup>[50]</sup>. The pursuit of minimally invasive techniques for harvesting the latissimus dorsi muscle flap has always been a goal. Endoscopic harvesting has been reported by numerous studies<sup>[51,52]</sup>. However, due to technical challenges such as line of sight around the curvature of the back and other limitations of endoscopic instrumentation, some centers have abandoned this technique<sup>[53]</sup>. Selber first described the robotic harvest of the latissimus dorsi muscle in the cadaver model<sup>[54]</sup>. Subsequently, robotic latissimus dorsi harvest was performed on seven patients, and it was found to be a novel and effective method for muscle procurement, with technical advantages over endoscopic harvesting and esthetic advantages over open techniques<sup>[55]</sup>. None of the seven patients experienced donor

**Table 1. Comparative study of robotic breast surgery**

Author (year)	Research type	Group (number)	Operation time	Complication (P value)	Reference
Filipe <i>et al.</i> (2022)	Meta analysis	R-NSM vs. C-NSM	-	0.07	[41]
Lai <i>et al.</i> (2020)	Retrospective study	R-NSM (51) vs. C-NSM (62)	0.01	0.98	[42]
Lai <i>et al.</i> (2020)	Retrospective study	R-NSM (40) vs. E-NSM (91)	0.01	0.284	[44]
Lai <i>et al.</i> (2024)	Prospectively study	R-NSM (76) vs. C-NSM (46) vs. E-NSM (84)	< 0.01 (unilateral mastectomy)	-	[43]
Winocour <i>et al.</i> (2020)	Retrospective study	R-LD (25) vs. C-LD (27)	0.002	0.034	[56]
Eo <i>et al.</i> (2023)	Prospective study	R-LD (20) vs. E-LD (17) vs. C-LD (20)	< 0.001	> 0.90	[57]
Lee <i>et al.</i> (2022)	Retrospective study	R-DIEP (21) vs. C-DIEP (186)	< 0.001	≥ 0.085	[69]
Tsai <i>et al.</i> (2023)	Retrospective study	R-DIEP (13) vs. C-DIEP (86)	0.006	0.348	[70]
Moreira <i>et al.</i> (2024)	Retrospective study	R-DIEP (46) vs. C-DIEP (48)	0.013	0.624	[71]
Elameen <i>et al.</i> (2024)	Meta analysis	R-ABR (263) vs. C-ABR (520)	< 0.001	≥ 0.150	[73]

R-NSM: Robot-assisted nipple-sparing mastectomy; C-NSM: conventional nipple-sparing mastectomy; E-NSM: endoscopic nipple-sparing mastectomy; R-LD: robot-assisted latissimus dorsi muscle harvest; E-LD: endoscopic latissimus dorsi muscle harvest; C-LD: conventional latissimus dorsi muscle harvest; R-DIEP: robot-assisted deep inferior epigastric perforator flap harvest; C-DIEP: conventional deep inferior epigastric perforator flap harvest; R-ABR: robotic autologous breast reconstruction; C-ABR: conventional autologous breast reconstruction.

site hematomas, seroma, or skin injuries, and the surgery time was reduced from over 2 h to approximately 1 h. Despite longer operative times and higher seroma rates, robotic-assisted latissimus dorsi muscle harvest remains an effective alternative to traditional open surgery, offering the benefits of no back scars and lower opioid requirements<sup>[56]</sup>. Patient satisfaction with robotic-assisted latissimus dorsi flap breast reconstruction after partial mastectomy is higher than with traditional open methods, particularly regarding donor site scarring<sup>[57]</sup>. Chen *et al.* conducted a study on robotic NSM followed by immediate breast reconstruction using the latissimus dorsi muscle without island flap<sup>[58]</sup>. They found that compared to reconstruction with latissimus dorsi muscle with skin island, the technique without island flap saved surgical time and reduced intraoperative blood loss, but also included the disadvantage of lacking a skin paddle for monitoring. Robotic breast reconstruction using the latissimus dorsi muscle without island flap shows potential for application in the future. A prospective, single-arm study on robotic-assisted latissimus dorsi muscle harvest has been approved by the U.S. FDA<sup>[59]</sup>. The primary safety endpoint is adverse events caused by the harvest, and the primary efficacy endpoints include muscle viability after the harvest and the conversion rate to open surgery. This study will further explore the feasibility and safety of robotic-assisted latissimus dorsi harvest.

### Robotic deep inferior epigastric perforator flap

Some patients are satisfied with autologous breast reconstruction<sup>[60]</sup>. The initial autologous breast reconstruction was based on abdominal tissue, relying on the pedicled transverse rectus abdominis myocutaneous (TRAM) flap, followed by the free TRAM flap. Currently, the deep inferior epigastric perforator (DIEP) flap has become the standard for autologous breast reconstruction<sup>[61-63]</sup>. Both pedicled and free TRAM flaps sacrifice the rectus abdominis muscle, which can lead to serious abdominal wall complications. In contrast, DIEP flaps, which preserve the rectus abdominis muscle, can significantly reduce the occurrence of abdominal wall complications<sup>[64]</sup>. Even with technical advancements, harvesting the DIEP flap through the traditional open approach still requires a long incision. Since the DIEP pedicle extends along the deep layer of the rectus abdominis muscle, dissecting it often necessitates splitting the muscle,

leading to direct muscle damage. This may damage the motor nerves of the rectus abdominis, resulting in muscle dysfunction and increased postoperative pain<sup>[1]</sup>. Robot-assisted DIEP can significantly reduce the fascial incision and perform the resection through a posterior approach, minimizing the occurrence of muscle and nerve damage<sup>[65]</sup>. Robotic-assisted DIEP flap harvest has been shown to be feasible in cadaveric models, and a totally extraperitoneal approach may further reduce donor site morbidity<sup>[66,67]</sup>. Preoperative computed tomography angiography (CTA) imaging is crucial for determining whether patients are suitable for a robotic approach<sup>[68]</sup>. Patients with a single perforator or two closely grouped perforators are better suited for a robotic DIEP. A retrospective study found that compared to the open DIEP, a robotic DIEP can improve postoperative recovery while reducing postoperative pain and shortening hospital stays<sup>[69]</sup>. Robotic bilateral DIEP flaps are also feasible without the need for port adjustments, thanks to novel port placement techniques<sup>[70]</sup>. Compared to open bilateral DIEP, robotic bilateral DIEP flap harvest can reduce the length of fascial incisions, decrease abdominal wall disruption, and avoid the need for fascial reinforcement with mesh<sup>[71]</sup>. Indocyanine green-guided near-infrared fluorescence aids in vascular dissection during robotic-assisted DIEP<sup>[72]</sup>. The successful harvest of robotic DIEP flaps also confirms the feasibility and safety of robotic autologous breast reconstruction<sup>[73]</sup>. The da Vinci SP® surgical system requires only a 2.5 cm diameter port, which allows surgeons to operate in confined space, such as DIEP surgery. Jung *et al.* successfully used the SP® surgical system for the first time to perform dissection of vascular pedicle through minimal fascial incisions<sup>[74]</sup>.

### Robotic microsurgery

The complete elimination of tremor and 10x magnification advantages of the robotic system make it an appealing choice for super-microsurgery. Robotic microsurgery can be used to treat upper limb lymphedema. In 2010, Selber *et al.* reported cases of oral robotic reconstruction of oropharyngeal defects, including the use of robotic microvascular anastomosis<sup>[75]</sup>. O'Brien first described the technique of lymphovenous bypass in dogs in 1977 as a means of improving lymphedema<sup>[76]</sup>. In the field of supermicrosurgery, the renowned surgeon Koshima made it possible to treat lymphedema with lymphovenous bypass surgery by performing anastomoses on vessels and lymphatics with diameters ranging from 0.3 to 0.6 mm<sup>[77]</sup>. In the first human study of robotic-assisted supermicrosurgery lymphovenous anastomosis (LVA), van Mulken *et al.* compared robotic-assisted anastomosis with manual anastomosis and demonstrated the feasibility of robotic LVA<sup>[78]</sup>. Robotic omentum lymph node flap harvest for the treatment of extremity lymphedema has also been shown to be safe and feasible<sup>[79]</sup>. Lindenblatt *et al.* reported the first use of the Symani surgical system for LVA and arterial anastomosis in lymphatic reconstruction in humans<sup>[26]</sup>. However, the learning curve for robotic-assisted LVA is very steep, as the procedure takes significantly longer than manual suturing techniques at the beginning<sup>[80]</sup>, but once surgeons become familiar with the technology, the time required for robotic-assisted LVA becomes comparable to that of manual suturing techniques. van Mulken *et al.* reported the application of the MUSA in humans for treating breast cancer-related lymphedema<sup>[27]</sup>. The advantages of the MUSA surgical system are also stability, scaling movements, and filtering vibration. The robotic arms of this system are equipped with true microsurgical instruments. Robotic-assisted techniques allow novice and intermediate surgeons to perform at a level comparable to experts, enabling novice learners to train early on in microsurgical anastomosis with the assistance of robotics<sup>[81]</sup>. The studies conducted thus far have demonstrated that the application of robotic-assisted anastomosis techniques in clinical settings is both safe and feasible. Overall, despite the limitation of a steep learning curve, prolonged operative time, and high costs, these advanced robotic systems are critical to the development of microsurgery.

## NEW TRENDS IN ROBOTIC BREAST SURGERY IN THE ERA OF AI

### AI in robotic surgery

AI was coined by John McCarthy in 1955 at Stanford University<sup>[82]</sup>. AI is the use of computer science and related technologies to enable machines to simulate human intelligent behavior<sup>[83]</sup>. From manufacturing to education and other industries, AI is gradually changing the way humans live and work. AI, in particular the deep-learning subtype, has been applied in various healthcare fields, and in some cases, it has even outperformed experienced doctors<sup>[84]</sup>. However, in surgical procedures, AI has not yet fully realized its potential due to some challenges such as safety and efficacy in the discipline<sup>[85]</sup>.

In the field of surgery, AI has a broad range of applications, such as simulation, intraoperative decision making, event and outcome prediction, postoperative progression, complication management, and re-credentialing of surgeons. In this context, the term “surgical data science” has recently been coined<sup>[85]</sup>. AI-based systems can predict future acute kidney injury based on data from hundreds of thousands of patients<sup>[86]</sup>. Similarly, AI can also be used to predict circulatory failure in intensive care units<sup>[87]</sup>.

With regard to surgery, digital surgical methods, such as master-slave manipulators used in robotic-assisted surgery, stand to benefit significantly from the advancements in AI. The most distant goal of the application of AI in robotic surgery is to achieve independent and autonomous completion of advanced surgery. To achieve this goal, robotic systems need to start with conventional anatomical structure recognition, and gradually intelligent learning to complete the long-term goal<sup>[88-91]</sup>. The automation level of robotic surgery ranges from the non-automated “master-slave” method to the fully automated level where the robot makes all surgical decisions without human involvement, divided into six levels<sup>[92,93]</sup>.

Wang *et al.* reported that a convolutional neural network (CNN) can assist in creating a smoke-free intraoperative view during surgery, thus enabling a clearer intraoperative field of view<sup>[94]</sup>. Eslamian *et al.* created a model that combines intraoperative visual field tracking, intraoperative image, and robotic kinematic data, enabling autonomous view transformation and determination of the best surgical visualization<sup>[95]</sup>. These combinations of AI with robotics can enhance the surgical field of view for surgeons during procedures. Kumazu *et al.* created a model by analyzing surgical videos of robot-assisted surgeries<sup>[96]</sup>. This model is capable of automatically segmenting loose connective tissue, demonstrating strong performance in identifying the safety plane. Marsden *et al.* proposed an AI model based on integrating fiber fluorescence imaging with robotic surgical platform cameras, and the model has the potential to assess intraoperative edges<sup>[97]</sup>. The sensitivity and specificity for identifying different anatomical structures of the oral cavity and oropharynx were 86% and 87%, respectively. Bianchi *et al.* used preoperative multi-parametric magnetic resonance imaging (MRI) images to generate a 3D model in order to accurately cut the lesion and identify the tumor margin, which could be combined with the robot field of view during the operation and instruct the intraoperative tumor margin cutting<sup>[98]</sup>. In this study, there were no significant differences between the model group and the control group in terms of operation time and postoperative complications. However, the rate of positive surgical margins in the model group (5%) was significantly lower than that in the control group (20%). These methods may also guide surgeons in identifying tumor boundaries during surgery to facilitate complete cancer resection in the future. In the age of AI, automated system performance indicators can be used for objective, real-time surgical performance assessment tools through dynamic and video analysis. The learning curve of robotic-assisted surgery is very long, and its training is mainly carried out through training courses, virtual reality, etc. Training in robotic surgical systems is extremely valuable, and machine learning and video feedback can enhance education in robotic-assisted surgery<sup>[99]</sup>. By predicting surgical proficiency based on the training intervals of key surgical tasks and video segments, it is possible to differentiate the performance of expert and novice surgeons<sup>[100]</sup>.

Together, these studies lay the foundation for the future automation of robotic surgery.

### AI in robotic breast surgery

Machine learning is a facet of AI and deep learning is a facet of machine learning. Deep learning has the ability to automatically extract features and process large amounts of data. It works by establishing an artificial neural network system capable of classifying and recognizing images. The role of deep learning in assisting image analysis for the early detection of cancer is gradually growing<sup>[101]</sup>. Liao *et al.* developed an ensemble deep learning model, EDL-BC, based on deep CNN. This model can extract morphological features of early breast lesions, perform effective and objective image analysis, improve the accuracy of differential diagnosis of breast nodules, and provide an accurate prognosis for early breast cancer<sup>[102]</sup>. Yoon *et al.* conducted a meta-analysis on the use of AI for breast cancer detection in screening digital mammography and digital breast tomosynthesis, finding that the performance of standalone AI in digital mammography is as good as that of radiologists<sup>[103]</sup>. Radiomics can analyze medical images and transform them into quantitative data. By using machine learning to extract quantitative imaging features at high throughput, it can further assess the tumor microenvironment and heterogeneity<sup>[104]</sup>. Zhao *et al.* developed a machine-learning-based radiomics model that can accurately predict the response to immune checkpoint inhibitors (ICIs)-based therapy in patients with advanced breast cancer<sup>[105]</sup>.

AI is more accurate and faster than humans in diagnosis and treatment decision making. IBM Watson for Oncology (WFO) is one of the most advanced AI decision-making systems, with a consistency rate of 56% between its intelligent decisions and those of doctors. It has shown excellent feasibility and standardization in the application of breast cancer treatment<sup>[106]</sup>. A retrospective multicenter study integrated MRI radiomics features, axillary lymph node (ALN), clinicopathological characteristics, and molecular subtypes through machine learning to propose a multi-omic signature. This signature can conveniently identify patients with ALN metastasis among different molecular subtypes in early-stage breast cancer, potentially guiding future surgery<sup>[107]</sup>. Existing postoperative pain assessments require extensive questionnaires and interaction with patients, while machine learning in breast surgery can predict postoperative pain immediately after surgery<sup>[108,109]</sup>. Machine learning models also indicate that artificial neural networks achieve superior predictive performance and are significantly better than other predictive models, which could predict breast cancer recurrence up to 10 years after surgery<sup>[110]</sup>. In a preliminary study, Mavioso *et al.* utilized machine learning to assess the vascular CT imaging of patients planned for DIEP breast reconstruction, discovering that this technology could reduce the preoperative planning time for DIEP flaps<sup>[111]</sup>. O'Neill *et al.* developed a machine learning algorithm to establish a clinical predictive model capable of identifying the risk of flap failure in patients undergoing DIEP flap reconstruction. The results indicated that age  $\geq 40$  years, BMI  $\geq 40$ , unilateral or bilateral reconstruction, and the presence of complications may contribute to flap failure<sup>[112]</sup>.

The application of AI in breast surgery is still largely conceptual, but computer vision technology can support surgical techniques, and be used in anatomical visualization and surgical navigation<sup>[113]</sup>. AI, involving machine learning and supported by deep learning, offers opportunities for real-time decision making in breast surgery and can also enhance the realism of robotic breast surgery. Anatomical visualization and surgical navigation are crucial for visualization and completely removing tumor tissue during robotic breast surgery, as well as predicting and resecting axillary positive lymph nodes during surgery. Matrix metalloproteinase-9 (MMP-9) and fibronectin are biomarkers for breast cancer. Cheng *et al.* have developed a new imaging probe, CREKA-GK8-QC, targeted at fibronectin and MMP-9 activation, for detecting *in situ* and metastatic breast cancer<sup>[114]</sup>. This probe can facilitate complete tumor resection and reduce iatrogenic damage during surgery under the guidance of fluorescence imaging. The probe may also be applicable to robotic breast surgery and even robotic breast-conserving surgery. Robotic ultrasound-assisted biopsy technology enables the integration of robotics and AI with MRI and ultrasound (US)

imaging, offering a high degree of precision in needle placement for breast cancer imaging and biopsy<sup>[115]</sup>.

Duarte *et al.* proposed an innovative deep learning technology that integrates CNN, long short-term memory networks (LSTM), and multilayer perceptrons (MLP) to generate depth maps from RGB images, which are then converted in real time into 3D meshes for the three-dimensional reconstruction of the breast<sup>[116]</sup>. This method can be used for robotic-guided breast biopsy. Additionally, this technology could be further utilized to predict breast size and volume for the calculation of the amount of prosthesis or tissue needed to extend to the field of breast reconstruction, and even for 3D printed breast reconstruction. Peng *et al.* conducted a study on a lesion morphology aware network, LMA-Net, segmenting breast tumors in MRI images, which includes two stages: breast segmentation and tumor segmentation<sup>[117]</sup>. Alqaoud *et al.* utilized deep learning based on nnU-Net to segment MRI images of the fatty, fibroglandular, and tumor tissues, thereby laying the foundation for robotic surgery planning and image-guided navigation<sup>[118]</sup>. The study also proposed a simulated breast phantom created from these segmentation results, introducing a polyvinyl alcohol cryogel (PVA-C) breast model for experimental evaluation and validation of the research. The 3D-printed PVA-C mold was produced according to the surfaces extracted from the segmentation outcomes. Although still in the research phase, these developments provide a solid foundation for future clinical applications in robotic breast surgery.

### The future

Although current surgical robots are fully remotely operated devices, the “semi-automation” and “automation” of robotic systems are advancing. Researchers are addressing the technical challenges of surgical process automation through the development of real-world task demonstrations, such as the da Vinci Research Kit (dVRK), KUKA LBR Med collaborative robot, and RAVEN robot<sup>[119]</sup>. Surgical intelligent vision is a challenging task, but if applied to robotic surgery, such as intraoperative decision support and image-guided surgery, it will lead to advancements in future surgical procedures<sup>[120]</sup>. One of the main differences between robotic-assisted surgery and open surgery is the sense of touch. Developing an intelligent robot with a complete tactile feedback loop requires high-resolution sensor skin, algorithms to interpret sensor information, and reliable feedback control that corresponds to the skin, nerves, and the brain. Bao *et al.* have integrated robotics with AI to study an integrated intelligent tactile system based on touch, achieving real-time tactile perception and grasp control of a humanoid robot’s intelligent robotic hand<sup>[121]</sup>. Guo *et al.* have developed a tailored wearable haptic controller for robot-assisted microsurgery systems. This haptic glove is designed to capture hand movements for remote operation of robot-assisted microsurgery, and enhance the user’s depth perception to reduce depth errors in microsurgical tasks<sup>[122]</sup>. The integration of AI with robotic breast surgery may also promote the improvement of medical materials. In the future, appropriate reconstruction materials can be selected through preoperative AI and imaging data modeling to evaluate breast size and density. Robotic remote breast surgery is an important direction for the future, particularly in addressing the surgical needs of patients in areas with limited medical resources or in dangerous regions. In robotic remote surgery, signal latency can cause image interference and oscillation of robotic instruments, thereby affecting surgical safety. Motion scaling may be a solution, and the issue of signal delay can be further resolved by improving motion scaling algorithms and customizing them for specific surgical tasks and conditions in the future<sup>[123]</sup>. Moreover, the automation of robotic breast surgery will inevitably face ethical constraints. Therefore, it is necessary to standardize autonomous robots in terms of ethics, standards, and regulations to establish public confidence in autonomous robotic systems. In the near future, the task of automating robotic surgery, enabling autonomous robotic programs to adapt to real environments and their resilience, will be the true challenge ahead.

## CONCLUSION

The integration of AI with robotic breast surgery represents a significant leap forward in the evolution of surgical procedures. With continued advancements, breast surgery is likely to become more precise, less invasive, and increasingly tailored to individual patients. Ultimately, this progress promises to improve the survival rates and enhance the quality of life for those affected by breast cancer.

## DECLARATIONS

### Authors' contributions

Made substantial contributions to the conception and design of the study: Li C

Manuscript writing: Li X, Li C

### Availability of data and materials

Not applicable.

### Financial support and sponsorship

None.

### Conflicts of interest

All authors declared that there are no conflicts of interest.

### Ethical approval and consent to participate

Not applicable.

### Consent for publication

Not applicable.

### Copyright

© The Author(s) 2025.

## REFERENCES

1. Bishop SN, Selber JC. Minimally invasive robotic breast reconstruction surgery. *Gland Surg.* 2021;10:469-78. [DOI PubMed PMC](#)
2. Zhang S, Xie Y, Liang F, Wang Y, Lv Q, Du Z. Endoscopic-assisted nipple-sparing mastectomy with direct-to-implant subpectoral breast reconstruction in the management of breast cancer. *Plast Reconstr Surg Glob Open.* 2021;9:e3978. [DOI PubMed PMC](#)
3. Leff DR, Vashisht R, Yongue G, Keshtgar M, Yang GZ, Darzi A. Endoscopic breast surgery: where are we now and what might the future hold for video-assisted breast surgery? *Breast Cancer Res Treat.* 2011;125:607-25. [DOI PubMed](#)
4. Pavone M, Baroni A, Campolo F, et al. Robotic assisted versus laparoscopic surgery for deep endometriosis: a meta-analysis of current evidence. *J Robot Surg.* 2024;18:212. [DOI PubMed PMC](#)
5. Chen K, M Beeraka N, Zhang J, et al. Efficacy of Da Vinci robot-assisted lymph node surgery than conventional axillary lymph node dissection in breast cancer - a comparative study. *Int J Med Robot.* 2021;17:e2307. [DOI PubMed](#)
6. Nessa A, Shaikh S, Fuller M, Masannat YA, Kastora SL. Postoperative complications and surgical outcomes of robotic versus conventional nipple-sparing mastectomy in breast cancer: meta-analysis. *Br J Surg.* 2024;111:znad336. [DOI PubMed PMC](#)
7. Bashir M, Harky A. Artificial intelligence in aortic surgery: the rise of the machine. *Semin Thorac Cardiovasc Surg.* 2019;31:635-7. [DOI PubMed](#)
8. Raghupathi W, Raghupathi V. Big data analytics in healthcare: promise and potential. *Health Inf Sci Syst.* 2014;2:3. [DOI PubMed PMC](#)
9. Zarkowsky DS, Stonko DP. Artificial intelligence's role in vascular surgery decision-making. *Semin Vasc Surg.* 2021;34:260-7. [DOI PubMed](#)
10. Eadie LH, Seifalian AM, Davidson BR. Telemedicine in surgery. *Br J Surg.* 2003;90:647-58. [DOI PubMed](#)
11. Loftus TJ, Tighe PJ, Filiberto AC, et al. Artificial intelligence and surgical decision-making. *JAMA Surg.* 2020;155:148-58. [DOI PubMed PMC](#)
12. Satava RM. Surgical robotics: the early chronicles: a personal historical perspective. *Surg Laparosc Endosc Percutan Tech.* 2002;12:6-16. [DOI PubMed](#)

13. Morrell ALG, Morrell-Junior AC, Morrell AG, et al. The history of robotic surgery and its evolution: when illusion becomes reality. *Rev Col Bras Cir.* 2021;48:e20202798. DOI PubMed PMC
14. Ghezzi T, Campos Corleta O. 30 years of robotic surgery. *World J Surg.* 2016;40:2550-7. DOI PubMed
15. Ranev D, Teixeira J. History of computer-assisted surgery. *Surg Clin North Am.* 2020;100:209-18. DOI PubMed
16. van Mulken TJM, Boymans CAEM, Schols RM, et al. Preclinical experience using a new robotic system created for microsurgery. *Plast Reconstr Surg.* 2018;142:1367-76. DOI PubMed
17. Guerrini GP, Esposito G, Magistri P, et al. Robotic versus laparoscopic gastrectomy for gastric cancer: the largest meta-analysis. *Int J Surg.* 2020;82:210-28. DOI PubMed
18. Guo Z, Shi Y, Song Z, et al. Single-incision robotic assisted surgery: a nonrandomized cohort pilot study on a novel surgical platform in colorectal surgery. *Int J Surg.* 2023;109:3417-29. DOI PubMed PMC
19. Schmelzle M, Feldbrügge L, Ortiz Galindo SA, et al. Robotic vs. laparoscopic liver surgery: a single-center analysis of 600 consecutive patients in 6 years. *Surg Endosc.* 2022;36:5854-62. DOI PubMed PMC
20. Kalata S, Thumma JR, Norton EC, Dimick JB, Sheetz KH. Comparative safety of robotic-assisted vs laparoscopic cholecystectomy. *JAMA Surg.* 2023;158:1303-10. DOI PubMed PMC
21. Catto JWF, Khetrapal P, Ricciardi F, et al; iROC Study Team. Effect of robot-assisted radical cystectomy with intracorporeal urinary diversion vs open radical cystectomy on 90-day morbidity and mortality among patients with bladder cancer: a randomized clinical trial. *JAMA.* 2022;327:2092-103. DOI PubMed PMC
22. Minamimura K, Aoki Y, Kaneya Y, et al. Current status of robotic hepatobiliary and pancreatic surgery. *J Nippon Med Sch.* 2024;91:10-9. DOI PubMed
23. Alkatout I, O'Sullivan O, Peters G, Maass N. Expanding robotic-assisted surgery in gynecology using the potential of an advanced robotic system. *Medicina.* 2023;60:53. DOI PubMed PMC
24. Bae HL, Wong JS, Kim SJ, et al. Surgical outcomes of robotic thyroidectomy for thyroid tumors over 4 cm via the bilateral axillo-breast approach. *Sci Rep.* 2024;14:11646. DOI PubMed PMC
25. Toesca A, Peradze N, Galimberti V, et al. Robotic nipple-sparing mastectomy and immediate breast reconstruction with implant: first report of surgical technique. *Ann Surg.* 2017;266:e28-30. DOI PubMed
26. Lindenblatt N, Grünherz L, Wang A, et al. Early experience using a new robotic microsurgical system for lymphatic surgery. *Plast Reconstr Surg Glob Open.* 2022;10:e4013. DOI PubMed PMC
27. van Mulken TJM, Schols RM, Scharmga AMJ, et al; MicroSurgical Robot Research Group. First-in-human robotic supermicrosurgery using a dedicated microsurgical robot for treating breast cancer-related lymphedema: a randomized pilot trial. *Nat Commun.* 2020;11:757. DOI PubMed PMC
28. Pandav K, Te AG, Tomer N, Nair SS, Tewari AK. Leveraging 5G technology for robotic surgery and cancer care. *Cancer Rep.* 2022;5:e1595. DOI PubMed PMC
29. O'Sullivan S, Leonard S, Holzinger A, et al. Operational framework and training standard requirements for AI-empowered robotic surgery. *Int J Med Robot.* 2020;16:1-13. DOI PubMed
30. Medina-Franco H, Vasconez LO, Fix RJ, et al. Factors associated with local recurrence after skin-sparing mastectomy and immediate breast reconstruction for invasive breast cancer. *Ann Surg.* 2002;235:814-9. DOI PubMed PMC
31. Ueda S, Tamaki Y, Yano K, et al. Cosmetic outcome and patient satisfaction after skin-sparing mastectomy for breast cancer with immediate reconstruction of the breast. *Surgery.* 2008;143:414-25. DOI PubMed
32. Yi M, Kronowitz SJ, Meric-Bernstam F, et al. Local, regional, and systemic recurrence rates in patients undergoing skin-sparing mastectomy compared with conventional mastectomy. *Cancer.* 2011;117:916-24. DOI PubMed PMC
33. Yamashita Y, Tsunoda H, Nagura N, et al. Long-term oncologic safety of nipple-sparing mastectomy with immediate reconstruction. *Clin Breast Cancer.* 2021;21:352-9. DOI PubMed
34. Sacchini V, Pinotti JA, Barros AC, et al. Nipple-sparing mastectomy for breast cancer and risk reduction: oncologic or technical problem? *J Am Coll Surg.* 2006;203:704-14. DOI PubMed
35. Wagner JL, Fearmonti R, Hunt KK, et al. Prospective evaluation of the nipple-areola complex sparing mastectomy for risk reduction and for early-stage breast cancer. *Ann Surg Oncol.* 2012;19:1137-44. DOI PubMed PMC
36. Gerber B, Krause A, Dieterich M, Kundt G, Reimer T. The oncological safety of skin sparing mastectomy with conservation of the nipple-areola complex and autologous reconstruction: an extended follow-up study. *Ann Surg.* 2009;249:461-8. DOI PubMed
37. Haslinger ML, Sosin M, Bartholomew AJ, et al. Positive nipple margin after nipple-sparing mastectomy: an alternative and oncologically safe approach to preserving the nipple-areolar complex. *Ann Surg Oncol.* 2018;25:2303-7. DOI PubMed
38. Mesdag V, Régis C, Tresch E, et al. Nipple sparing mastectomy for breast cancer is associated with high patient satisfaction and safe oncological outcomes. *J Gynecol Obstet Hum Reprod.* 2017;46:637-42. DOI PubMed
39. Lai HW, Chen ST, Chen DR, et al. Current trends in and indications for endoscopy-assisted breast surgery for breast cancer: results from a six-year study conducted by the taiwan endoscopic breast surgery cooperative group. *PLoS One.* 2016;11:e0150310. DOI PubMed PMC
40. Toesca A, Peradze N, Manconi A, et al. Robotic nipple-sparing mastectomy for the treatment of breast cancer: feasibility and safety study. *Breast.* 2017;31:51-6. DOI PubMed PMC
41. Filipe MD, de Bock E, Postma EL, et al. Robotic nipple-sparing mastectomy complication rate compared to traditional nipple-sparing mastectomy: a systematic review and meta-analysis. *J Robot Surg.* 2022;16:265-72. DOI PubMed PMC

42. Lai HW, Chen ST, Mok CW, et al. Robotic versus conventional nipple sparing mastectomy and immediate gel implant breast reconstruction in the management of breast cancer- a case control comparison study with analysis of clinical outcome, medical cost, and patient-reported cosmetic results. *J Plast Reconstr Aesthet Surg.* 2020;73:1514-25. DOI PubMed
43. Lai HW, Chen DR, Liu LC, et al. Robotic versus conventional or endoscopic-assisted nipple-sparing mastectomy and immediate prosthesis breast reconstruction in the management of breast cancer: a prospectively designed multicenter trial comparing clinical outcomes, medical cost, and patient-reported outcomes (RCENSM-P). *Ann Surg.* 2024;279:138-46. DOI PubMed PMC
44. Lai HW, Chen ST, Tai CM, et al. Robotic- versus endoscopic-assisted nipple-sparing mastectomy with immediate prosthesis breast reconstruction in the management of breast cancer: a case-control comparison study with analysis of clinical outcomes, learning curve, patient-reported aesthetic results, and medical cost. *Ann Surg Oncol.* 2020;27:2255-68. DOI PubMed
45. Garden EB, Al-Alao O, Razdan S, Mullen GR, Florman S, Palese MA. Robotic single-port donor nephrectomy with the da vinci sp® surgical system. *JSL.S.* 2021;25:e2021.00062. DOI PubMed PMC
46. Park HS, Lee J, Lee H, Lee K, Song SY, Toesca A. Development of robotic mastectomy using a single-port surgical robot system. *J Breast Cancer.* 2020;23:107-12. DOI PubMed PMC
47. Go J, Ahn JH, Park JM, et al. Analysis of robot-assisted nipple-sparing mastectomy using the da Vinci SP system. *J Surg Oncol.* 2022;126:417-24. DOI PubMed
48. Farr DE, Haddock NT, Tellez J, et al. Safety and feasibility of single-port robotic-assisted nipple-sparing mastectomy. *JAMA Surg.* 2024;159:269-76. DOI PubMed PMC
49. Bostwick J 3rd, Vasconez LO, Jurkiewicz MJ. Breast reconstruction after a radical mastectomy. *Plast Reconstr Surg.* 1978;61:682-93. DOI PubMed
50. Lin CH, Wei FC, Levin LS, Chen MC. Donor-site morbidity comparison between endoscopically assisted and traditional harvest of free latissimus dorsi muscle flap. *Plast Reconstr Surg.* 1999;104:1070-7. PubMed
51. Miller MJ, Robb GL. Endoscopic technique for free flap harvesting. *Clin Plast Surg.* 1995;22:755-73. PubMed
52. Pomel C, Missana MC, Lasser P. [Endoscopic harvesting of the latissimus dorsi flap in breast reconstructive surgery. Feasibility study and review of the literature]. *Ann Chir.* 2002;127:337-42. DOI PubMed
53. Fine NA, Orgill DP, Pribaz JJ. Early clinical experience in endoscopic-assisted muscle flap harvest. *Ann Plast Surg.* 1994;33:465-9. DOI PubMed
54. Selber JC, Baumann DP, Holsinger CF. Robotic harvest of the latissimus dorsi muscle: laboratory and clinical experience. *J Reconstr Microsurg.* 2012;28:457-64. DOI PubMed
55. Selber JC, Baumann DP, Holsinger FC. Robotic latissimus dorsi muscle harvest: a case series. *Plast Reconstr Surg.* 2012;129:1305-12. DOI PubMed
56. Winocour S, Tarassoli S, Chu CK, Liu J, Clemens MW, Selber JC. Comparing outcomes of robotically assisted latissimus dorsi harvest to the traditional open approach in breast reconstruction. *Plast Reconstr Surg.* 2020;146:1221-5. DOI PubMed
57. Eo PS, Kim H, Lee JS, Lee J, Park HY, Yang JD. Robot-assisted latissimus dorsi flap harvest for partial breast reconstruction: comparison with endoscopic and conventional approaches. *Aesthet Surg J.* 2023;44:38-46. DOI PubMed
58. Chen K, Zhang J, Beeraka NM, Lu P. Robotic nipple sparing mastectomy and immediate breast reconstruction: significant attempts with the latissimus dorsi muscle without island flap. *Minerva Surg.* 2024;79:411-8. DOI PubMed
59. Shuck J, Asaad M, Liu J, Clemens MW, Selber JC. Prospective pilot study of robotic-assisted harvest of the latissimus dorsi muscle: a 510(k) approval study with U.S. Food and Drug Administration investigational device exemption. *Plast Reconstr Surg.* 2022;149:1287-95. DOI PubMed
60. von Glinski M, Holler N, Kümmel S, et al. The partner perspective on autologous and implant-based breast reconstruction. *Aesthetic Plast Surg.* 2023;47:1324-31. DOI PubMed PMC
61. Schefflan M, Hartrampf CR, Black PW. Breast reconstruction with a transverse abdominal island flap. *Plast Reconstr Surg.* 1982;69:908-9. PubMed
62. Speck NE, Grufman V, Farhadi J. Trends and innovations in autologous breast reconstruction. *Arch Plast Surg.* 2023;50:240-7. DOI PubMed PMC
63. Tønseth KA, Hokland BM, Tindholdt TT, Abyholm FE, Stavem K. Quality of life, patient satisfaction and cosmetic outcome after breast reconstruction using DIEP flap or expandable breast implant. *J Plast Reconstr Aesthet Surg.* 2008;61:1188-94. DOI PubMed
64. Bruce JC, Batchinsky M, Van Spronsen NR, Sinha I, Bharadia D. Analysis of online materials regarding DIEP and TRAM flap autologous breast reconstruction. *J Plast Reconstr Aesthet Surg.* 2023;82:81-91. DOI PubMed
65. Khan MTA, Won BW, Baumgardner K, et al. Literature review: robotic-assisted harvest of deep inferior epigastric flap for breast reconstruction. *Ann Plast Surg.* 2022;89:703-8. DOI PubMed
66. Struk S, Sarfati B, Leymarie N, et al. Robotic-assisted DIEP flap harvest: a feasibility study on cadaveric model. *J Plast Reconstr Aesthet Surg.* 2018;71:259-61. DOI PubMed
67. Manrique OJ, Bustos SS, Mohan AT, et al. Robotic-assisted DIEP flap harvest for autologous breast reconstruction: a comparative feasibility study on a cadaveric model. *J Reconstr Microsurg.* 2020;36:362-8. DOI PubMed
68. Kurlander DE, Le-Petross HT, Shuck JW, Butler CE, Selber JC. Robotic DIEP patient selection: analysis of CT angiography. *Plast Reconstr Surg Glob Open.* 2021;9:e3970. DOI PubMed PMC
69. Lee MJ, Won J, Song SY, et al. Clinical outcomes following robotic versus conventional DIEP flap in breast reconstruction: a retrospective matched study. *Front Oncol.* 2022;12:989231. DOI PubMed PMC

70. Tsai CY, Kim BS, Kuo WL, et al. Novel port placement in robot-assisted DIEP flap harvest improves visibility and bilateral DIEP access: early controlled cohort study. *Plast Reconstr Surg.* 2023;152:590e-5e. DOI PubMed
71. Moreira A, Bailey EA, Chen B, et al. A new era in perforator flap surgery for breast reconstruction: a comparative study of robotic versus standard harvest of bilateral deep inferior epigastric artery perforator flaps. *J Reconstr Microsurg.* 2024;Epub ahead of print. DOI PubMed
72. Nelson W, Murariu D, Moreira AA. Indocyanine green-guided near-infrared fluorescence enhances vascular anatomy in robot-assisted DIEP flap harvest. *Plast Reconstr Surg.* 2024;153:796-8. DOI PubMed
73. Elameen AM, Dahy AA. Surgical outcomes of robotic versus conventional autologous breast reconstruction: a systematic review and meta-analysis. *J Robot Surg.* 2024;18:189. DOI PubMed PMC
74. Jung JH, Jeon YR, Lee DW, et al. Initial report of extraperitoneal pedicle dissection in deep inferior epigastric perforator flap breast reconstruction using the da Vinci SP. *Arch Plast Surg.* 2022;49:34-8. DOI PubMed PMC
75. Selber JC. Transoral robotic reconstruction of oropharyngeal defects: a case series. *Plast Reconstr Surg.* 2010;126:1978-87. DOI PubMed
76. O'Brien BM, Sykes P, Threlfall GN, Browning FS. Microlymphaticovenous anastomoses for obstructive lymphedema. *Plast Reconstr Surg.* 1977;60:197-211. DOI PubMed
77. Koshima I, Inagawa K, Urushibara K, Moriguchi T. Supermicrosurgical lymphaticovenular anastomosis for the treatment of lymphedema in the upper extremities. *J Reconstr Microsurg.* 2000;16:437-42. DOI PubMed
78. van Mulken TJM, Wolfs JAGN, Qiu SS, et al; MicroSurgical Robot Research Group. One-year outcomes of the first human trial on robot-assisted lymphaticovenous anastomosis for breast cancer-related lymphedema. *Plast Reconstr Surg.* 2022;149:151-61. DOI PubMed
79. Teven CM, Yi J, Hammond JB, et al. Expanding the horizon: single-port robotic vascularized omentum lymphatic transplant. *Plast Reconstr Surg Glob Open.* 2021;9:e3414. DOI PubMed PMC
80. Weinzierl A, Barbon C, Gousopoulos E, et al. Benefits of robotic-assisted lymphatic microsurgery in deep anatomical planes. *JPRAS Open.* 2023;37:145-54. DOI PubMed PMC
81. Frieberg H, Winter JM, Engström O, Önefält D, Nilsson A, Mani M. Robot-assisted microsurgery-what does the learning curve look like? *JPRAS Open.* 2024;42:33-41. DOI PubMed PMC
82. Moglia A, Georgiou K, Georgiou E, Satava RM, Cuschieri A. A systematic review on artificial intelligence in robot-assisted surgery. *Int J Surg.* 2021;95:106151. DOI PubMed
83. Sung JY. Introduction to artificial intelligence in medicine. *Singapore Med J.* 2024;65:132. DOI PubMed PMC
84. Topol EJ. High-performance medicine: the convergence of human and artificial intelligence. *Nat Med.* 2019;25:44-56. DOI PubMed
85. Maier-Hein L, Vedula SS, Speidel S, et al. Surgical data science for next-generation interventions. *Nat Biomed Eng.* 2017;1:691-6. DOI PubMed
86. Bajaj T, Koyner JL. Artificial intelligence in acute kidney injury prediction. *Adv Chronic Kidney Dis.* 2022;29:450-60. DOI PubMed PMC
87. Li S, Hickey GW, Lander MM, Kanwar MK. Artificial intelligence and mechanical circulatory support. *Heart Fail Clin.* 2022;18:301-9. DOI PubMed
88. Zhou XY, Guo Y, Shen M, Yang GZ. Application of artificial intelligence in surgery. *Front Med.* 2020;14:417-30. DOI PubMed
89. Ma R, Vanstrum EB, Lee R, Chen J, Hung AJ. Machine learning in the optimization of robotics in the operative field. *Curr Opin Urol.* 2020;30:808-16. DOI PubMed PMC
90. Kassahun Y, Yu B, Tibebu AT, et al. Surgical robotics beyond enhanced dexterity instrumentation: a survey of machine learning techniques and their role in intelligent and autonomous surgical actions. *Int J Comput Assist Radiol Surg.* 2016;11:553-68. DOI PubMed
91. Moustris GP, Hiridis SC, Deliparaschos KM, Konstantinidis KM. Evolution of autonomous and semi-autonomous robotic surgical systems: a review of the literature. *Int J Med Robot.* 2011;7:375-92. DOI PubMed
92. Panesar S, Cagle Y, Chander D, Morey J, Fernandez-Miranda J, Kliot M. Artificial intelligence and the future of surgical robotics. *Ann Surg.* 2019;270:223-6. DOI PubMed
93. Knudsen JE, Ghaffar U, Ma R, Hung AJ. Clinical applications of artificial intelligence in robotic surgery. *J Robot Surg.* 2024;18:102. DOI PubMed PMC
94. Wang F, Sun X, Li J. Surgical smoke removal via residual Swin transformer network. *Int J Comput Assist Radiol Surg.* 2023;18:1417-27. DOI PubMed
95. Eslamian S, Reisner LA, Pandya AK. Development and evaluation of an autonomous camera control algorithm on the da Vinci surgical system. *Int J Med Robot.* 2020;16:e2036. DOI PubMed
96. Kumazu Y, Kobayashi N, Kitamura N, et al. Automated segmentation by deep learning of loose connective tissue fibers to define safe dissection planes in robot-assisted gastrectomy. *Sci Rep.* 2021;11:21198. DOI PubMed PMC
97. Marsden M, Weyers BW, Bec J, et al. Intraoperative margin assessment in oral and oropharyngeal cancer using label-free fluorescence lifetime imaging and machine learning. *IEEE Trans Biomed Eng.* 2021;68:857-68. DOI PubMed PMC
98. Bianchi L, Chessa F, Angiolini A, et al. The use of augmented reality to guide the intraoperative frozen section during robot-assisted radical prostatectomy. *Eur Urol.* 2021;80:480-8. DOI PubMed
99. Azadi S, Green IC, Arnold A, Truong M, Potts J, Martino MA. Robotic surgery: the impact of simulation and other innovative

- platforms on performance and training. *J Minim Invasive Gynecol.* 2021;28:490-5. DOI PubMed
100. Juarez-Villalobos L, Hevia-Montiel N, Perez-Gonzalez J. Machine learning based classification of local robotic surgical skills in a training tasks set. *Annu Int Conf IEEE Eng Med Biol Soc.* 2021;2021:4596-9. DOI
  101. Tran KA, Kondrashova O, Bradley A, Williams ED, Pearson JV, Waddell N. Deep learning in cancer diagnosis, prognosis and treatment selection. *Genome Med.* 2021;13:152. DOI PubMed PMC
  102. Liao J, Gui Y, Li Z, et al. Artificial intelligence-assisted ultrasound image analysis to discriminate early breast cancer in Chinese population: a retrospective, multicentre, cohort study. *EclinicalMedicine.* 2023;60:102001. DOI PubMed PMC
  103. Yoon JH, Strand F, Baltzer PAT, et al. Standalone AI for breast cancer detection at screening digital mammography and digital breast tomosynthesis: a systematic review and meta-analysis. *Radiology.* 2023;307:e222639. DOI PubMed PMC
  104. Lambin P, Leijenaar RTH, Deist TM, et al. Radiomics: the bridge between medical imaging and personalized medicine. *Nat Rev Clin Oncol.* 2017;14:749-62. DOI PubMed
  105. Zhao J, Sun Z, Yu Y, et al. Radiomic and clinical data integration using machine learning predict the efficacy of anti-PD-1 antibodies-based combinational treatment in advanced breast cancer: a multicentered study. *J Immunother Cancer.* 2023;11:e006514. DOI PubMed PMC
  106. van de Sande D, Sharabiani M, Bluemink H, et al. Artificial intelligence based treatment planning of radiotherapy for locally advanced breast cancer. *Phys Imaging Radiat Oncol.* 2021;20:111-6. DOI PubMed PMC
  107. Yu Y, He Z, Ouyang J, et al. Magnetic resonance imaging radiomics predicts preoperative axillary lymph node metastasis to support surgical decisions and is associated with tumor microenvironment in invasive breast cancer: a machine learning, multicenter study. *EBioMedicine.* 2021;69:103460. DOI PubMed PMC
  108. Juwara L, Arora N, Gornitsky M, Saha-Chaudhuri P, Velly AM. Identifying predictive factors for neuropathic pain after breast cancer surgery using machine learning. *Int J Med Inform.* 2020;141:104170. DOI PubMed
  109. Lötsch J, Sipilä R, Dimova V, Kalso E. Machine-learned selection of psychological questionnaire items relevant to the development of persistent pain after breast cancer surgery. *Br J Anaesth.* 2018;121:1123-32. DOI PubMed
  110. Lou SJ, Hou MF, Chang HT, et al. Machine learning algorithms to predict recurrence within 10 years after breast cancer surgery: a prospective cohort study. *Cancers.* 2020;12:3817. DOI PubMed PMC
  111. Mavioso C, Araújo RJ, Oliveira HP, et al. Automatic detection of perforators for microsurgical reconstruction. *Breast.* 2020;50:19-24. DOI PubMed PMC
  112. O'Neill AC, Yang D, Roy M, Sebastiaipillai S, Hofer SOP, Xu W. Development and evaluation of a machine learning prediction model for flap failure in microvascular breast reconstruction. *Ann Surg Oncol.* 2020;27:3466-75. DOI PubMed
  113. Soh CL, Shah V, Arjomandi Rad A, et al. Present and future of machine learning in breast surgery: systematic review. *Br J Surg.* 2022;109:1053-62. DOI PubMed PMC
  114. Cheng Z, Jin Y, Li J, et al. Fibronectin-targeting and metalloproteinase-activatable smart imaging probe for fluorescence imaging and image-guided surgery of breast cancer. *J Nanobiotechnology.* 2023;21:112. DOI PubMed PMC
  115. Welleweerd MK, Siepel FJ, Groenhuis V, Veltman J, Stramigioli S. Design of an end-effector for robot-assisted ultrasound-guided breast biopsies. *Int J Comput Assist Radiol Surg.* 2020;15:681-90. DOI PubMed PMC
  116. Duarte B, Oliveira B, Torres HR, Morais P, Fonseca JC, Vilaca JL. Robust 3D breast reconstruction based on monocular images and artificial intelligence for robotic guided oncological interventions. *Annu Int Conf IEEE Eng Med Biol Soc.* 2023;2023:1-4. DOI PubMed
  117. Peng C, Zhang Y, Meng Y, et al. LMA-Net: a lesion morphology aware network for medical image segmentation towards breast tumors. *Comput Biol Med.* 2022;147:105685. DOI PubMed
  118. Alqaoud M, Plemmons J, Feliberti E, et al. nnUNet-based multi-modality breast mri segmentation and tissue-delineating phantom for robotic tumor surgery planning. *Annu Int Conf IEEE Eng Med Biol Soc.* 2022;2022:3495-501. DOI PubMed
  119. Fiorini P, Goldberg KY, Liu Y, Taylor RH. Concepts and trends in autonomy for robot-assisted surgery. *Proc IEEE Inst Electr Electron Eng.* 2022;110:993-1011. DOI PubMed PMC
  120. Kitaguchi D, Takeshita N, Hasegawa H, Ito M. Artificial intelligence-based computer vision in surgery: recent advances and future perspectives. *Ann Gastroenterol Surg.* 2022;6:29-36. DOI PubMed PMC
  121. Bao R, Tao J, Zhao J, Dong M, Li J, Pan C. Integrated intelligent tactile system for a humanoid robot. *Sci Bull.* 2023;68:1027-37. DOI PubMed
  122. Guo X, McFall F, Jiang P, Liu J, Lepora N, Zhang D. A lightweight and affordable wearable haptic controller for robot-assisted microsurgery. *Sensors.* 2024;24:2676. DOI PubMed PMC
  123. Orosco RK, Lurie B, Matsuzaki T, et al. Compensatory motion scaling for time-delayed robotic surgery. *Surg Endosc.* 2021;35:2613-8. DOI PubMed