



Original Article

Open Access



The growing prevalence of nonalcoholic fatty liver disease (NAFLD), determined by fatty liver index, amongst young adults in the United States. A 20-year experience

Wenhao Li^{1,†}, Cheng Han Ng^{2,†} , Jingxuan Quek^{2,†}, Kai En Chan², Caitlyn Tan², Rebecca Wenling Zeng², Jie Ning Yong², Hannah Tay², Darren Jun Hao Tan², Wen Hui Lim², Douglas Chee⁶, Jinyang Ho⁵, Nicholas W.S. Chew^{2,3}, Lung Yi Mak⁴, Mohammad Shadab Siddiqui⁵, Arun Sanyal⁵, William Alazawi¹, Naim Alkouri⁹, Mark Muthiah^{2,6,7,†}, Mazen Nouredin^{8,†} 

¹Blizard Institute, Queen Mary University of London, London E1 2AT, UK.

²MBBS Programme, Yong Loo Lin School of Medicine, National University of Singapore, Singapore 117597, Singapore.

³Department of Cardiology, National University Heart Centre, National University Hospital, Singapore 119074, Singapore.

⁴Department of Medicine, University of Hong Kong, Hong Kong 999077, China.

⁵Division of Gastroenterology, Hepatology and Nutrition, Department of Internal Medicine, Virginia Commonwealth University, Richmond, VA 23284, USA.

⁶Division of Gastroenterology and Hepatology, Department of Medicine, National University Hospital of Singapore, Singapore 119074, Singapore.

⁷National University Centre for Organ Transplantation, National University Health System, Singapore 119228, Singapore.

⁸Houston Liver Institute, Houston, TX 77079, USA.

⁹Texas Liver Institute, San Antonio, TX 78215, USA.

[†]These authors contributed equally to this work and share first authorship.

[†]The authors equally supervise the manuscript.

Correspondence to: Dr. Cheng Han Ng, MBBS Programme, Yong Loo Lin School of Medicine, National University of Singapore, Singapore 10 Medical Dr, Singapore 117597, Singapore. E-mail: chenhanng@gmail.com; Dr. Mazen Nouredin, MD, MHSc, Director, Houston Liver Institute, 1155 Dairy Ashford, Houston, TX 77079, USA. E-mail: noureddinmd@gmail.com

How to cite this article: Li W, Ng CH, Quek J, Chan KE, Tan C, Zeng RW, Yong JN, Tay H, Tan DJH, Lim WH, Chee D, Ho J, Chew NWS, Mak LY, Siddiqui MS, Sanyal A, Alazawi W, Alkouri N, Muthiah M, Nouredin M. The growing prevalence of nonalcoholic fatty liver disease (NAFLD), determined by fatty liver index, amongst young adults in the United States. A 20-year experience. *Metab Target Organ Damage* 2022;2:19. <https://dx.doi.org/10.20517/mtod.2022.24>

Received: 5 Aug 2022 **First Decision:** 26 Aug 2022 **Revised:** 14 Sep 2022 **Accepted:** 21 Oct 2022 **Published:** 2 Nov 2022

Academic Editor: Silvia R. Fargion **Copy Editor:** Peng-Juan Wen **Production Editor:** Peng-Juan Wen

Abstract

Aim: The Global burden of nonalcoholic fatty liver disease (NAFLD) has significantly increased recently, with its prevalence mirroring increasing obesity and diabetes. However, population-specific evidence for young adults



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, sharing, adaptation, distribution and reproduction in any medium or format, for any purpose, even commercially, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.



remains limited. Herein, we provide a 20-year trend analysis of NAFLD in young adults and examine factors associated with NAFLD and major adverse cardiovascular events (MACE) prevalence.

Methods: This study uses data from the United States National Health and Nutrition Examination Survey (NHANES) 1999-2018. Fatty liver was examined with the fatty liver index (FLI) and United States-FLI (US-FLI), and advanced fibrosis was examined with the fibrosis-4 index. Clustered multivariate logistic regression analysis on the year of study was applied to obtain odds ratios (OR) for the estimation of events.

Results: 13.31% (95%CI: 12.71% to 13.94%) of young adults had NAFLD. The prevalence increased from 9.98% in 1999 to 19.49% in 2018, with a statistically significant trend ($P < 0.001$). 9.52% and 5.29% of patients have clinically significant and advanced fibrosis, respectively. In multivariate analysis, diabetes (3.48, 95%CI: 2.37 to 5.11), hypertension (2.03, 95%CI: 1.62 to 2.55), elevated body mass index (1.22, 95%CI: 1.20 to 1.23, $P < 0.001$) significantly increases odds of NAFLD. The largest increase in odds was related to obesity (OR: 21.61, 95%CI: 16.95 to 27.55, $P < 0.001$). Young adults with NAFLD had a borderline non-significant increase in the prevalence of MACE compared to individuals without NAFLD (OR: 1.603, 95%CI: 0.949 to 2.708, $P = 0.078$).

Conclusion: The rising prevalence of NAFLD in young adults depicts the changing landscape of NAFLD and its association with a significant increase in MACE. The challenge of effective risk stratification and education of these individuals remains.

Keywords: NAFLD, prevalence, young Adults, epidemiology

INTRODUCTION

Non-alcoholic fatty liver disease (NAFLD) is the most common cause of chronic liver disease worldwide, affecting approximately 25%-33% of the population^[1-4], and represents a spectrum of liver diseases that ranges from simple liver steatosis to non-alcoholic steatohepatitis (NASH). NASH is the more progressive form of the condition characterized by inflammation, hepatocyte ballooning and hepatocellular injury with or without fibrosis^[5]. The progression of NAFLD is associated with a range of liver-related sequelae such as cirrhosis, liver failure and hepatocellular carcinoma, with the leading cause of mortality being cardiovascular disease (CVD)^[6,7], which shares many remediable cardiometabolic risk factors encapsulated by metabolic syndrome, including obesity, atherogenic dyslipidemia, hypertension and hyperglycaemia^[8]. However, one should consider that there are circumstances in which this association does not apply^[9]; for example, isolated hepatic steatosis without further histological characteristics does not appear to influence stage-dependent CVD risk. Therefore, there is current dissonance on whether NAFLD confers additional independent CVD risk or whether an increase in CVD risk in NAFLD is due to associated CVD risk factors^[10]. Nevertheless, CVD mortality accounts for approximately 40% of all deaths in NAFLD patients^[11,12].

The prevalence of NAFLD in young adults has seen an alarming rise in NAFLD prevalence which mirrors the rising prevalence of obesity and type 2 diabetes (T2DM). Previous studies have shown NAFLD and NASH to have the highest prevalence among patients with DM, as opposed to other components of metabolic syndrome^[9,13]. Mrad *et al.* reported a rise in NAFLD prevalence in young adults in the United States from 9.6% in 1988-1994 to 24.0% in 2005-2010^[14]. Similarly, Lawlor *et al.* and Abeysekera *et al.* reported a rise in NAFLD prevalence from 2.5% to 20.7% among young adults from the Avon Longitudinal Study of Parents and Children (ALSPAC) cohort in the United Kingdom from a mean age of 17.9 to 24^[15,16]. An updated analysis of the epidemic of NAFLD in young adults has to be examined with more recent population data despite recent drastic increases in obesity and diabetes in young adults^[17]. Additionally, the relationship between cardiovascular health and NAFLD in young adults remains limited. Therefore, we

sought to examine the prevalence and risk factors associated with NAFLD and the risk of cardiovascular disease in the young adult population using patients recruited in the United States National Health and Nutrition Examination Survey (NHANES) between 1999-2018.

METHODS

The NHANES study examines aggregated health-related data from a clustered sampled national survey involving general and noninstitutionalised individuals in the United States between 1999-2018. The study involved participants undergoing comprehensive interviews, medical examinations, and laboratory assessments^[18]. Ethics approval by the Institutional Review Board was exempted due to the anonymous nature of the data made publicly available by the National Centre for Health Statistics (NCHS). Baseline characteristics such as but are not limited to age, gender, body mass index (BMI), ethnicity, income levels, low-density lipoprotein (LDL) cholesterol, high-density lipoprotein (HDL) cholesterol, total cholesterol, triglyceride, fasting blood glucose, glycohemoglobin, homeostatic model assessment for insulin resistance (HOMA-IR), platelet count, aspartate aminotransferase (AST), alanine aminotransferase (ALT) and past medical history (diabetes, hypertension, obesity, smoking status) were collected. Information on longitudinal outcomes of patients, including major adverse cardiovascular events (MACE), was also collected.

The definition of NAFLD was adapted based on the American Association for the Study of Liver Disease (AASLD) guidelines for NAFLD^[19]. We defined NAFLD as the presence of steatosis in the absence of substantial alcohol use (≥ 3 drinks a day in men, ≥ 2 drinks a day in women). The presence of steatosis in NAFLD was quantified with either the fatty liver index (FLI) or the United States fatty liver index (US-FLI) with a cut-off of ≥ 60 ^[20] and ≥ 30 ^[21], respectively^[8,22]. Metabolic unhealthy patients were defined as individuals with fatty liver and concomitant type 2 diabetes and/or ≥ 2 metabolic risk abnormalities^[23]: (1) waist circumference ≥ 102 cm in men and 88 cm in women ($\geq 90/80$ cm in Asian men and women); (2) blood pressure $\geq 130/85$ mm Hg or receiving antihypertensives; (3) plasma triglycerides ≥ 2.3 mmol/L or receiving specific drug treatment; (4) plasma high-density lipoprotein < 1.00 mmol/L for men and < 1.30 mmol/L for women or specific drug treatment; (5) fasting blood glucose ≥ 100 mg/dL. Young adults were defined as individuals aged 18 to 30, while obesity status was defined as BMI ≥ 30.0 kg/m² for Caucasians and BMI ≥ 27.5 kg/m² for Asians^[24]. Diabetes was defined as glycohemoglobin $\geq 6.5\%$, fasting plasma glucose ≥ 7 mmol/L, self-reported diabetes or the use of anti-diabetic medications^[25]. Hypertension was defined as a systolic or diastolic blood pressure $\geq 130/85$ or the use of antihypertensive^[26]. Major adverse cardiovascular events (MACE) was an umbrella definition encompassing heart failure events, stroke, myocardial infarction and mortality^[27]. Advanced fibrosis was assessed by fibrosis-4 index (FIB-4), where a cut-off value of FIB < 1.3 ^[28] was defined as having a low risk of advanced fibrosis^[29]. A sensitivity analysis was then conducted to examine the prevalence of NAFLD with vibration-controlled transient elastography (VCTE). The NHANES 2017-2018 cycle was the only completed cycle with VCTE assessment of liver steatosis and fibrosis. A controlled attenuation parameter (CAP) score of ≥ 288 dB/m was selected for assessing liver steatosis^[30]. Clinically significant fibrosis (F2-4) was assessed with a liver stiffness (LSM) of ≥ 8.8 kPa^[31] and an LSM of ≥ 11.7 kPa^[32] for advanced fibrosis, respectively.

All statistical analysis was performed using STATA (16.1). Continuous variables were examined with Wilcoxon ranked sum test and Kruskal-Wallis analysis of variance, while binary variables were examined with chi-square test and Fisher's exact test where appropriate. Univariate and multivariate logistic regression analysis was used to obtain odds ratios (OR) for the estimation of common events and clinical interpretability in NAFLD and non-NAFLD young adults. A cluster analysis was also included based on the year of study to account for relevant heterogeneity introduced, using a cluster variable within multivariate

logistic regression that was constructed with important confounders that include age, gender, ethnicity, diabetes, smoking status and BMI. A trend analysis using the Cochran-Armitage test was also used to describe the relationship between NAFLD prevalence and the year of study.

RESULTS

Prevalence of NAFLD in young adults

In total, a total of 46,094 individuals were included in the analysis. A total of 27.09% (95%CI: 26.69% to 27.50%) had a diagnosis of NAFLD [Figure 1] and the prevalence of NAFLD by year groups is presented in Figure 2. A breakdown of age found a total population of 14,628 were between 18 to 30 years of age, 9228 were 31 to 40 years of age, 8907 were 41 to 50 years of age, 8329 were 51 to 60 years of age, 8517 were 61 to 70 years of age and 9422 were 70 years and older. The prevalence of NAFLD was 13.31% (95%CI: 12.71% to 13.94%), 24.96% (95%CI: 23.99% to 25.97%), 30.87% (95%CI: 29.81% to 31.96%), 35.40% (95%CI: 34.25% to 36.56%), 37.54% (95%CI: 36.39% to 38.70%) and 30.94% (95%CI: 29.83% to 32.07%) respectively in the age groups [Figure 3].

An analysis specific to young adults (18 to 30 years) based on the year of study on the proportion of NAFLD is presented in Figure 4. In total, 13.31% (95%CI: 12.71% to 13.94%) of young adults have NAFLD, of which 29.29% (25.66% to 33.20%) were classified as metabolically unhealthy. In young adults, the prevalence increased from 9.98% to 19.49% from 1999 to 2018. Trend analysis by the Cochran-Armitage test found a significant relationship between the year of study and the prevalence of NAFLD in young adults (Figure 4, $P < 0.001$). A sensitivity analysis was conducted to examine the prevalence of NAFLD based on VCTE only. However, VCTE was only available for the NHANES 2017-2018. A total of 1041 individuals were identified as young adults in the NHANES 2017-2018 cycle and the prevalence of NAFLD amongst these individuals was 18.16% (95%CI: 15.93 to 20.62) with a CAP score of 288 dB/m.

Most of the young adults were at low risk of advanced fibrosis ($FIB4 < 1.30$; 99.03%, 95%CI: 98.40% to 99.94%), with only a small proportion at intermediate-high risk of advanced fibrosis ($FIB4 \geq 1.30$, 0.07%, 95%CI: 0.06% to 1.60%). As FIB-4 is inaccurate in young adults, a sensitivity analysis was conducted to examine the prevalence of NAFLD based on liver stiffness from VCTE. The prevalence of clinically significant fibrosis (F2-4) and advanced fibrosis (F3-4) among young adults with NAFLD was 9.52% (95%CI: 6.06% to 14.65%) and 5.29% (95%CI: 2.86% to 9.59%), respectively.

Associated factors of NAFLD in young adults

A comparative analysis of the baseline characteristics between NAFLD and non-NAFLD is presented in Table 1. NAFLD individuals in young adults were found to be slightly older (24.70 vs. 22.87, $P < 0.001$) without any significant gender differences. Diabetes and hypertension were significantly associated with the presence of NAFLD (0.05 vs. 0.01, $P < 0.001$; 0.18 vs. 0.09, $P < 0.001$, respectively). Measures of lipids, including low-density lipoproteins, triglycerides and total cholesterol, were significantly higher in NAFLD individuals ($P < 0.001$). Similarly, the median liver enzyme levels were higher in NAFLD compared to non-NAFLD individuals ($P < 0.001$). The median BMI of young adults with NAFLD was significantly higher than non-NAFLD individuals (34.20 vs. 24.62, $P < 0.001$) and most of the young adults with NAFLD were found to be obese ($P < 0.001$).

There was a borderline non-statistical difference in ethnicity difference between NAFLD and non-NAFLD ($P = 0.051$). Multivariate logistic regression with a cluster variable on the year of study was used to examine baseline factors associated with NAFLD [Table 1]. Increasing age amongst young adults was a statistically significant factor resulting in NAFLD (OR: 1.11, 95%CI: 1.04 to 1.18, $P = 0.001$). Both BMI and waist

Table 1. Differences in baseline characteristics in young adults aged 18 to 30 with and without NAFLD

	NAFLD (n = 1562)	Non-NAFLD (n = 10,170)	P-value	Multivariate OR	P value
Age	24.70 (IQR: 22.00 to 28.00)	22.87 (IQR: 19.00 to 26.00)	< 0.001	1.11 (95%CI: 1.04 to 1.18)	0.001
Diabetes	0.05 (95%CI: 0.04 to 0.06)	0.01 (95%CI: 0.01 to 0.02)	< 0.001	3.48 (95%CI: 2.37 to 5.11)	< 0.001
BMI (kg/m ²)	34.20 (IQR: 31.09 to 38.70)	24.62 (IQR: 21.80 to 28.49)	< 0.001	1.22 (95%CI: 1.20 to 1.23)	< 0.001
Waist circumference (cm)	109.60 (IQR: 102.70 to 118.70)	85.20 (IQR: 95.40 to 77.30)	< 0.001	1.09 (95%CI: 1.08 to 1.10)	< 0.001
Obesity status					
Non-obese	0.19 (IQR: 0.18 to 0.20)	0.84 (IQR: 0.82 to 0.85)	< 0.001	Ref	
Obese	0.81 (IQR: 0.80 to 0.82)	0.16 (IQR: 0.15 to 0.18)		21.61 (95%CI: 16.95 to 27.55)	< 0.001
HTN	0.18 (95%CI: 0.16 to 0.20)	0.09 (95%CI: 0.08 to 0.09)	< 0.001	2.03 (95%CI: 1.62 to 2.55)	< 0.001
Gender					
			0.396		
Male	0.46 (95%CI: 0.44 to 0.49)	0.47 (95%CI: 0.46 to 0.48)		Ref	
Female	0.54 (95%CI: 0.51 to 0.56)	0.53 (95%CI: 0.52 to 0.54)		0.74 (95%CI: 0.58 to 0.94)	0.012
Platelet count	276.85 (IQR: 231.00 to 315.50)	260.05 (IQR: 216.00 to 296.00)	< 0.001	1.00 (95%CI: 0.99 to 1.01)	0.06
Glycohemoglobin (%)	5.37 (IQR: 5.10 to 5.50)	5.17 (IQR: 4.90 to 5.40)	< 0.001	1.47 (95%CI: 1.33 to 1.61)	< 0.001
HOMA	1.98 (IQR: 1.31 to 3.09)	4.24 (IQR: 2.75 to 6.52)	< 0.001	1.18 (95%CI: 1.12 to 1.23)	< 0.001
Fasting blood glucose (mg/dL)	96.00 (IQR: 89.00 to 103.00)	92.00 (IQR: 86.80 to 98.00)	< 0.001	1.01 (95%CI: 1.00 to 1.01)	0.001
Total cholesterol (mg/dL)	193.75 (IQR: 165.00 to 218.00)	175.41 (IQR: 149.00 to 196.00)	< 0.001	1.01 (95%CI: 1.01 to 1.02)	< 0.001
LDL (mg/dL)	114.13 (IQR: 91.00 to 135.00)	100.80 (IQR: 79.00 to 118.00)	< 0.001	1.01 (95%CI: 1.01 to 1.02)	< 0.001
HDL (mg/dL)	45.97 (IQR: 37.00 to 52.00)	53.94 (IQR: 44.00 to 62.00)	< 0.001	0.95 (95%CI: 0.93 to 0.97)	< 0.001
Triglycerides (mg/dL)	191.81 (IQR: 112.00 to 234.00)	107.87 (IQR: 59.00 to 129.00)	< 0.001	1.01 (95%CI: 1.01 to 1.02)	< 0.001
AST (IU/L)	22.00 (IQR: 18.00 to 28.00)	21.00 (IQR: 18.00 to 25.00)	< 0.001	1.01 (95%CI: 1.00 to 1.01)	0.024
ALT (IU/L)	24.00 (IQR: 17.00 to 37.00)	18.00 (IQR: 14.00 to 25.00)	< 0.001	1.01 (95%CI: 1.00 to 1.01)	< 0.001
Smoking status					
			< 0.001		
Non-smoker	0.73 (95%CI: 0.71 to 0.75)	0.66 (95%CI: 0.65 to 0.67)		Ref	
Former smoker	0.10 (95%CI: 0.09 to 0.12)	0.11 (95%CI: 0.11 to 0.12)		0.77 (95%CI: 0.61 to 0.97)	0.025
Current smoker	0.17 (95%CI: 0.15 to 0.19)	0.23 (95%CI: 0.22 to 0.24)		0.66 (95%CI: 0.56 to 0.76)	< 0.001
Ethnicity					
			0.051		
Mexican American	0.24 (95%CI: 0.22 to 0.27)	0.25 (95%CI: 0.24 to 0.26)		Ref	
Other Hispanic	0.07 (95%CI: 0.06 to 0.09)	0.08 (95%CI: 0.08 to 0.09)		0.94 (95%CI: 0.79 to 1.11)	0.452
Caucasian	0.33 (95%CI: 0.31 to 0.36)	0.35 (95%CI: 0.34 to 0.36)		1.00 (95%CI: 0.84 to 1.19)	0.977
African American	0.26 (95%CI: 0.24 to 0.28)	0.22 (95%CI: 0.21 to 0.23)		0.95 (95%CI: 0.76 to 1.19)	0.644
Others	0.09 (95%CI: 0.08 to 0.11)	0.09 (95%CI: 0.09 to 0.10)		1.36 (95%CI: 1.11 to 1.67)	0.003
Income levels					
			0.032		
Below 10,000	0.08 (95%CI: 0.07 to 0.10)	0.11 (95%CI: 0.10 to 0.12)		Ref	
10,000 to 24,999	0.25 (95%CI: 0.23 to 0.28)	0.26 (95%CI: 0.25 to 0.27)		1.23 (95%CI: 1.00 to 1.50)	0.045
25,000 to 44,999	0.28 (95%CI: 0.25 to 0.30)	0.26 (95%CI: 0.25 to 0.27)		1.25 (95%CI: 1.03 to 1.51)	0.023
45,000 to 74,999	0.24 (95%CI: 0.22 to 0.27)	0.23 (95%CI: 0.22 to 0.24)		1.21 (95%CI: 0.96 to 1.53)	0.114
Above 75,000	0.14 (95%CI: 0.13 to 0.16)	0.13 (95%CI: 0.13 to 0.15)		1.21 (95%CI: 1.00 to 1.46)	0.052

Bolded P-value ≤ 0.05 denotes statistical significance. NAFLD: Non-alcoholic fatty liver disease; OR: odds ratio; BMI: body mass index; LDL: low-density lipoprotein; HDL: high-density lipoprotein; HTN: hypertension; HOMA: homeostatic model assessment; AST: aspartate aminotransferase; ALT: alanine transaminase; IQR: interquartile range; 95%CI: 95% confidence interval.



Figure 1. Prevalence of NAFLD in the total and young adult population. NAFLD: Nonalcoholic fatty liver disease.




Figure 2. Prevalence of NAFLD by year of study. NAFLD: Nonalcoholic fatty liver disease.

circumference significantly increase the odds of NAFLD (1.22, 95%CI: 1.20 to 1.23, $P < 0.001$ and 1.09, 95%CI: 1.08 to 1.10, $P < 0.001$). Additionally, the female gender was less likely to be associated with NAFLD compared to the male gender (OR: 0.74, 95%CI: 0.58 to 0.94, $P = 0.012$) and lipid dysregulation significantly affected the odds of NAFLD [Table 1]. Interestingly, smoking history was associated with reduced odds of NAFLD as with higher income levels [Table 1]. The presence of diabetes affected only 5% of NAFLD individuals but resulted in a large increase in the odds of NAFLD (OR: 3.48, 95%CI: 2.37 to 5.11, $P < 0.001$). The largest increase in odds of NAFLD in young adults, however, was related to the presence of obesity (OR: 21.61, 95%CI: 16.95 to 27.55, $P < 0.001$)

MACE events in young adults with NAFLD

Young adults with NAFLD were then examined for an increased association between MACE and cardiovascular risk. MACE events were defined as any events of stroke, heart failure, myocardial infarction and cardiovascular mortality based on ICD. There were a total of 15 MACE events in 1416 NAFLD individuals and 41 MACE events in 6978 non-NAFLD individuals. An unadjusted clustered logistic regression found a significant increase in MACE events with young adults with NAFLD (OR: 1.81, 95%CI: 1.05 to 3.13, $P = 0.033$). A multivariate clustered logistic regression was then conducted, adjusting for age, gender, BMI, race, diabetes, and smoking status, and found a borderline non-significant increase in odds of




Figure 3. Prevalence of NAFLD by age group. NAFLD: Nonalcoholic fatty liver disease.

MACE events with NAFLD (OR: 1.59, 95%CI: 0.95 to 2.67, $P = 0.081$).

DISCUSSION

The epidemic of NAFLD has been well described in general adults and shown to significantly increase the risk of mortality^[33]. A lesser-known entity, however, lies in the prevalence of NAFLD in young adults who are often overlooked. In this study, we show that the overall prevalence of NAFLD in young adults (aged between 18 to 30) in this population database from the United States is estimated to be 13.31%. Within the young adult population, the proportion of NAFLD diagnoses doubled from 9.98% in 1999-2000 to 19.49% in 2017-2018 using FLI. Sensitivity analysis by VCTE with the NHANES 2017-2018 cycle similarly found that 18.16% (95%CI: 15.93 to 20.62) of young adults were affected by NAFLD. The findings are congruent with previous studies^[34,14] and emphasize the growing prevalence of NAFLD in young adults in the United States. However, the current study also shows the potential of NAFLD in young adults that may predispose the onset of MACE events (OR: 1.603, 95%CI: 0.949 to 2.708, $P = 0.078$), albeit with borderline non-significance.

There was an evident association between metabolic dysfunction where young adults with NAFLD had a higher prevalence of obesity, diabetes, hypertension, obesity, insulin resistance, dyslipidemia, and abnormal liver function test results. However, the largest increase in odds of NAFLD was found to be from obesity which had a 21 times increase in odds of NAFLD. Nevertheless, while an estimated 40%-45%^[8] of NAFLD are associated with diabetes and HTN, only 5% and 18% of young adults with NAFLD had HTN and diabetes. Significantly, only 29.29% of the young adult with NAFLD had metabolic dysfunction and the higher odds of NAFLD may relate to metabolically healthy obesity (MHO) status. MHO is a unique subtype




Figure 4. Prevalence of NAFLD in young adults by year of study. NAFLD: Nonalcoholic fatty liver disease.

of obesity more frequently found in younger individuals without significant metabolic dysfunction despite the presence of obesity^[35,36] which has been shown to predispose the risk of NAFLD^[37]. Additionally, the onset of NAFLD in young adults before evidence of other systemic dysregulation also alludes to the potential of fatty liver driving the development of other metabolic diseases. Only 5% and 18% of young adults with NAFLD have HTN or diabetes, respectively, which is a common commodity of NAFLD^[8,22], and NAFLD has been shown to longitudinally increase the risk of prediabetes and diabetes^[8]. This potentially alludes that the presence of hepatic fat may not be a bystander but rather a systemic driver of other metabolic disease^[38].

Female gender and smokers are associated with reduced odds of NAFLD diagnosis, and prior studies have attributed the protective effects of estrogen in pre-menopausal females to be protective towards NAFLD^[39]. Additionally, it was interesting to find that NAFLD individuals were less likely to be smokers. However, the negative association between NAFLD and smoking should be interpreted with caution as previous studies have reported contrasting associations between smoking and NAFLD^[40-42], and the effect of smoking may be too premature to manifest in young adults. Interestingly, lower income levels are also significantly associated with NAFLD diagnosis. This is in line with previous studies by Giammarino *et al.* and Golovaty *et al.* in predominantly older adults, which demonstrated the strong association between socioeconomic factors, including food insecurity and NAFLD in the United States^[43,44]. The availability, affordability and recognition of healthy food options and lifestyles need to be addressed in young adults, particularly those with lower income, to tackle the growing prevalence of NAFLD^[45].

Although cardiovascular events are rare in individuals below the age of 30, interestingly, there were a borderline 1.6 times odds of a non-significant increase in MACE events in young adults with NAFLD. A recently published long-term mortality data by Simon *et al.* showed that children and young adults with biopsy-proven NAFLD showed a significantly increased risk of long-term CVD-related mortality in NAFLD^[46]. However, risk stratification tools such as Framingham Heart Score, which are used to predict future MACE in older adults, perform poorly in individuals under 30^[47]. Therefore, there remains an unmet need to develop prognostication tools to identify individuals at risk of future cardiovascular disease in young adults with NAFLD.

Limitations

The current study has several limitations. NAFLD diagnosis was made through non-invasive means relying on FLI as a surrogate measure of steatosis but was deemed suitable considering the size of this population study. Additionally, FIB-4 score performs poorly in NAFLD patients under 35^[48]; therefore, the estimation of fibrosis may be inaccurate in the current study. A sensitivity analysis with VCTE instead showed that 5%-10% are likely to have clinically significant and advanced fibrosis. The population data was prevalent and did not collect the age of the first presentation of MACE among patients. Lastly, longitudinal outcomes of mortality were not assessed due to the significant duration of time needed for events to occur in young adults.

In conclusion, the prevalence of NAFLD is rising rapidly among the young adult population in the United States, with significant association with the obesity epidemic in young adults and the presence of which is associated with increased odds of MACE events. However, risk stratification of young adults with NAFLD remains challenging due to the lack of stratification tools in individuals below 35 and awareness of the disease can be significantly lower in this population.

DECLARATIONS

Authors' contributions

Conceptualisation and design: Li W, Ng CH, Muthiah M, Nouredin M

Acquisition of data: Quek J, Chan KE, Tan C, Zeng RW, Yong JN, Tay H, Tan DJH, Lim WH

Analysis and interpretation of data: Li W, Ng CH, Quek J, Chan KE, Tan C, Zeng RW, Yong JN, Tay H, Tan DJH, Lim WH

Writing - original draft: Li W, Ng CH, Quek J, Chan KE, Tan C

Writing - review & editing: Chee D, Ho J, Chew NWS, Mak LY, Siddiqui MS, Sanyal A, Alazawi W, Alkouri N, Muthiah M, Nouredin M

All authors approved the final version of the manuscript, including the authorship list and agreed to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Availability of data and materials

Data were retrieved from the National Health and Nutrition Examination Survey Registry.

Financial support and sponsorship

None.

Conflicts of interests

Sanyal A is President of Sanyal Biotechnology and has stock options in Genfit, Akarna, Tiziana, Indalo, Durect and Galmed. He has served as a consultant to Astra Zeneca, Nitto Denko, Enyo, Ardelyx, Conatus, Nimbus, Amarin, Salix, Tobira, Takeda, Janssen, Gilead, Terns, Birdrock, Merck, Valeant, Boehringer-

Ingelheim, Lilly, Hemoshear, Zafgen, Novartis, Novo Nordisk, Pfizer, Exhalenz and Genfit. He has been an unpaid consultant to Intercept, Echosens, Immuron, Galectin, Fractyl, Syntlogis, Affimune, Chemomab, Zydus, Nordic Bioscience, Albireo, Prosciento, Surrozen and Bristol Myers Squibb. His institution has received grant support from Gilead, Salix, Tobira, Bristol Myers, Shire, Intercept, Merck, Astra Zeneca, Malinckrodt, Cumberland and Novartis. He receives royalties from Elsevier and UptoDate. Nouredin M has been on the advisory board/consultant for 89BIO, Altimune, Gilead, cohBar, Cytodyn, Intercept, Pfizer, Novo Nordisk, Blade, EchoSens, Fractyl, Madrgial, NorthSea, Prespecturm, Terns, Siemens and Roche diagnostic; He has received research support from Allergan, BMS, Gilead, Galmed, Galectin, Genfit, Conatus, Enanta, Madrigal, Novartis, Pfizer, Shire, Viking and Zydus; He is a shareholder or has stocks in Anaetos, Chrownwell, Ciema, Rivus Pharma and Viking. All other authors declared that there are no conflicts of interest.

Ethical approval and consent to participate

Ethics approval by the Institutional Review Board was exempted due to the anonymous nature of the data made publicly available by the National Centre for Health Statistics (NCHS). Simultaneously, it does not require the author's informed permission..

Consent for publication

Not applicable.

Copyright

The Author(s) 2022.

REFERENCES

1. Paik JM, Kabbara K, Eberly KE, Younossi Y, Henry L, Younossi ZM. Global burden of NAFLD and chronic liver disease among adolescents and young adults. *Hepatology* 2022;75:1204-17. [DOI PubMed](#)
2. Sanyal AJ. Past, present and future perspectives in nonalcoholic fatty liver disease. *Nat Rev Gastroenterol Hepatol* 2019;16:377-86. [DOI PubMed](#)
3. Lim GEH, Tang A, Ng CH, et al. An observational data meta-analysis on the differences in prevalence and risk factors between MAFLD vs. NAFLD. *Clin Gastroenterol Hepatol* 2021;11:S1542-3565(21)01276. [DOI PubMed](#)
4. Le MH, Yeo YH, Li X, et al. 2019 global NAFLD prevalence: a systematic review and meta-analysis. *Clin Gastroenterol Hepatol* ;2021:S1542-3565(21)01280. [DOI PubMed](#)
5. Muthiah MD, Cheng Han N, Sanyal AJ. A clinical overview of non-alcoholic fatty liver disease: a guide to diagnosis, the clinical features, and complications-What the non-specialist needs to know. *Diabetes Obes Metab* 2022;24 Suppl 2:3-14. [DOI PubMed](#)
6. Targher G, Byrne CD, Tilg H. NAFLD and increased risk of cardiovascular disease: clinical associations, pathophysiological mechanisms and pharmacological implications. *Gut* 2020;69:1691-705. [DOI PubMed](#)
7. Chew NWS, Ng CH, Muthiah MD, Sanyal AJ. Comprehensive review and updates on holistic approach towards non-alcoholic fatty liver disease management with cardiovascular disease. *Curr Atheroscler Rep* 2022;24:515-32. [DOI PubMed](#)
8. Ng CH, Chan KE, Chin YH, et al. The effect of diabetes and prediabetes on the prevalence, complications and mortality in non-alcoholic fatty liver disease. *Clin Mol Hepatol* 2022;28:565-74. [DOI PubMed PMC](#)
9. Stefan N, Cusi K. A global view of the interplay between non-alcoholic fatty liver disease and diabetes. *Lancet Diabetes Endocrinol* 2022;10:284-96. [DOI PubMed](#)
10. Francque SM, van der Graaff D, Kwanten WJ. Non-alcoholic fatty liver disease and cardiovascular risk: pathophysiological mechanisms and implications. *J Hepatol* 2016;65:425-43. [DOI PubMed](#)
11. Ekstedt M, Hagström H, Nasr P, et al. Fibrosis stage is the strongest predictor for disease-specific mortality in NAFLD after up to 33 years of follow-up. *Hepatology* 2015;61:1547-54. [DOI PubMed](#)
12. Angulo P, Kleiner DE, Dam-Larsen S, et al. Liver fibrosis, but no other histologic features, is associated with long-term outcomes of patients with nonalcoholic fatty liver disease. *Gastroenterology* 2015;149:389-97.e10. [DOI PubMed PMC](#)
13. Younossi ZM, Golabi P, de Avila L, et al. The global epidemiology of NAFLD and NASH in patients with type 2 diabetes: a systematic review and meta-analysis. *J Hepatol* 2019;71:793-801. [DOI PubMed](#)
14. Mrad RA, Merjaneh N, Mubarak G, Lopez R, Zein NN, Alkhoury N. The increasing burden of nonalcoholic fatty liver disease among young adults in the United States: a growing epidemic. *Hepatology* 2016;64:1386-7. [DOI PubMed](#)
15. Lawlor DA, Callaway M, Macdonald-Wallis C, et al. Nonalcoholic fatty liver disease, liver fibrosis, and cardiometabolic risk factors in adolescence: a cross-sectional study of 1874 general population adolescents. *J Clin Endocrinol Metab* 2014;99:E410-7. [DOI PubMed PMC](#)

16. Abeysekera KWM, Fernandes GS, Hammerton G, et al. Prevalence of steatosis and fibrosis in young adults in the UK: a population-based study. *Lancet Gastroenterol Hepatol* 2020;5:295-305. DOI PubMed PMC
17. Angi A, Chiarelli F. Obesity and diabetes: a sword of damocles for future generations. *Biomedicines* 2020;8:478. DOI PubMed PMC
18. National Health and Nutrition Examination Survey (NHANES) MEC Laboratory Procedures Manual. Centers for disease control and prevention; 2017. Available from: https://www.cdc.gov/nchs/data/nhanes/2017-2018/manuals/2017_MEC_Laboratory_Procedures_Manual.pdf [Last accessed on 28 Oct 2022].
19. Ando Y, Jou JH. Nonalcoholic fatty liver disease and recent guideline updates. *Clin Liver Dis (Hoboken)* 2021;17:23-8. DOI PubMed PMC
20. Ruhl CE, Everhart JE. Fatty liver indices in the multiethnic United States National Health and Nutrition Examination Survey. *Aliment Pharmacol Ther* 2015;41:65-76. DOI PubMed
21. Bedogni G, Bellentani S, Miglioli L, et al. The fatty liver index: a simple and accurate predictor of hepatic steatosis in the general population. *BMC Gastroenterol* 2006;6:33. DOI PubMed PMC
22. Ng CH, Wong ZY, Chew NWS, et al. Hypertension is prevalent in non-alcoholic fatty liver disease and increases all-cause and cardiovascular mortality. *Front Cardiovasc Med* 2022;9:942753. DOI PubMed PMC
23. Alberti KG, Eckel RH, Grundy SM, et al. Harmonizing the metabolic syndrome: a joint interim statement of the International Diabetes Federation Task Force on Epidemiology and Prevention; National Heart, Lung, and Blood Institute; American Heart Association; World Heart Federation; International Atherosclerosis Society; and International Association for the Study of Obesity. *Circulation* 2009;120:1640-5. DOI PubMed
24. Expert Consultation. Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies. *Lancet* 2004;363:157-63. DOI PubMed
25. Punthakee Z, Goldenberg R, Katz P; Diabetes Canada Clinical Practice Guidelines Expert Committee. Definition, classification and diagnosis of diabetes, prediabetes and metabolic syndrome. *Can J Diabetes* 2018;42 Suppl 1:S10-5. DOI PubMed
26. Whelton PK, Carey RM, Aronow WS, et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA guideline for the prevention, detection, evaluation, and management of high blood pressure in adults: a report of the american college of cardiology/american heart association task force on clinical practice guidelines. *Hypertension* 2018;71:e13-e115. DOI PubMed
27. Bosco E, Hsueh L, McConeghy KW, Gravenstein S, Saade E. Major adverse cardiovascular event definitions used in observational analysis of administrative databases: a systematic review. *BMC Med Res Methodol* 2021;21:241. DOI PubMed PMC
28. Blanco-Grau A, Gabriel-Medina P, Rodriguez-Algarra F, et al. Assessing liver fibrosis using the FIB4 index in the community setting. *Diagnostics (Basel)* 2021;11:2236. DOI PubMed PMC
29. Mózes FE, Lee JA, Selvaraj EA, et al; LITMUS Investigators. Diagnostic accuracy of non-invasive tests for advanced fibrosis in patients with NAFLD: an individual patient data meta-analysis. *Gut* 2022;71:1006-19. DOI PubMed PMC
30. Caussy C, Alquraish MH, Nguyen P, et al. Optimal threshold of controlled attenuation parameter with MRI-PDFF as the gold standard for the detection of hepatic steatosis. *Hepatology* 2018;67:1348-59. DOI PubMed PMC
31. Ziol M, Handra-Luca A, Kettaneh A, et al. Noninvasive assessment of liver fibrosis by measurement of stiffness in patients with chronic hepatitis C. *Hepatology* 2005;41:48-54. DOI PubMed
32. Campos-Murguía A, Ruiz-Margáin A, González-Regueiro JA, Macías-Rodríguez RU. Clinical assessment and management of liver fibrosis in non-alcoholic fatty liver disease. *World J Gastroenterol* 2020;26:5919-43. DOI PubMed PMC
33. Simon TG, Roelstraete B, Khalili H, Hagström H, Ludvigsson JF. Mortality in biopsy-confirmed nonalcoholic fatty liver disease: results from a nationwide cohort. *Gut* 2021;70:1375-82. DOI PubMed PMC
34. Arshad T, Paik JM, Biswas R, Alqahtani SA, Henry L, Younossi ZM. Nonalcoholic fatty liver disease prevalence trends among adolescents and young adults in the United States, 2007-2016. *Hepatol Commun* 2021;5:1676-88. DOI PubMed PMC
35. Tsatsoulis A, Paschou SA. Metabolically healthy obesity: criteria, epidemiology, controversies, and consequences. *Curr Obes Rep* 2020;9:109-20. DOI PubMed
36. Chang Y, Jung HS, Cho J, et al. Metabolically healthy obesity and the development of nonalcoholic fatty liver disease. *Am J Gastroenterol* 2016;111:1133-40. DOI PubMed
37. Vusirikala A, Thomas T, Bhala N, Tahrani AA, Thomas GN, Nirantharakumar K. Impact of obesity and metabolic health status in the development of non-alcoholic fatty liver disease (NAFLD): a United Kingdom population-based cohort study using the health improvement network (THIN). *BMC Endocr Disord* 2020;20:96. DOI PubMed PMC
38. Yki-järvinen H. Non-alcoholic fatty liver disease as a cause and a consequence of metabolic syndrome. *Lancet Diabetes Endocrinol* 2014;2:901-10. DOI PubMed
39. Lonardo A, Nascimbeni F, Ballestri S, et al. Sex differences in nonalcoholic fatty liver disease: state of the art and identification of research gaps. *Hepatology* 2019;70:1457-69. DOI PubMed PMC
40. Shen H, Peng JL, Tayarachakul S, Liangpunsakul S. Association between serum cotinine level and prevalence of non-alcoholic fatty liver disease: a cross-sectional study from the Third National Health and Nutrition Examination Survey. *J Investig Med* 2017;65:43-8. DOI PubMed PMC
41. Liu Y, Dai M, Bi Y, et al. Active smoking, passive smoking, and risk of nonalcoholic fatty liver disease (NAFLD): a population-based study in China. *J Epidemiol* 2013;23:115-21. DOI PubMed PMC
42. Caballería L, Pera G, Auladell MA, et al. Prevalence and factors associated with the presence of nonalcoholic fatty liver disease in an adult population in Spain. *Eur J Gastroenterol Hepatol* 2010;22:24-32. DOI PubMed

43. Giammarino AM, Qiu H, Bulsara K, et al. Community socioeconomic deprivation predicts nonalcoholic steatohepatitis. *Hepatology* 2022;6:550-60. [DOI](#) [PubMed](#) [PMC](#)
44. Golovaty I, Tien PC, Price JC, Sheira L, Seligman H, Weiser SD. Food insecurity may be an independent risk factor associated with nonalcoholic fatty liver disease among low-income adults in the United States. *J Nutr* 2020;150:91-8. [DOI](#) [PubMed](#) [PMC](#)
45. Rich NE, Nouredin M, Kanwal F, Singal AG. Racial and ethnic disparities in non-alcoholic fatty liver disease in the USA. *Lancet Gastroenterol Hepatol* 2021;6:422-4. [DOI](#) [PubMed](#)
46. Simon TG, Roelstraete B, Hartjes K, et al. Non-alcoholic fatty liver disease in children and young adults is associated with increased long-term mortality. *J Hepatol* 2021;75:1034-41. [DOI](#) [PubMed](#) [PMC](#)
47. Berry JD, Lloyd-Jones DM, Garside DB, Greenland P. Framingham risk score and prediction of coronary heart disease death in young men. *Am Heart J* 2007;154:80-6. [DOI](#) [PubMed](#) [PMC](#)
48. McPherson S, Hardy T, Dufour JF, et al. Age as a confounding factor for the accurate non-invasive diagnosis of advanced NAFLD fibrosis. *Am J Gastroenterol* 2017;112:740-51. [DOI](#) [PubMed](#) [PMC](#)