

Review

Open Access



Non-coding RNA and drug resistance in head and neck cancer

Yulong Zhang^{1,#}, Yingming Peng^{1,#}, Bingqin Lin^{1,2,#}, Shuai Yang³, Feiqiang Deng^{1,2}, Xuan Yang¹, An Li¹, Wanyi Xia¹, Chenxi Gao¹, Shaona Lei¹, Wei Liao⁴, Qi Zeng^{1,2}

¹Cancer Center, The Fifth Affiliated Hospital of Sun Yat-sen University, Zhuhai 519000, Guangdong, China.

²Key Laboratory of Traditional Chinese Medicine for the Prevention and Treatment of Infectious Diseases, Traditional Chinese Medicine Bureau of Guangdong Province, the Fifth Affiliated Hospital, Sun Yat-sen University, Zhuhai 519000, Guangdong, China.

³Department of Radiotherapy and Minimally Invasive Surgery, Cancer Center, The Fifth Affiliated Hospital of Sun Yat-sen University, Zhuhai 519000, Guangdong, China.

⁴Department of Otolaryngology and Head and Neck Surgery, The Fifth Affiliated Hospital of Sun Yat-sen University, Zhuhai 519000, Guangdong, China.

#Authors contributed equally.

Correspondence to: Prof. Zeng Qi, Cancer Center, The Fifth Affiliated Hospital of Sun Yat-sen University, Meihua East Street, Zhuhai 519000, Guangdong, China. E-mail: zengqi37@mail.sysu.edu.cn; Wei Liao, Department of Otolaryngology and Head and Neck Surgery, The Fifth Affiliated Hospital of Sun Yat-sen University, Meihua East Street, Zhuhai 519000, Guangdong, China. E-mail: liaowei23@mail.sysu.edu.cn

How to cite this article: Zhang Y, Peng Y, Lin B, Yang S, Deng F, Yang X, Li A, Xia W, Gao C, Lei S, Liao W, Zeng Q. Non-coding RNA and drug resistance in head and neck cancer. *Cancer Drug Resist* 2024;7:34. <https://dx.doi.org/10.20517/cdr.2024.59>

Received: 14 May 2024 **First Decision:** 1 Aug 2024 **Revised:** 17 Aug 2024 **Accepted:** 29 Aug 2024 **Published:** 20 Sep 2024

Academic Editors: Hailin Tang, Godefridus J. Peters **Copy Editor:** Dong-Li Li **Production Editor:** Dong-Li Li

Abstract

Head and neck cancer (HNC) is ranked as the sixth most common malignant tumor, and the overall survival rate with current treatment options remains concerning, primarily due to drug resistance that develops following antitumor therapy. Recent studies indicate that non-coding RNAs play a crucial role in drug resistance among HNC patients. This article systematically reviews the current research landscape, explores novel targets and treatment strategies related to non-coding RNAs and HNC resistance, raises some unresolved issues, and discusses five promising research directions in this field: ferroptosis, nanomedicine, exosomes, proteolysis-targeting chimeras (PROTACs), and artificial intelligence. We hope that our work will contribute to advancing research on overcoming HNC resistance through the regulation of non-coding RNAs.

Keywords: Head and neck cancer, non-coding RNA, microRNA, long non-coding RNA, circular RNA, drug resistance



© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, sharing, adaptation, distribution and reproduction in any medium or format, for any purpose, even commercially, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.



INTRODUCTION

Head and neck cancer (HNC) ranks as the sixth most prevalent malignant neoplasm, presenting significant challenges in terms of treatment^[1]. It affects various anatomical sites including the lips, oral cavity, pharynx, larynx, nose, salivary glands, and thyroid. Squamous cell carcinoma (SCC) and its variants constitute more than 90% of the histopathological types^[2]. Major risk factors associated with HNC include tobacco and alcohol use, as well as infections with human papillomavirus (HPV) and Epstein-Barr virus (EBV)^[3]. Globally, the incidence of HNC has been gradually declining, primarily attributed to the reduction in tobacco use and lifestyle modifications^[4,5].

Treatment decisions are guided by the precise location, stage, and pathologic characteristics of the disease. For around 30%-40% of patients diagnosed with early-stage disease (Stage I or II), the typical recommendation involves a single treatment modality such as surgery or radiotherapy. Conversely, roughly 60% of patients presenting with locally or regionally advanced disease upon diagnosis typically receive a multidisciplinary approach encompassing a combination of treatments. These may involve surgery, radiotherapy, chemotherapy, and immunotherapy, as well as additional measures such as nutritional support, psychological counseling, supportive care, and rehabilitation^[2]. Various studies have indicated that the overall 5-year survival rate for individuals with HNC falls below 50%, underscoring the persisting challenges in current treatment outcomes^[2,6]. The inadequate response of patients to antitumor therapy, and in some cases, the development of drug resistance, contribute to unsatisfactory clinical outcomes and stand as significant factors leading to mortality^[7].

Unfortunately, the key determinants underlying this resistance phenomenon remain largely elusive. Advances in molecular biology and gene sequencing have unveiled that approximately 98% of human DNA is designated non-protein coding^[8]. Non-coding RNAs constitute a diverse group of RNA transcripts that lack protein-coding potential. Significant subtypes include microRNAs (miRNAs), circular RNAs (circRNAs), and long non-coding RNAs (lncRNAs)^[9]. Emerging research has revealed the potential of non-coding RNAs to modulate multiple facets of cellular functions, encompassing growth, proliferation, differentiation, development, metabolism, infection, immunity, cell death, organelle biogenesis, messenger signaling, DNA repair, and self-renewal^[10-12]. Moreover, non-coding RNAs exhibit close associations with various common diseases, particularly cancer^[9,13-15]. Additionally, non-coding RNAs function as widespread regulators of various cancer-related characteristics, including proliferation, apoptosis, invasion, metastasis, and genomic instability, thus playing a crucial role in mediating resistance to different cancer therapies^[16,17]. Consequently, gaining a comprehensive understanding of the mechanisms underlying non-coding RNAs and drug resistance holds significant importance in the context of HNC treatment. In addition, the research on overcoming tumor drug resistance through the regulation of non-coding RNAs offers many advantages^[16,18-28] [Table 1].

This review aims to provide a comprehensive summary of the intricate relationship between non-coding RNAs and drug resistance in HNC. It delves into the various types of non-coding RNAs and their potential roles in mediating resistance to different therapeutic approaches utilized in HNC treatment. Through an analysis of the current literature, this review seeks to enhance our understanding of the mechanisms underlying drug resistance in HNC and identify potential avenues for improving treatment outcomes.

miRNAs IN HNC CELL DRUG RESISTANCE

miRNAs, approximately 22 nucleotides in length, constitute a class of endogenous non-coding RNAs. These molecules regulate diverse biological processes, including cell proliferation, differentiation, and apoptosis, through specific binding to the 3'-untranslated region (3'-UTR) of target gene mRNAs, thereby inducing

Table 1. Advantages of targeting non-coding RNAs to overcome tumor drug resistance

Aspect	Summary	Ref.
Complexity of regulatory networks	ncRNAs form intricate regulatory networks that influence various biological processes, including gene expression, signal transduction, and the cell cycle. Targeting these nodes can impact tumor growth and drug resistance	[18]
Specific role	ncRNAs may be uniquely expressed in tumor cells or play critical roles in tumor development and drug resistance, enabling the development of targeted therapies	[19]
Multi-target intervention	ncRNAs can affect multiple signaling pathways and biological processes simultaneously, making them suitable for multi-target interventions to address multidrug resistance	[20]
Drug tolerance	ncRNAs may have a lower risk of tolerance development compared to protein targets due to their distinct regulatory mechanisms	[21]
Diverse treatment strategies	ncRNAs can act as direct drug targets, carriers, or therapeutic agents, offering a range of treatment strategies	[16]
New drug discovery	Research on ncRNAs provides new molecular targets for drug development, aiding in the discovery of novel antitumor agents	[22]
Personalized medicine	ncRNA-based biomarkers can guide personalized treatment plans, helping physicians select the most suitable therapy for individual patients	[23-25]
Combination therapy	ncRNAs can be combined with other therapies (e.g., chemotherapy, radiotherapy, immunotherapy) to improve efficacy and reduce drug resistance	[26]
Treatment monitoring and evaluation	ncRNA expression levels can be used as biomarkers to monitor treatment efficacy, tumor response, and drug resistance development	[27]
Gene editing applications	Gene editing technologies like CRISPR/Cas9 offer tools for targeting ncRNAs, enabling precise regulation of their expression and influencing drug resistance	[28]

ncRNAs: Non-coding RNAs.

mRNA degradation or inhibiting translation^[29-31]. Numerous studies have demonstrated the involvement of miRNA dysregulation in the development and chemoresistance of HNC^[32-34].

miRNAs dysregulation is closely related to chemoresistance in HNC

Research indicates a close association between miRNA dysregulation and chemoresistance in HNC. Zhang *et al.* observed the downregulation of miR-216a-5p and ZEB1 in laryngeal squamous cell carcinoma (LSCC) tissues. Moreover, they demonstrated that overexpression of miR-216a-5p could reverse the malignant phenotype and cisplatin (CDDP) resistance of LSCC cells by targeting ZEB1^[35]. Gao *et al.* identified that circ_0109291 promoted CDDP resistance in oral squamous cell carcinoma (OSCC) cells. They elucidated its mechanism by showing that circ_0109291 sponged miR-188-3p, leading to upregulation of ABCB1 expression^[36]. Cao *et al.* revealed that HOTAIR induced CDDP resistance in nasopharyngeal carcinoma (NPC) by sponging miR-106a-5p, consequently upregulating SOX4 expression^[37]. The miR-200 family constitutes a pivotal group of miRNAs implicated in the regulation of epithelial-mesenchymal transition (EMT). Their downregulation is closely linked to chemoresistance in HNC. For instance, reduced expression of Let-7, miR-200, and miR-203 correlates with docetaxel resistance in OSCC^[38-42]. Moreover, the miR-200c/c-myc negative feedback loop orchestrates EMT, stemness, and chemoresistance in NPC^[43,44]. Additionally, dysregulation of miR-138, miR-222, miR-101, miR-23a, miR-214, and various other miRNAs has been associated with chemoresistance in HNC^[45-47]. In summary, the dysregulation of specific miRNAs plays a key role in the acquired chemoresistance of HNC.

miRNAs affect chemoresistance by regulating autophagy signaling pathways

Autophagy and miRNAs are intricately linked to chemoresistance in HNC. H19 induces CDDP resistance in LSCC cells by upregulating the expression of autophagy-related proteins Atg5 and Beclin1 through the miR-107/HMGB1 axis^[48]. MiR-155 inhibitor-loaded exosomes reverse CDDP resistance in OSCC by inducing autophagy through the upregulation of FOXO3a^[49]. Yang *et al.* found that upregulated miR-214 expression could inhibit autophagy in OSCC cells by targeting autophagy-related genes ULK1 and ATG5,

consequently enhancing chemosensitivity^[50]. CircAP1M2 promotes CDDP resistance and autophagy in OSCC by inhibiting miR-1249-3p to upregulate ATG9A expression^[51]. MiR-1278 inhibits autophagy and CDDP resistance in NPC cells by targeting ATG2B^[52]. Therefore, targeting autophagy pathways through the regulation of miRNAs may enhance the therapeutic sensitivity of HNC.

miRNAs affect chemoresistance by regulating the cell cycle and metabolism

miRNAs can also modulate the sensitivity of tumor cells to treatment by regulating energy metabolism and cell cycle. Fan *et al.* discovered that downregulation of mitochondrial miR-2392 expression could lead to impaired mitochondrial oxidative phosphorylation by inhibiting MT-CO3 translation, resulting in a shift of glucose metabolism from oxidative phosphorylation to anaerobic glycolysis, ultimately causing CDDP resistance in tongue squamous cell carcinoma (TSCC)^[53]. Chen *et al.* discovered that miR-5787 contributes to CDDP resistance in TSCC by reprogramming glucose metabolism via inhibiting the translation of MT-CO3^[54]. In another study, Mortezaagholi *et al.* demonstrated that miR-34 could reverse paclitaxel resistance in OECM-1 oral cancer cells by inducing DNA damage and apoptosis in a p53-dependent manner^[55]. HOXA11-AS enhances the proliferation, invasion, survival, and drug resistance of OSCC by sponging miR-494 to promote NQO1 expression and recruiting EZH2 to the NQO2 promoter to suppress NQO2 expression^[56]. Kang *et al.* demonstrated that cancer-associated fibroblast (CAF)-derived extracellular vesicles carrying miR-876-3p can modulate CDDP resistance in OSCC by targeting insulin-like growth factor binding protein 3 (IGFBP3)^[57]. Li *et al.* identified miR-194 as a tumor suppressor in LSCC that inhibits cell proliferation and enhances chemosensitivity by targeting Wee1^[58].

miRNAs affect chemoresistance via exosome-mediated mechanisms

Exosomes, as an important mediator of intercellular communication, can carry functional molecules such as miRNAs to mediate information exchange between tumor cells and the microenvironment, playing a key role in tumor progression and chemoresistance^[59-61]. Qin *et al.* found that CAF-derived exosomal miR-196a could confer CDDP resistance to head and neck squamous carcinoma cells by targeting CDKN1B and ING5^[62]. Li *et al.* discovered that exosomal miR-106a-5p derived from CDDP-resistant cells could promote NPC cell proliferation and suppress apoptosis by targeting ARNT2 and activating AKT phosphorylation, thereby regulating tumorigenesis^[63]. This study revealed that tumor cells could remodel the tumor microenvironment through exosomal miRNAs to acquire a chemoresistant phenotype.

miRNAs affect chemoresistance by regulating cancer stem cells

Cancer stem cells play a key role in tumor development, metastasis, and chemoresistance^[64-66]. MiRNAs are involved in the chemoresistance process of HNC by regulating cancer stemness. Recent studies have shown that miRNA-485-5p can regulate the stemness and chemotherapy resistance of OSCC by targeting keratin 17 (KRT17)^[67]. MiR-21-3p overexpression maintained the stemness of this subpopulation. Lin *et al.* discovered that activation of the miR-371/372/373 cluster could enhance the tumorigenicity and drug tolerance of OSCC cells^[68]. Cai *et al.* found that EBV-encoded miR-BART7-3p enhanced the stemness and chemotherapy resistance of NPC by inhibiting SMAD7 to activate the TGF- β pathway^[69].

miRNAs affect chemoresistance by regulating other signaling pathways

In addition, miRNAs are also involved in chemoresistance of HNC by regulating multiple signaling pathways. Zhang *et al.* found that miR-205-5p could induce EMT and CDDP resistance in NPC by targeting PTEN to activate the PI3K/Akt pathway^[70]. Gu *et al.* discovered that miR-552 could promote the proliferation and metastasis of laryngeal cancer cells by targeting p53^[71]. Sheng *et al.* discovered that miR-21 enhances proliferation, apoptosis inhibition, and CDDP resistance in head and neck squamous cell carcinoma (HNSCC) through the PTEN/PI3K/AKT pathway^[72]. Wu *et al.* found that the miR-577/EIF5A2 axis suppresses the proliferation of CDDP-resistant NPC by blocking the TGF- β signaling pathway and

inhibiting EMT^[73]. In a study by Chen *et al.*, miR-132 was shown to inhibit proliferation and invasion while enhancing CDDP chemosensitivity in OSCC cells via the TGF- β 1/Smad2/3 signaling pathway^[74]. Furthermore, miRNA-296-5p was found to increase the sensitivity of NPC cells to CDDP by targeting and suppressing the STAT3/KLF4 signaling axis^[75]. These results highlight specific miRNAs play a key role in the acquired chemoresistance of HNC by fine-tuning multiple signaling pathways.

miRNAs regulate chemoresistance in HNC by competitively binding to upstream lncRNAs

miRNAs also regulate chemoresistance in HNC by competitively binding to upstream lncRNAs. FOXD1 promotes OSCC chemoresistance by upregulating LPP expression via sponging miR-1252-5p and miR-3148 through CYTOR^[76]. PVT1 promotes cetuximab resistance in head and neck squamous carcinoma by inhibiting miR-124-3p^[77]. KCNQ1OT1 promotes CDDP resistance in NPC through the miR-454/USP47 axis^[78]. NEAT1 promotes CDDP resistance in thyroid cancer(TC) through the miR-9-5p/SPAG9 axis^[79]. CircRNAs also regulate chemoresistance in HNC by sponging miRNAs, such as circCRIM1 promoting docetaxel resistance in NPC through the miR-422a/FOXQ1 axis^[80].

Furthermore, miRNAs can also be involved in the chemoresistance of HNC by affecting drug resistance-related enzymes, membrane transporters, angiogenesis, and other factors. Yuan *et al.* found that LINC-PINT could reverse laryngeal cancer stemness and chemotherapy resistance through the miR-425-5p/PTCH1/SHH axis^[81]. MiR-340 reverses multidrug resistance in NPC by inhibiting P-gp and BCRP^[82]. Docetaxel can induce IL-8 secretion in OSCC cells, thereby promoting angiogenesis.

In recent years, strategies targeting miRNAs to reverse chemoresistance have received widespread attention. Li *et al.* found that miR-101-3p mimics could inhibit the proliferation and CDDP resistance of NPC cells through ZIC5^[47]. Song *et al.* discovered that miR-619-5p inhibitors could enhance the CDDP sensitivity of OSCC cells by activating ATXN3^[83] [Table 2 and Figure 1]. Although miRNAs have shown good antitumor and chemoresistance reversal effects at the animal level, their clinical application still faces many challenges, such as the construction of miRNA delivery systems, in vivo stability, and off-target effects^[84-87]. In addition, the role of circRNAs and lncRNAs as endogenous miRNA sponges in reversing chemoresistance requires further in-depth research.

LONG NON-CODING RNAs IN HNC CELL DRUG RESISTANCE

lncRNAs are extensively expressed and have specific interactions with DNA, RNA, and proteins, allowing them to regulate chromatin function, modulate the assembly and function of membraneless nucleosomes, influence the stability and translation of cytoplasmic mRNA, and participate in the regulation of signaling pathways^[88,89]. Moreover, lncRNAs play a crucial role in regulating processes such as apoptosis, drug efflux, drug metabolism, DNA repair, EMT, autophagy, and ferroptosis. Consequently, they emerge as pivotal mediators of tumor drug resistance^[90-93]. In recent years, there has been a notable increase in the number of lncRNAs associated with cancer development and progression. These lncRNAs can be explored and investigated in meticulously curated databases like Lnc2Cancer3.0^[94] or Cancer lncRNA Census^[95].

CDDP resistance

CDDP stands as a highly extensively employed drug in the treatment of diverse solid tumors^[96]. Its principal mode of action involves disrupting DNA repair mechanisms, leading to DNA damage and consequently triggering apoptosis in cancer cells^[97,98]. Nonetheless, drug resistance poses an inherent challenge in the clinical utilization of CDDP, primarily stemming from three molecular mechanisms: heightened DNA repair, modified cellular accumulation, and augmented drug inactivation^[99]. In 2014, Galluzzi *et al.* Systematically categorized CDDP resistance into four distinct stages: Pre-Target, On-Target, Post-Target,

Table 2. HNC cell drug resistance-related microRNAs

Tumor type	MiRNAs	Cell line	Expression level	Target	Functions	Corresponding drugs	Ref.
NPC	miR-106a-5p	C666-1, CNE2	Down	SOX4	Enhance DDP resistance in NPC cells	CDDP	[37]
NPC	miR-200c	CNE1, CNE2, 5-8F	Up	c-Myc	Promising approach to overcome the oncogenic role of c-Myc in NPC	Cisplatin	[43]
NPC	miR-101-3p	HNE-1/DDP, C666-1/DDP	Up	SOX2	Inhibit cisplatin resistance through miR-101-3p/SOX2/ZIC5 axis	CDDP	[47]
NPC	miR-1278	CNE-1, CNE-2, C666-1, 5-8F, HONE-1	Up	ATG2B	Sensitize NPC cells to DDP and reduce autophagy	CDDP	[52]
NPC	miR-106a-5p	CNE1	Down	ARNT2	Improve recipient cell proliferation, metastasis, and chemoresistance	CDDP	[63]
NPC	miR-205-5p	HNE1/DDP	Down	PTEN	Restrain EMT progression of HNE1/DDP cells	CDDP	[70]
NPC	miR-454	5-8F, SUNE-1	Down	USP47	Suppress NPC cell viability and DDP resistance	CDDP	[78]
NPC	miR-422a	S18, S26	Down	FOXQ1	Promote NPC cell metastasis and EMT	CDDP	[80]
NPC	miR-181a	C666-1, SUNE1	Up	KDM5C	Delay NPC cell progression	/	[82]
LSCC	miR-216a-5p	Tu-686, SNU899, SNU46, Tu-177	Up	ZEB1	Downregulate the cell proliferation and migration and invasive ability	/	[35]
LSCC	miR-107	TU-177, AMC-HN-8	Down	HMGB1	Increase cisplatin resistance	CDDP	[48]
LSCC	miR-425-5p	Hep-2	Up	PTCH1	Regulate laryngeal carcinoma cells through miR 425 5p/PTCH1	CDDP	[81]
OSCC	miR-188-3p	SCC-4, SCC-9, CAL-27, UM1, UM2	Down	ABCB1	Promote cisplatin resistance of Oral Squamous Cell Carcinoma	CDDP	[36]
OSCC	miR-155	UPCI-SCC-131, UPCI-SCC-131R	Down	FOXO3a	Suppress the stem-cell-like property and drug efflux transporter protein expression	CDDP	[49]
OSCC	miR-1249-3p	CAL27, SCC15	Down	ATG9A	Promote autophagy and induce cisplatin resistance	CDDP	[51]
OSCC	miR-34	OECM-1/PTX	Up	p53	Increase DNA damage and apoptosis in a p53-dependend manner	Paclitaxel	[55]
OSCC	miR-494	HSC3, HSC4	Down	NQO1	Induce drug resistance and increase stemness	CDDP	[56]
OSCC	miR-371/372/373	SAS subclones	Up	AKT, β -catenin and Src	Increase both oncogenicity and drug resistance	CDDP	[68]
OSCC	miR-1252-5p	CAL-27 and SCC4	Down	FOXD1	Upregulate LPP expression	CDDP	[76]
OSCC	miR-3148	CAL-27 and SCC4	Down	FOXD1	Upregulate LPP expression	CDDP	[76]
OSCC	miR-619-5p	HN6 and CAL27	Up	ATXN3	Inhibit proliferation and arrest cell cycle progression	CDDP	[84]
TC	miR-9-5p	Nthy-ori 3-1, SW1736, 8505C	Up	SPAG9	Sensitize ATC cells to DDP	CDDP	[79]
TSCC	miR-200c	HSC-3	Down	TUBB3 and PPP2R1B	Increase resistance to DTX, migration, and invasion and decrease apoptosis	Paclitaxel	[38]
TSCC	miR-214	CAL-27 and CP-H203	Up	ULK1	Antagonize antitumor effect	CDDP, paclitaxel	[50]
TSCC	miR-2392	CAL-27 and SCC-9	Up	AGO2	Inhibit apoptosis and cisplatin sensitivity	CDDP	[53]
TSCC	miR-5787	Cal27	Down	MT-CO3	Inhibit the translation of MT-CO3 to regulate cisplatin resistance	CDDP	[54]

HNC: Head and neck cancer; NPC: nasopharyngeal carcinoma; DDP: cisplatin; CDDP: cisplatin; LSCC: laryngeal squamous cell carcinoma; OSCC: oral squamous cell carcinoma; TC: thyroid cancer; TSCC: tongue squamous cell carcinoma; DTX: docetaxel.




Figure 1. Overview of the main molecular mechanisms of microRNAs in HNC drug resistance. HNC: Head and neck cancer.

and Off-Target^[100]. In recent years, the advancement of molecular biology and multi-omics approaches has unveiled additional molecules and mechanisms associated with CDDP resistance in non-NHC. These include epigenetic mechanisms, as well as regulators such as HNRNPU, CHD4, TRPV1, MAFG-AS, and MAST1, among others^[101-106]. Furthermore, a study demonstrated that in non-small cell lung cancer, MEK inhibitors successfully surmounted resistance to combination immunotherapy with CDDP by inducing CXCL10 expression in cancer cells^[107].

lncRNAs mediate drug resistance of NPC

NPC represents an EBV-associated malignancy that is particularly common in southern China, Southeast Asia, and North Africa. The age-standardized incidence rate in these regions ranges from 4 to 25 cases per 100,000 individuals, as reported by GLOBOCAN^[5,108]. Patients diagnosed with NPC generally exhibit a favorable prognosis, with a minimum 5-year survival rate of 80% in cases of locally advanced NPC. However, the 5-year survival rate drops to approximately 20% in recurrent or distant metastatic NPC^[109,110]. The study found that CDDP-resistant NPC exhibits a high expression of lncRNA TINCR, which contributes to the development of chemoresistance via the TINCR/ACLY/PADI1/MAPK/MMP2/9 axis^[111]. Another study identified elevated expression of lncRNA NEAT1 in NPC cells resistant to histone deacetylase inhibitors (HDACis). This upregulation is mediated by the NEAT1/miR-129/Bcl-2 axis, which plays a role in NPC's resistance to HDACis^[112]. Consequently, an epigenetic therapeutic approach involving the use of HDACis in combination with other targeted agents holds promise as a novel strategy for future NPC treatment^[113,114].

lncRNAs mediate drug resistance of HNC

The majority of HNCs originate from the mucosal epithelium of the oral cavity, pharynx, and larynx, making them the most prevalent malignant tumors in the head and neck region^[115]. In head and neck squamous cell carcinoma (HNSCC), STAT3 promotes the transcription of lncRNA HOTAIR and its

interaction with pEZH2-S21, which contributes to resistance against both CDDP and cetuximab^[116]. Another study demonstrated that VN1R5 upregulates lncRNA POP1-1, leading to the promotion of CDDP resistance in HNSCC through its interaction with MCM5^[117]. LSCC represents a highly prevalent subtype of laryngeal cancer, ranking second in incidence among respiratory tumors. Chemotherapy is regarded as the established first-line treatment for patients with advanced LSCC^[118]. A study reported that the activation of the STAT3 signaling pathway by lncRNA FOXD2-AS1 enhances CDDP resistance in LSCC^[119]. OSCC is the prevailing malignant tumor of the oral mucosa, with 57.5% of global oral cancer cases being reported in Asia, particularly in India^[120]. FOXD1 upregulates the expression of lncRNA CYTOR, promoting EMT and conferring CDDP resistance in OSCC^[76]. Additionally, it has been reported that in OSCC, HOXA11-AS upregulates the expression of NQO1 by sponging microRNA-494 while downregulating the expression of NQO2. This regulatory mechanism promotes tumor progression and contributes to drug resistance^[56]. TSCC is characterized by a significant upregulation in the expression of miRNA processing-related lncRNA (MPRL), which disrupts pre-miRNA processing. This disruption promotes mitochondrial fission and enhances CDDP chemosensitivity in TSCC^[121]. TC is a prevalent endocrine malignancy, comprising approximately 1% of all malignant tumors. Moreover, its incidence is consistently rising on a global scale^[122]. Papillary thyroid cancer (PTC) is the most common histological subtype of TC, accounting for 89.1% of cases. Studies have shown that the lncRNA Glycolysis-Associated Regulator of LDHA post-transcriptional modification (GLTC) is significantly upregulated in PTC tissues and correlates with more extensive distant metastasis, increased tumor size, and poorer prognosis. GLTC facilitates the succinylation-dependent activation of LDHA, thus promoting resistance to radioiodine therapy. These findings provide a theoretical foundation for considering the GLTC-LDHA pathway a potential target for therapeutic intervention in PTC^[123].

Several lncRNAs have been found to have close associations with the malignant biology of tumors. In LSCC, HCP5 exhibits high expression levels. The knockdown of HCP5 has been shown to inhibit malignant biological functions through the regulation of miR-216a-5p/ZEB1 signaling pathway^[35]. In OSCC, lnc-p23154 is believed to play a significant role in glut1-mediated glycolysis and promote tumor metastasis by suppressing the transcription of miR-378a-3p^[124]. Additionally, the mitochondria-localized lncRNA growth-arrest-specific 5 (GAS5) has been identified as a tumor suppressor that plays a critical role in maintaining cellular energy homeostasis^[125]. The targeting of epidermal growth factor receptor (EGFR) has proven to be an effective therapeutic strategy for the treatment of SCCs. Notably, studies have indicated that lnc-EGFR-AS1 is involved in mediating EGFR addiction and influencing treatment responses in SCCs^[126] [Table 3 and Figure 2].

Substantial evidence supports the dependence of cellular homeostasis on lncRNAs^[125,127]. However, despite a small fraction of the thousands of expressed lncRNAs potentially having functional roles in cancer cells, the extent of their involvement remains inadequately studied^[128]. Further research is warranted to explore the role of lncRNAs in various aspects concerning tumor chemotherapy, targeted therapy, and immunotherapy, as well as their impact on the tumor microenvironment^[129,130].

circRNAs IN HNC CELL DRUG RESISTANCE

The discovery of single-stranded covalently closed circular RNA dates back to 1976, with subsequent findings revealing their common presence in both viruses and mammals^[131]. circRNAs exhibit a diverse range of functions, encompassing their role as protein scaffolds or miRNA sponges, and their capacity for translation into polypeptides^[132]. The unique structure of circRNAs grants them a longer half-life compared to linear RNA and provides resistance against RNase R degradation^[133]. Consequently, circRNAs have garnered significant attention as reliable diagnostic and prognostic biomarkers in cancer diagnosis, treatment, and prevention^[13,16,134-137].

Table 3. HNC cell drug resistance-related lncRNAs

Tumor type	lncRNA	Cell line	Expression level	Target	Functions	Corresponding drugs	Ref.
NPC	TINCR	NP69 N2Tert CNE-1, HK-1	Up	ACLY	Activate TINCR-ACLY-PADI1-MAPK-MMP2/9 axis	CDDP	[111]
NPC	NEAT1	C666-1, CNE-1, CNE-2	Up	miR-129	Modulate the miR-129/Bcl-2 axis	HDACis	[112]
HNSCC	HOTA1	SCC25, Cal27, UM1	Up	EZH2	Promote the growth of HNSCC cells.	CDDP/cetuximab	[116]
HNSCC	POP1-1	HN4, HN30	Up	MCM5	Facilitate the repair of DNA damage	CDDP	[117]
LSCC	FOXD2-AS1	Hep2, TU-212	Up	STAT3	Promote STAT3 transcriptional activity	CDDP	[119]
OSCC	CYTOR	CAL-27, SCC4	Up	ceRNA	Activate CYTOR/LPP axis	CDDP	[76]
OSCC	HOXA11-AS	HSC3, HSC4	Up	microRNA-494	Facilitate tumor growth	Dicoumarol	[56]
OSCC	EGFR-AS1	primary cell cultures	Up	/	Resistance to EGFR TKIs	TKIs	[126]
PTC	GLTC	BCPAP, TPC-1, KTC-1	Up	LDHA	Resistance to RAI	RAI	[123]

HNC: Head and neck cancer; NPC: nasopharyngeal carcinoma; CDDP: cisplatin; HNSCC: head and neck squamous cell carcinoma; LSCC: laryngeal squamous cell carcinoma; OSCC: oral squamous cell carcinoma; EGFR: epidermal growth factor receptor; TKIs: tyrosine kinase inhibitors; PTC: papillary thyroid cancer; GLTC: glycolysis-associated regulator of LDHA post-transcriptional modification; LDHA: lactate dehydrogenase A; RAI: radioiodine.




Figure 2. Overview of the main molecular mechanisms of lncRNAs in HNC drug resistance. HNC: Head and neck cancer.

circRNAs mediate drug resistance of NPC

Patients with distant metastases of NPC exhibit specific overexpression of circIPO7, and its knockdown enhances sensitivity to CDDP treatment by inhibiting YBX1 nuclear localization^[138]. Furthermore, studies have reported an association between aging and tumor metastasis and chemoresistance. Notably, circWDR37 facilitates the activation of PKR, leading to the initiation of SASP transcription through NF- κ B, thereby promoting metastasis and chemoresistance in NPC^[139]. A separate study demonstrated the role of circCRIM1 as a ceRNA in promoting metastasis and conferring docetaxel chemoresistance in NPC through the upregulation of FOXQ1. Additionally, the study successfully developed a prognostic model based on circCRIM1 expression and TMN staging to assess the risk of NPC metastasis^[80].

circRNAs mediate drug resistance of HNC

CDDP resistance in HNC is strongly linked to autophagy, a process essential for maintaining protein homeostasis, organelle integrity, cellular homeostasis, cell viability, and the degradation/recirculation of various cellular components to the lysosome^[140,141]. In OSCC, significant upregulation of circAP1M2 leads to the induction of autophagy-associated CDDP resistance via the miR-1249-3p/ATG9A axis^[51]. In LSCC, circPARD3 hinders autophagy by serving as a sponge for miR-145-5p, which activates the PRKCI/Akt/mTOR pathway, thereby promoting tumor progression and contributing to CDDP resistance^[142]. TC is characterized by the promotion of autophagy and increased CDDP resistance due to the regulatory effects of circEIF6 on the miR-144-3p/TGF- α axis^[143] [Table 4 and Figure 3]. Interestingly, autophagy exhibits dynamic tumor-suppressive or tumor-promoting roles in diverse contexts and stages of cancer development^[144]. Therefore, further research is needed to better define the specific roles of autophagy in various types and stages of cancer, and to gain a deeper understanding of how tumors rely on autophagy. Furthermore, several studies have demonstrated the reciprocal regulation between circRNAs and N⁶-methyladenosine (m⁶A) modifications in HNC, highlighting their potential impact on cancer progression and treatment response^[145-148]. Additionally, there have been reports indicating the upregulation of circCUX1 expression in radiotherapy-resistant hypopharyngeal SCC patients, and the knockdown of circCUX1 has been shown to enhance the sensitivity of hypopharyngeal cancer cells to radiotherapy^[149].

FUTURE PERSPECTIVES

Based on our persistent dedication to this field and thorough literature research, we firmly assert that the following areas of research exhibit dynamism and merit sustained attention.

Ferroptosis

Ferroptosis, also known as iron-induced apoptosis, is a type of intracellular cell death that relies on iron and is distinguished by the excessive accumulation of reactive oxygen species (ROS) and lipid peroxidation. It is distinct from apoptosis, necrosis, and autophagy^[150,151]. Numerous studies have reported that tumor cells can enhance their defense mechanisms against oxidative stress by inhibiting ferroptosis, ultimately promoting their survival and drug resistance^[152,153]. Studies have demonstrated that in NPC, infection with EBV leads to the upregulation of GPX4 expression, resulting in the inhibition of ferroptosis. Consequently, the elevated levels of GPX4 contribute to the progression of NPC and its resistance to chemotherapy through the activation of the TAK1-JNK and IKK/NF- κ B signaling pathways. Moreover, clearance of the EBV genome has been shown to enhance the sensitivity of NPC cells to ferroptosis^[154]. Artesunate exhibits selective cytotoxicity against HNC cells. In specific cases of CDDP-resistant HNC, inhibiting the Nrf2-ARE pathway enhances the sensitivity of these cells to artesunate and reverses their resistance to ferroptosis^[155,156]. Additional studies have further demonstrated that inhibiting GLRX5 renders CDDP-resistant HNC cells more susceptible to ferroptosis^[157]. Furthermore, the upregulation of HMGA1 in ESCC serves as a crucial factor responsible for CDDP resistance by suppressing ferroptosis. This effect is achieved through HMGA1's role in maintaining intracellular redox homeostasis via its assistance to ATF4 in activating

Table 4. HNC cell drug resistance-related circRNAs

Tumor type	circRNA	Cell line	Expression level	Target	Functions	Corresponding drugs	Ref.
NPC	IPO7	CNE2, HNE1, HONE1, SUNE1, HK1, C666-1	Up	YBX1	Promote cell migration, invasion, and cisplatin resistance	CDDP	[138]
NPC	WDR37	S18, S26	Up	PKR	CCND1	CDDP, gemcitabine	[139]
NPC	CRIM1	S18, S26	Up	miR-422a	Promote metastasis and EMT	Docetaxel	[80]
OSCC	AP1M2	CAL27/CDDP	Up	miR-1249-3p	modulate miR-1249-3p-ATG9A axis	CDDP	[51]
LSCC	PARD3	Tu 177, HOK, FD-LSC-1	Up	miR-145-5p w	Activate the Akt-mTOR axis	CDDP	[142]
ATC	EIF6	TPC1, BHT101	Up	miR-144-3p	Regulate miR144-3p/TGF- α axis	CDDP	[143]

HNC: Head and neck cancer; NPC: nasopharyngeal carcinoma; CDDP: cisplatin; CCND1: stimulate cyclin D1; EMT: epithelial-mesenchymal transition; OSCC: oral squamous cell carcinoma; LSCC: laryngeal squamous cell carcinoma; ATC: anaplastic thyroid carcinoma.




Figure 3. Overview of the main molecular mechanisms of circRNAs in HNC drug resistance. HNC: Head and neck cancer.

SLC7A11 transcription. Inhibition of HMGA1 has been shown to enhance the sensitivity of ESCC to ferroptosis^[158]. Multiple studies have now established that regulating ferroptosis can impact the effectiveness of cancer treatment and potentially overcome resistance to chemotherapy, targeted therapy, and immunotherapy^[159-164]. Several non-coding RNAs, including miR-324-3p, miR-375, miR-144-3p, miR-27a-3p, miR-3173-5p, circRNA-101093, and lncRNA-PMAN, have been implicated in the regulation of tumor ferroptosis^[165-168]. Currently, there are no relevant studies investigating the role of non-coding RNAs in tumor drug resistance through the regulation of ferroptosis specifically in HNC.

Nanodrugs

CDDP, carboplatin, and oxaliplatin are commonly employed in tumor therapy. Nevertheless, their clinical utility is severely restricted due to the side effects associated with platinum drugs, including poor selectivity, high systemic toxicity, and drug resistance^[169]. A novel CDDP nanocarrier system with dual targeting properties has been developed. This system selectively attaches to the A54 receptor, which is highly expressed on the cell surface of hepatocellular carcinoma (HCC) cells. Additionally, it utilizes the drug-resistance gene NOR1 shRNA as a piggyback mechanism to enable precise tumor-targeted therapy and overcome drug resistance^[170]. Numerous studies have also highlighted the effectiveness of platinum nanocarriers in reducing systemic toxicity and overcoming drug resistance^[171-174]. Currently, there is considerable research focus on strategies involving nanomedicines for regulating ferroptosis, as well as nanomedicines utilizing non-coding RNAs as carriers to target tumors^[85,175-179].

Exosomes

Exosomes are nanovesicles derived from cells, ranging in size from 30 to 150 nm. They are released when multivesicular bodies fuse with the cell surface and play a crucial role in intercellular communication by transporting nucleic acids, proteins, and lipids. Furthermore, exosomes can activate signaling pathways in target cells^[180,181]. In recent years, there has been significant interest in the role of exosomes carrying non-coding RNA in tumor drug resistance^[93,163,182-185]. Several studies have reported that intervention with RNA carried by exosomes can potentially overcome tumor chemoresistance^[61,186,187]. For instance, CAFs emerge as crucial regulators of CDDP resistance in HNC. They achieve this by transporting functional miR-196a from CAFs to tumor cells through exosomes^[62]. Although our understanding of the role of RNAs carried within exosomes in the mechanisms of drug resistance in HNC is still in the early stages, exosomal circRNAs have emerged as innovative genetic information carriers. These molecules enable communication between tumor cells and cells in the microenvironment, thereby regulating critical aspects of cancer progression. As a result, they contribute significantly to chemotherapeutic drug resistance across different types of cancer. Based on this, synthetic exosomal circRNA holds the potential to introduce new avenues for cancer therapy^[188,189].

Proteolysis-targeting chimeras

Proteolysis-targeting chimeras (PROTACs) are emerging as promising therapeutic modalities for the degradation of disease-causing proteins. They are composed of a ligand that binds to a protein of interest (POI) and another ligand that recruits the E3 ubiquitin ligase. This recruitment induces chemical proximity between the POI and the E3 ligase, resulting in ubiquitination and subsequent degradation of the POI through the ubiquitin-proteasome system^[190-193]. PROTACs have been developed to target several proteins, including PD-L1, BTK, STAT3, EGFR, MEK1/2, VEGFR2, FLT-3, and SHP2^[194]. These compounds offer advantages such as high enzymatic reaction efficiency, overall degradation of target proteins, and the ability to target small-molecule non-druggable proteins. Several studies have reported the successful inhibition of HNSCC growth and metastasis using PROTAC technology^[195,196]. This finding highlights the importance of continued attention to PROTAC and related technologies due to their ability to specifically degrade oncogenic proteins or RNAs^[194,197,198].

CONCLUSION

In summary, chemotherapy, targeted therapy, and immunotherapy are crucial components of integrated tumor therapy. However, drug resistance significantly limits the effectiveness of clinical drug therapy for HNC. Non-coding RNA dysregulation plays a critical role in the known mechanisms of drug resistance in HNC. It is essential to continually focus on new technologies and theories to address the challenges in HNC drug resistance research. This includes the construction or utilization of clinical databases similar to UK biobank^[199], and integrating clinical information with multi-omics sequencing data. These approaches aim to establish a robust foundation for further research on integrative oncology treatment^[200]. Additionally,

leveraging ChatGPT automated development and analysis (ADA) technology to build machine learning models based on real clinical trial data^[201] can facilitate efficient research on cancer precision treatment, prognosis, and drug resistance markers.

DECLARATIONS

Acknowledgments

We appreciate and acknowledge the support from Dr. Rensen Ran and Lei Chen for their guidance and assistance in drawing.

Authors' contributions

Conceived the contents and structure: Yang S, Deng F, Yang X, Li A, Xia W, Gao C, Lei S

Wrote the original manuscript: Zhang Y, Peng Y, Lin B

Revised and improved the manuscript: Liao W, Zeng Q

All authors have read and agreed to the published version of the manuscript.

Financial support and sponsorship

Project of Administration of Traditional Chinese Medicine of Guangdong Province of China (No. 20231070).

Conflicts of interest

All authors declared that there are no conflicts of interest.

Ethical approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Copyright

© The Author(s) 2024.

REFERENCES

1. Li Q, Tie Y, Alu A, Ma X, Shi H. Targeted therapy for head and neck cancer: signaling pathways and clinical studies. *Signal Transduct Target Ther* 2023;8:31. [DOI](#) [PubMed](#) [PMC](#)
2. Pfister DG, Spencer S, Adelstein D, et al. Head and neck cancers, version 2.2020, NCCN clinical practice guidelines in oncology. *J Natl Compr Canc Netw* 2020;18:873-98. [DOI](#) [PubMed](#)
3. Mody MD, Rocco JW, Yom SS, Haddad RI, Saba NF. Head and neck cancer. *Lancet* 2021;398:2289-99. [DOI](#) [PubMed](#)
4. Chow LQM. Head and neck cancer. *N Engl J Med* 2020;382:60-72. [DOI](#) [PubMed](#)
5. Wong KCW, Hui EP, Lo KW, et al. Nasopharyngeal carcinoma: an evolving paradigm. *Nat Rev Clin Oncol* 2021;18:679-95. [DOI](#) [PubMed](#)
6. Leemans CR, Braakhuis BJ, Brakenhoff RH. The molecular biology of head and neck cancer. *Nat Rev Cancer* 2011;11:9-22. [DOI](#) [PubMed](#)
7. Dai F, Dai L, Zheng X, et al. Non-coding RNAs in drug resistance of head and neck cancers: a review. *Biomed Pharmacother* 2020;127:110231. [DOI](#) [PubMed](#)
8. Romano G, Veneziano D, Acunzo M, Croce CM. Small non-coding RNA and cancer. *Carcinogenesis* 2017;38:485-91. [DOI](#) [PubMed](#) [PMC](#)
9. Nemeth K, Bayraktar R, Ferracin M, Calin GA. Non-coding RNAs in disease: from mechanisms to therapeutics. *Nat Rev Genet* 2024;25:211-32. [DOI](#) [PubMed](#)
10. Patop IL, Wüst S, Kadener S. Past, present, and future of circRNAs. *EMBO J* 2019;38:e100836. [DOI](#) [PubMed](#) [PMC](#)
11. Kabekkodu SP, Shukla V, Varghese VK, D'Souza J, Chakrabarty S, Satyamoorthy K. Clustered miRNAs and their role in biological functions and diseases. *Biol Rev Camb Philos Soc* 2018;93:1955-86. [DOI](#) [PubMed](#)
12. Ransohoff JD, Wei Y, Khavari PA. The functions and unique features of long intergenic non-coding RNA. *Nat Rev Mol Cell Biol*

- 2018;19:143-57. DOI PubMed PMC
13. Kristensen LS, Jakobsen T, Hager H, Kjems J. The emerging roles of circRNAs in cancer and oncology. *Nat Rev Clin Oncol* 2022;19:188-206. DOI PubMed
 14. Wen SY, Qadir J, Yang BB. Circular RNA translation: novel protein isoforms and clinical significance. *Trends Mol Med* 2022;28:405-20. DOI PubMed
 15. Williams M, Cheng YY, Phimmachanh M, Winata P, van Zandwijk N, Reid G. Tumour suppressor microRNAs contribute to drug resistance in malignant pleural mesothelioma by targeting anti-apoptotic pathways. *Cancer Drug Resist* 2019;2:1193-206. DOI PubMed PMC
 16. Chen B, Dragomir MP, Yang C, Li Q, Horst D, Calin GA. Targeting non-coding RNAs to overcome cancer therapy resistance. *Signal Transduct Target Ther* 2022;7:121. DOI PubMed PMC
 17. Zhang X, Xie K, Zhou H, et al. Role of non-coding RNAs and RNA modifiers in cancer therapy resistance. *Mol Cancer* 2020;19:47. DOI PubMed PMC
 18. Mattick JS, Amaral PP, Carninci P, et al. Long non-coding RNAs: definitions, functions, challenges and recommendations. *Nat Rev Mol Cell Biol* 2023;24:430-47. DOI PubMed PMC
 19. Huarte M. The emerging role of lncRNAs in cancer. *Nat Med* 2015;21:1253-61. DOI PubMed
 20. Yuan JH, Yang F, Wang F, et al. A long noncoding RNA activated by TGF- β promotes the invasion-metastasis cascade in hepatocellular carcinoma. *Cancer Cell* 2014;25:666-81. DOI PubMed
 21. Winkle M, El-Daly SM, Fabbri M, Calin GA. Noncoding RNA therapeutics - challenges and potential solutions. *Nat Rev Drug Discov* 2021;20:629-51. DOI PubMed PMC
 22. Slack FJ, Chinnaiyan AM. The role of non-coding RNAs in oncology. *Cell* 2019;179:1033-55. DOI PubMed PMC
 23. Calin GA, Croce CM. MicroRNA signatures in human cancers. *Nat Rev Cancer* 2006;6:857-66. DOI PubMed
 24. Cavaliere AF, Perelli F, Zaami S, et al. Towards personalized medicine: non-coding RNAs and endometrial cancer. *Healthcare* 2021;9:965. DOI PubMed PMC
 25. Piergentili R, Basile G, Nocella C, et al. Using ncRNAs as tools in cancer diagnosis and treatment-the way towards personalized medicine to improve patients' health. *Int J Mol Sci* 2022;23:9353. DOI PubMed PMC
 26. Rupaimoole R, Slack FJ. MicroRNA therapeutics: towards a new era for the management of cancer and other diseases. *Nat Rev Drug Discov* 2017;16:203-22. DOI PubMed
 27. Naorem LD, Prakash VS, Muthaiyan M, Venkatesan A. Comprehensive analysis of dysregulated lncRNAs and their competing endogenous RNA network in triple-negative breast cancer. *Int J Biol Macromol* 2020;145:429-36. DOI PubMed
 28. Mahato RK, Bhattacharya S, Khullar N, et al. Targeting long non-coding RNAs in cancer therapy using CRISPR-Cas9 technology: a novel paradigm for precision oncology. *J Biotechnol* 2024;379:98-119. DOI PubMed
 29. Kilikevicius A, Meister G, Corey DR. Reexamining assumptions about miRNA-guided gene silencing. *Nucleic Acids Res* 2022;50:617-34. DOI PubMed PMC
 30. Shang R, Lee S, Senavirathne G, Lai EC. microRNAs in action: biogenesis, function and regulation. *Nat Rev Genet* 2023;24:816-33. DOI PubMed PMC
 31. Agbu P, Carthew RW. MicroRNA-mediated regulation of glucose and lipid metabolism. *Nat Rev Mol Cell Biol* 2021;22:425-38. DOI PubMed PMC
 32. He B, Zhao Z, Cai Q, et al. miRNA-based biomarkers, therapies, and resistance in Cancer. *Int J Biol Sci* 2020;16:2628-47. DOI PubMed PMC
 33. Pan G, Liu Y, Shang L, Zhou F, Yang S. EMT-associated microRNAs and their roles in cancer stemness and drug resistance. *Cancer Commun* 2021;41:199-217. DOI PubMed PMC
 34. Meng X, Lou QY, Yang WY, et al. The role of non-coding RNAs in drug resistance of oral squamous cell carcinoma and therapeutic potential. *Cancer Commun* 2021;41:981-1006. DOI PubMed PMC
 35. Zhang S, Huangfu H, Zhao Q, Li Y, Wu L. Downregulation of long noncoding RNA HCP5/miR-216a-5p/ZEB1 axis inhibits the malignant biological function of laryngeal squamous cell carcinoma cells. *Front Immunol* 2022;13:1022677. DOI PubMed PMC
 36. Gao F, Han J, Wang Y, Jia L, Luo W, Zeng Y. Circ_0109291 promotes cisplatin resistance of oral squamous cell carcinoma by sponging miR-188-3p to increase ABCB1 expression. *Cancer Biother Radiopharm* 2022;37:233-45. DOI PubMed
 37. Cao W, Sun Y, Liu L, et al. HOTAIR mediates cisplatin resistance in nasopharyngeal carcinoma by regulating miR-106a-5p/SOX4 axis. *Bioengineered* 2022;13:6567-78. DOI PubMed PMC
 38. Cui J, Wang H, Zhang X, Sun X, Zhang J, Ma J. Exosomal miR-200c suppresses chemoresistance of docetaxel in tongue squamous cell carcinoma by suppressing TUBB3 and PPP2R1B. *Aging* 2020;12:6756-73. DOI PubMed PMC
 39. Proença C, Freitas M, Ribeiro D, Rufino AT, Fernandes E, Ferreira de Oliveira JMP. The role of flavonoids in the regulation of epithelial-mesenchymal transition in cancer: a review on targeting signaling pathways and metastasis. *Med Res Rev* 2023;43:1878-945. DOI PubMed
 40. Garinet S, Didelot A, Denize T, et al. Clinical assessment of the miR-34, miR-200, ZEB1 and SNAIL EMT regulation hub underlines the differential prognostic value of EMT miRs to drive mesenchymal transition and prognosis in resected NSCLC. *Br J Cancer* 2021;125:1544-51. DOI PubMed PMC
 41. Chien CS, Wang ML, Chu PY, et al. Lin28B/Let-7 regulates expression of Oct4 and Sox2 and reprograms oral squamous cell carcinoma cells to a stem-like state. *Cancer Res* 2015;75:2553-65. DOI PubMed

42. Manikandan M, Deva Magendhra Rao AK, Arunkumar G, et al. Oral squamous cell carcinoma: microRNA expression profiling and integrative analyses for elucidation of tumorigenesis mechanism. *Mol Cancer* 2016;15:28. DOI PubMed PMC
43. Yang J, Wu SP, Wang WJ, et al. A novel miR-200c/c-myc negative regulatory feedback loop is essential to the EMT process, CSC biology and drug sensitivity in nasopharyngeal cancer. *Exp Cell Res* 2020;391:111817. DOI PubMed
44. Deng X, Liu Z, Liu X, et al. miR-296-3p negatively regulated by nicotine stimulates cytoplasmic translocation of c-Myc via MK2 to suppress chemotherapy resistance. *Mol Ther* 2018;26:1066-81. DOI PubMed PMC
45. Huni KC, Cheung J, Sullivan M, Robison WT, Howard KM, Kingsley K. Chemotherapeutic drug resistance associated with differential miRNA expression of miR-375 and miR-27 among oral cancer cell lines. *Int J Mol Sci* 2023;24:1244. DOI PubMed PMC
46. Shaw P, Raymond G, Senthilnathan R, et al. Clinical theragnostic relationship between chemotherapeutic resistance, and sensitivity and miRNA expressions in head and neck cancers: a systematic review and meta-analysis protocol. *Genes* 2021;12:2029. DOI PubMed PMC
47. Li T, Zhang G, Li W, et al. MicroRNA-101-3p inhibits nasopharyngeal carcinoma cell proliferation and cisplatin resistance through ZIC5 down-regulation by targeting SOX2. *Biol Chem* 2023;404:961-75. DOI PubMed
48. Chen L, Xu Z, Zhao J, et al. H19/miR-107/HMGB1 axis sensitizes laryngeal squamous cell carcinoma to cisplatin by suppressing autophagy in vitro and in vivo. *Cell Biol Int* 2021;45:674-85. DOI PubMed
49. Sayyed AA, Gondaliya P, Mali M, et al. MiR-155 inhibitor-laden exosomes reverse resistance to cisplatin in a 3D tumor spheroid and xenograft model of oral cancer. *Mol Pharm* 2021;18:3010-25. DOI PubMed
50. Yang Y, Sun X, Li M, Li L, Wang S, Zhu Y. miR-214 modulates the growth and migration of oral cancer before and after chemotherapy through mediating ULK1. *J Immunol Res* 2022;2022:4589182. DOI PubMed PMC
51. Wenhao R, Yali C, Shaoming L, Jingjing Z, Ling G, Keqian Z. circAP1M2 activates ATG9A-associated autophagy by inhibiting miR-1249-3p to promote cisplatin resistance in oral squamous cell carcinoma. *J Cell Physiol* 2023;238:2612-24. DOI PubMed
52. Zhao Y, Wang P, Wu Q. miR-1278 sensitizes nasopharyngeal carcinoma cells to cisplatin and suppresses autophagy via targeting ATG2B. *Mol Cell Probes* 2020;53:101597. DOI PubMed
53. Fan S, Tian T, Chen W, et al. Mitochondrial miRNA determines chemoresistance by reprogramming metabolism and regulating mitochondrial transcription. *Cancer Res* 2019;79:1069-84. DOI PubMed PMC
54. Chen W, Wang P, Lu Y, et al. Decreased expression of mitochondrial miR-5787 contributes to chemoresistance by reprogramming glucose metabolism and inhibiting MT-CO3 translation. *Theranostics* 2019;9:5739-54. DOI PubMed PMC
55. Mortezagholi B, Nasiri K, Movahed E, et al. MiR-34 by targeting p53 induces apoptosis and DNA damage in paclitaxel-resistant human oral squamous carcinoma cells. *Chem Biol Drug Des* 2023;102:285-91. DOI PubMed
56. Nakashima C, Fujiwara-Tani R, Mori S, et al. An axis between the long non-coding RNA HOXA11-AS and NQOs enhances metastatic ability in oral squamous cell carcinoma. *Int J Mol Sci* 2022;23:10704. DOI PubMed PMC
57. Kang SH, Oh SY, Lee KY, et al. Differential effect of cancer-associated fibroblast-derived extracellular vesicles on cisplatin resistance in oral squamous cell carcinoma via miR-876-3p. *Theranostics* 2024;14:460-79. DOI PubMed PMC
58. Li P, Yang Y, Liu H, et al. MiR-194 functions as a tumor suppressor in laryngeal squamous cell carcinoma by targeting Wee1. *J Hematol Oncol* 2017;10:32. DOI PubMed PMC
59. Luo X, Li Y, Hua Z, et al. Exosomes-mediated tumor metastasis through reshaping tumor microenvironment and distant niche. *J Control Release* 2023;353:327-36. DOI PubMed
60. Dai J, Su Y, Zhong S, et al. Exosomes: key players in cancer and potential therapeutic strategy. *Signal Transduct Target Ther* 2020;5:145. DOI PubMed PMC
61. Huang C, Zhou Y, Feng X, Wang J, Li Y, Yao X. Delivery of engineered primary tumor-derived exosomes effectively suppressed the colorectal cancer chemoresistance and liver metastasis. *ACS Nano* 2023;17:10313-26. DOI PubMed
62. Qin X, Guo H, Wang X, et al. Exosomal miR-196a derived from cancer-associated fibroblasts confers cisplatin resistance in head and neck cancer through targeting CDKN1B and ING5. *Genome Biol* 2019;20:12. DOI PubMed PMC
63. Li J, Hu C, Chao H, et al. Exosomal transfer of miR-106a-5p contributes to cisplatin resistance and tumorigenesis in nasopharyngeal carcinoma. *J Cell Mol Med* 2021;25:9183-98. DOI PubMed PMC
64. Babaei G, Aziz SG, Jaghi NZZ. EMT, cancer stem cells and autophagy; The three main axes of metastasis. *Biomed Pharmacother* 2021;133:110909. DOI PubMed
65. Rao X, Zhang C, Luo H, et al. Targeting gastric cancer stem cells to enhance treatment response. *Cells* 2022;11:2828. DOI PubMed PMC
66. Lu H, Ju DD, Yang GD, et al. Targeting cancer stem cell signature gene SMOC-2 Overcomes chemoresistance and inhibits cell proliferation of endometrial carcinoma. *EBioMedicine* 2019;40:276-89. DOI PubMed PMC
67. Jang TH, Huang WC, Tung SL, et al. MicroRNA-485-5p targets keratin 17 to regulate oral cancer stemness and chemoresistance via the integrin/FAK/Src/ERK/β-catenin pathway. *J Biomed Sci* 2022;29:42. DOI PubMed PMC
68. Lin SC, Wu HL, Yeh LY, Yang CC, Kao SY, Chang KW. Activation of the miR-371/372/373 miRNA cluster enhances oncogenicity and drug resistance in oral carcinoma cells. *Int J Mol Sci* 2020;21:9442. DOI PubMed PMC
69. Cai L, Long Y, Chong T, et al. EBV-miR-BART7-3p imposes stemness in nasopharyngeal carcinoma cells by suppressing SMAD7. *Front Genet* 2019;10:939. DOI PubMed PMC
70. Zhang P, Lu X, Shi Z, et al. miR-205-5p regulates epithelial-mesenchymal transition by targeting PTEN via PI3K/AKT signaling

- pathway in cisplatin-resistant nasopharyngeal carcinoma cells. *Gene* 2019;710:103-13. DOI PubMed
71. Gu J, Han T, Sun L, Yan AH, Jiang XJ. miR-552 promotes laryngocarcinoma cells proliferation and metastasis by targeting p53 pathway. *Cell Cycle* 2020;19:1012-21. DOI PubMed PMC
 72. Sheng S, Su W, Mao D, et al. MicroRNA-21 induces cisplatin resistance in head and neck squamous cell carcinoma. *PLoS One* 2022;17:e0267017. DOI PubMed PMC
 73. Wu R, Zhong Q, Liu H, Liu S. MicroRNA-577/EIF5A2 axis suppressed the proliferation of DDP-resistant nasopharyngeal carcinoma cells by blocking TGF- β signaling pathway. *Chem Biol Drug Des* 2023;102:815-27. DOI PubMed
 74. Chen L, Zhu Q, Lu L, Liu Y. MiR-132 inhibits migration and invasion and increases chemosensitivity of cisplatin-resistant oral squamous cell carcinoma cells via targeting TGF- β 1. *Bioengineered* 2020;11:91-102. DOI PubMed PMC
 75. Luo HQ, Wang Y, Ren J, et al. MiRNA-296-5p promotes the sensitivity of nasopharyngeal carcinoma cells to cisplatin via targeted inhibition of STAT3/KLF4 signaling axis. *Sci Rep* 2024;14:6681. DOI PubMed PMC
 76. Chen S, Yang M, Wang C, et al. Forkhead box D1 promotes EMT and chemoresistance by upregulating lncRNA CYTOR in oral squamous cell carcinoma. *Cancer Lett* 2021;503:43-53. DOI PubMed
 77. Yang S, Yuan ZJ, Zhu YH, Chen X, Wang W. lncRNA PVT1 promotes cetuximab resistance of head and neck squamous cell carcinoma cells by inhibiting miR-124-3p. *Head Neck* 2021;43:2712-23. DOI PubMed
 78. Yuan F, Lou Z, Zhou Z, Yan X. [Retracted] Long non-coding RNA KCNQ1OT1 promotes nasopharyngeal carcinoma cell cisplatin resistance via the miR-454/USP47 axis. *Int J Mol Med* 2024;53:31. DOI PubMed PMC
 79. Yan P, Su Z, Zhang Z, Gao T. lncRNA NEAT1 enhances the resistance of anaplastic thyroid carcinoma cells to cisplatin by sponging miR-95p and regulating SPAG9 expression. *Int J Oncol* 2019;55:988-1002. DOI PubMed PMC
 80. Hong X, Liu N, Liang Y, et al. Circular RNA CRIM1 functions as a ceRNA to promote nasopharyngeal carcinoma metastasis and docetaxel chemoresistance through upregulating FOXQ1. *Mol Cancer* 2020;19:33. DOI PubMed PMC
 81. Yuan Z, Xiu C, Liu D, et al. Long noncoding RNA LINC-PINT regulates laryngeal carcinoma cell stemness and chemoresistance through miR-425-5p/PTCH1/SHH axis. *J Cell Physiol* 2019;234:23111-22. DOI PubMed
 82. Liu J, Zhu M, Tang Q. Human umbilical cord mesenchymal stem cells-derived exosomal microRNA-181a retards nasopharyngeal carcinoma development by mediating KDM5C. *J Cancer Res Clin Oncol* 2021;147:2867-77. DOI PubMed
 83. Song A, Wu Y, Chu W, et al. Involvement of miR-619-5p in resistance to cisplatin by regulating ATXN3 in oral squamous cell carcinoma. *Int J Biol Sci* 2021;17:430-47. DOI PubMed PMC
 84. Diener C, Keller A, Meese E. Emerging concepts of miRNA therapeutics: from cells to clinic. *Trends Genet* 2022;38:613-26. DOI PubMed
 85. Kara G, Calin GA, Ozpolat B. RNAi-based therapeutics and tumor targeted delivery in cancer. *Adv Drug Deliv Rev* 2022;182:114113. DOI PubMed
 86. Jiménez-Morales JM, Hernández-Cuenca YE, Reyes-Abrahantes A, et al. MicroRNA delivery systems in glioma therapy and perspectives: a systematic review. *J Control Release* 2022;349:712-30. DOI PubMed
 87. Han J, LaVigne CA, Jones BT, Zhang H, Gillett F, Mendell JT. A ubiquitin ligase mediates target-directed microRNA decay independently of tailing and trimming. *Science* 2020;370:eabc9546. DOI PubMed PMC
 88. Statello L, Guo CJ, Chen LL, Huarte M. Gene regulation by long non-coding RNAs and its biological functions. *Nat Rev Mol Cell Biol* 2021;22:96-118. DOI PubMed PMC
 89. Nojima T, Proudfoot NJ. Mechanisms of lncRNA biogenesis as revealed by nascent transcriptomics. *Nat Rev Mol Cell Biol* 2022;23:389-406. DOI PubMed
 90. Zhang Y, Luo M, Cui X, O'Connell D, Yang Y. Long noncoding RNA NEAT1 promotes ferroptosis by modulating the miR-362-3p/MIOX axis as a ceRNA. *Cell Death Differ* 2022;29:1850-63. DOI PubMed PMC
 91. Singh D, Assaraf YG, Gacche RN. Long non-coding RNA mediated drug resistance in breast cancer. *Drug Resist Updat* 2022;63:100851. DOI PubMed
 92. Eptaminitaki GC, Stellas D, Bonavida B, Baritaki S. Long non-coding RNAs (lncRNAs) signaling in cancer chemoresistance: From prediction to druggability. *Drug Resist Updat* 2022;65:100866. DOI PubMed
 93. Yu Z, Tang H, Chen S, et al. Exosomal LOC85009 inhibits docetaxel resistance in lung adenocarcinoma through regulating ATG5-induced autophagy. *Drug Resist Updat* 2023;67:100915. DOI PubMed
 94. Gao Y, Shang S, Guo S, et al. lnc2Cancer 3.0: an updated resource for experimentally supported lncRNA/circRNA cancer associations and web tools based on RNA-seq and scRNA-seq data. *Nucleic Acids Res* 2021;49:D1251-8. DOI PubMed PMC
 95. Carlevaro-Fita J, Lanzós A, Feuerbach L, et al; PCAWG Drivers and Functional Interpretation Group; PCAWG Consortium. Cancer lncRNA Census reveals evidence for deep functional conservation of long noncoding RNAs in tumorigenesis. *Commun Biol* 2020;3:56. DOI PubMed PMC
 96. Ghosh S. Cisplatin: The first metal based anticancer drug. *Bioorg Chem* 2019;88:102925. DOI PubMed
 97. Dasari S, Tchounwou PB. Cisplatin in cancer therapy: molecular mechanisms of action. *Eur J Pharmacol* 2014;740:364-78. DOI PubMed PMC
 98. Perego P. Tackling cisplatin resistance in ovarian cancer: what can we do? *Cancer Drug Resist* 2021;4:755-7. DOI PubMed PMC
 99. Amable L. Cisplatin resistance and opportunities for precision medicine. *Pharmacol Res* 2016;106:27-36. DOI PubMed
 100. Galluzzi L, Vitale I, Michels J, et al. Systems biology of cisplatin resistance: past, present and future. *Cell Death Dis* 2014;5:e1257. DOI PubMed PMC

101. Xie R, Cheng L, Huang M, et al. NAT10 drives cisplatin chemoresistance by enhancing ac4C-associated DNA repair in bladder cancer. *Cancer Res* 2023;83:1666-83. DOI PubMed
102. Shi ZD, Hao L, Han XX, et al. Targeting HNRNPU to overcome cisplatin resistance in bladder cancer. *Mol Cancer* 2022;21:37. DOI PubMed PMC
103. Wu J, Zhou Z, Li J, et al. CHD4 promotes acquired chemoresistance and tumor progression by activating the MEK/ERK axis. *Drug Resist Updat* 2023;66:100913. DOI PubMed
104. Oh SJ, Lim JY, Son MK, et al. TRPV1 inhibition overcomes cisplatin resistance by blocking autophagy-mediated hyperactivation of EGFR signaling pathway. *Nat Commun* 2023;14:2691. DOI PubMed PMC
105. Xiang L, Zeng Q, Liu J, et al. MAFG-AS1/MAFG positive feedback loop contributes to cisplatin resistance in bladder urothelial carcinoma through antagonistic ferroptosis. *Sci Bull* 2021;66:1773-88. DOI PubMed
106. Tyagi A, Kaushal K, Chandrasekaran AP, et al. CRISPR/Cas9-based genome-wide screening for deubiquitinase subfamily identifies USP1 regulating MAST1-driven cisplatin-resistance in cancer cells. *Theranostics* 2022;12:5949-70. DOI PubMed PMC
107. Limagne E, Nuttin L, Thibaudin M, et al. MEK inhibition overcomes chemoimmunotherapy resistance by inducing CXCL10 in cancer cells. *Cancer Cell* 2022;40:136-52.e12. DOI PubMed
108. Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 2018;68:394-424. DOI PubMed
109. Hong S, Zhang Y, Yu G, et al. Gemcitabine plus cisplatin versus fluorouracil plus cisplatin as first-line therapy for recurrent or metastatic nasopharyngeal carcinoma: final overall survival analysis of GEM20110714 phase III study. *J Clin Oncol* 2021;39:3273-82. DOI PubMed PMC
110. Tang QN, Liu LT, Qi B, et al. Effect of concurrent chemoradiotherapy with nedaplatin vs cisplatin on the long-term outcomes of survival and toxic effects among patients with stage II to IVB nasopharyngeal carcinoma: a 5-year follow-up secondary analysis of a randomized clinical trial. *JAMA Netw Open* 2021;4:e2138470. DOI PubMed PMC
111. Zheng ZQ, Li ZX, Guan JL, et al. Long noncoding RNA TINCR-mediated regulation of acetyl-CoA metabolism promotes nasopharyngeal carcinoma progression and chemoresistance. *Cancer Res* 2020;80:5174-88. DOI PubMed
112. Xue F, Cheng Y, Xu L, et al. LncRNA NEAT1/miR-129/Bcl-2 signaling axis contributes to HDAC inhibitor tolerance in nasopharyngeal cancer. *Aging* 2020;12:14174-88. DOI PubMed PMC
113. Yang J, Xu J, Wang W, Zhang B, Yu X, Shi S. Epigenetic regulation in the tumor microenvironment: molecular mechanisms and therapeutic targets. *Signal Transduct Target Ther* 2023;8:210. DOI PubMed PMC
114. Tang J, Wang X, Xiao D, Liu S, Tao Y. The chromatin-associated RNAs in gene regulation and cancer. *Mol Cancer* 2023;22:27. DOI PubMed PMC
115. Johnson DE, Burtress B, Leemans CR, Lui VWY, Bauman JE, Grandis JR. Head and neck squamous cell carcinoma. *Nat Rev Dis Primers* 2020;6:92. DOI PubMed PMC
116. Sun S, Wu Y, Guo W, et al. STAT3/HOTAIR signaling axis regulates HNSCC growth in an EZH2-dependent manner. *Clin Cancer Res* 2018;24:2665-77. DOI PubMed
117. Jiang Y, Guo H, Tong T, et al. lncRNA lnc-POP1-1 upregulated by VN1R5 promotes cisplatin resistance in head and neck squamous cell carcinoma through interaction with MCM5. *Mol Ther* 2022;30:448-67. DOI PubMed PMC
118. Steuer CE, El-Deiry M, Parks JR, Higgins KA, Saba NF. An update on larynx cancer. *CA Cancer J Clin* 2017;67:31-50. DOI PubMed
119. Li R, Chen S, Zhan J, et al. Long noncoding RNA FOXD2-AS1 enhances chemotherapeutic resistance of laryngeal squamous cell carcinoma via STAT3 activation. *Cell Death Dis* 2020;11:41. DOI PubMed PMC
120. Pillai J, Chincholkar T, Dixit R, Pandey M. A systematic review of proteomic biomarkers in oral squamous cell cancer. *World J Surg Oncol* 2021;19:315. DOI PubMed PMC
121. Tian T, Lv X, Pan G, et al. Long noncoding RNA MPRL promotes mitochondrial fission and cisplatin chemosensitivity via disruption of pre-miRNA processing. *Clin Cancer Res* 2019;25:3673-88. DOI PubMed PMC
122. Filetti S, Durante C, Hartl D, et al; the ESMO Guidelines Committee. Thyroid cancer: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol* 2019;30:1856-83. DOI PubMed
123. Shi L, Duan R, Sun Z et al. LncRNA GLTC targets LDHA for succinylation and enzymatic activity to promote progression and radioiodine resistance in papillary thyroid cancer. *Cell Death Differ* 2023;30:1517-32. DOI PubMed PMC
124. Wang Y, Zhang X, Wang Z, et al. LncRNA-p23154 promotes the invasion-metastasis potential of oral squamous cell carcinoma by regulating Glut1-mediated glycolysis. *Cancer Lett* 2018;434:172-83. DOI PubMed
125. Sang L, Ju HQ, Yang Z, et al. Mitochondrial long non-coding RNA GAS5 tunes TCA metabolism in response to nutrient stress. *Nat Metab* 2021;3:90-106. DOI PubMed
126. Tan DSW, Chong FT, Leong HS, et al. Long noncoding RNA EGFR-AS1 mediates epidermal growth factor receptor addiction and modulates treatment response in squamous cell carcinoma. *Nat Med* 2017;23:1167-75. DOI PubMed
127. Zhang L, Meng X, Zhu XW, et al. Long non-coding RNAs in Oral squamous cell carcinoma: biologic function, mechanisms and clinical implications. *Mol Cancer* 2019;18:102. DOI PubMed PMC
128. Nair L, Chung H, Basu U. Regulation of long non-coding RNAs and genome dynamics by the RNA surveillance machinery. *Nat Rev Mol Cell Biol* 2020;21:123-36. DOI PubMed PMC
129. Zhang L, Xu X, Su X. Noncoding RNAs in cancer immunity: functions, regulatory mechanisms, and clinical application. *Mol Cancer*

- 2020;19:48. [DOI PubMed PMC](#)
130. Jiang W, Pan S, Chen X, Wang ZW, Zhu X. The role of lncRNAs and circRNAs in the PD-1/PD-L1 pathway in cancer immunotherapy. *Mol Cancer* 2021;20:116. [DOI PubMed PMC](#)
131. Zhou WY, Cai ZR, Liu J, Wang DS, Ju HQ, Xu RH. Circular RNA: metabolism, functions and interactions with proteins. *Mol Cancer* 2020;19:172. [DOI PubMed PMC](#)
132. Cheng J, Li G, Wang W, Stovall DB, Sui G, Li D. Circular RNAs with protein-coding ability in oncogenesis. *Biochim Biophys Acta Rev Cancer* 2023;1878:188909. [DOI PubMed](#)
133. Zhang Q, Wang W, Zhou Q, et al. Roles of circRNAs in the tumour microenvironment. *Mol Cancer* 2020;19:14. [DOI PubMed PMC](#)
134. Pisignano G, Michael DC, Visal TH, Pirlog R, Ladomery M, Calin GA. Going circular: history, present, and future of circRNAs in cancer. *Oncogene* 2023;42:2783-800. [DOI PubMed PMC](#)
135. Niu D, Wu Y, Lian J. Circular RNA vaccine in disease prevention and treatment. *Signal Transduct Target Ther* 2023;8:341. [DOI PubMed PMC](#)
136. Wu S, Lu J, Zhu H, et al. A novel axis of circKIF4A-miR-637-STAT3 promotes brain metastasis in triple-negative breast cancer. *Cancer Lett* 2024;581:216508. [DOI PubMed](#)
137. Wang Z, Yang L, Wu P, et al. The circROBO1/KLF5/FUS feedback loop regulates the liver metastasis of breast cancer by inhibiting the selective autophagy of afadin. *Mol Cancer* 2022;21:29. [DOI PubMed PMC](#)
138. Hong X, Li Q, Li J, et al. CircIPO7 promotes nasopharyngeal carcinoma metastasis and cisplatin chemoresistance by facilitating YBX1 nuclear localization. *Clin Cancer Res* 2022;28:4521-35. [DOI PubMed](#)
139. Li Q, Zhao YH, Xu C, et al. Chemotherapy-induced senescence reprogramming promotes nasopharyngeal carcinoma metastasis by circRNA-mediated PKR activation. *Adv Sci* 2023;10:e2205668. [DOI PubMed PMC](#)
140. Debnath J, Gammoh N, Ryan KM. Autophagy and autophagy-related pathways in cancer. *Nat Rev Mol Cell Biol* 2023;24:560-75. [DOI PubMed PMC](#)
141. Allen EA, Baehrecke EH. Autophagy in animal development. *Cell Death Differ* 2020;27:903-18. [DOI PubMed PMC](#)
142. Gao W, Guo H, Niu M, et al. circPARD3 drives malignant progression and chemoresistance of laryngeal squamous cell carcinoma by inhibiting autophagy through the PRKCI-Akt-mTOR pathway. *Mol Cancer* 2020;19:166. [DOI PubMed PMC](#)
143. Liu F, Zhang J, Qin L, et al. Circular RNA EIF6 (Hsa_circ_0060060) sponges miR-144-3p to promote the cisplatin-resistance of human thyroid carcinoma cells by autophagy regulation. *Aging* 2018;10:3806-20. [DOI PubMed PMC](#)
144. Li X, He S, Ma B. Autophagy and autophagy-related proteins in cancer. *Mol Cancer* 2020;19:12. [DOI PubMed PMC](#)
145. Ji X, Lv C, Huang J, Dong W, Sun W, Zhang H. ALKBH5-induced circular RNA NRIP1 promotes glycolysis in thyroid cancer cells by targeting PKM2. *Cancer Sci* 2023;114:2318-34. [DOI PubMed PMC](#)
146. Cui Y, Liu J, Liu L, et al. m⁶A-modified circFOXK2 targets GLUT1 to accelerate oral squamous cell carcinoma aerobic glycolysis. *Cancer Gene Ther* 2023;30:163-71. [DOI PubMed](#)
147. Zhao W, Liu J, Wu J, et al. High-throughput microarray reveals the epitranscriptome-wide landscape of m⁶A-modified circRNA in oral squamous cell carcinoma. *BMC Genomics* 2022;23:611. [DOI PubMed PMC](#)
148. Li J, Cao H, Yang J, Wang B. IGF2BP2-m6A-circMMP9 axis recruits ETS1 to promote TRIM59 transcription in laryngeal squamous cell carcinoma. *Sci Rep* 2024;14:3014. [DOI PubMed PMC](#)
149. Wu P, Fang X, Liu Y, et al. N6-methyladenosine modification of circCUX1 confers radioresistance of hypopharyngeal squamous cell carcinoma through caspase1 pathway. *Cell Death Dis* 2021;12:298. [DOI PubMed PMC](#)
150. Jiang X, Stockwell BR, Conrad M. Ferroptosis: mechanisms, biology and role in disease. *Nat Rev Mol Cell Biol* 2021;22:266-82. [DOI PubMed PMC](#)
151. Sun S, Shen J, Jiang J, Wang F, Min J. Targeting ferroptosis opens new avenues for the development of novel therapeutics. *Signal Transduct Target Ther* 2023;8:372. [DOI PubMed PMC](#)
152. Zhu J, Wang X, Su Y, et al. Multifunctional nanolocks with GSH as the key for synergistic ferroptosis and anti-chemotherapeutic resistance. *Biomaterials* 2022;288:121704. [DOI PubMed](#)
153. Zhang J, Ye ZW, Chakraborty P, et al. Microsomal glutathione transferase 1 controls metastasis and therapeutic response in melanoma. *Pharmacol Res* 2023;196:106899. [DOI PubMed PMC](#)
154. Yuan L, Li S, Chen Q, et al. EBV infection-induced GPX4 promotes chemoresistance and tumor progression in nasopharyngeal carcinoma. *Cell Death Differ* 2022;29:1513-27. [DOI PubMed PMC](#)
155. Roh JL, Kim EH, Jang H, Shin D. Nrf2 inhibition reverses the resistance of cisplatin-resistant head and neck cancer cells to artesunate-induced ferroptosis. *Redox Biol* 2017;11:254-62. [DOI PubMed PMC](#)
156. Shin D, Kim EH, Lee J, Roh JL. Nrf2 inhibition reverses resistance to GPX4 inhibitor-induced ferroptosis in head and neck cancer. *Free Radic Biol Med* 2018;129:454-62. [DOI PubMed](#)
157. Lee J, You JH, Shin D, Roh JL. Inhibition of glutaredoxin 5 predisposes cisplatin-resistant head and neck cancer cells to ferroptosis. *Theranostics* 2020;10:7775-86. [DOI PubMed PMC](#)
158. Yang JY, Lei XY, He KY, et al. HMGA1 drives chemoresistance in esophageal squamous cell carcinoma by suppressing ferroptosis. *Cell Death Dis* 2024;15:158. [DOI PubMed PMC](#)
159. Angeli JP, Krysko DV, Conrad M. Ferroptosis at the crossroads of cancer-acquired drug resistance and immune evasion. *Nat Rev Cancer* 2019;19:405-14. [DOI PubMed](#)

160. Li B, Yang L, Peng X, et al. Emerging mechanisms and applications of ferroptosis in the treatment of resistant cancers. *Biomed Pharmacother* 2020;130:110710. DOI PubMed
161. Sun X, Niu X, Chen R, et al. Metallothionein-1G facilitates sorafenib resistance through inhibition of ferroptosis. *Hepatology* 2016;64:488-500. DOI PubMed PMC
162. Jiang Z, Lim SO, Yan M, et al. TYRO3 induces anti-PD-1/PD-L1 therapy resistance by limiting innate immunity and tumoral ferroptosis. *J Clin Invest* 2021;131:139434. DOI PubMed PMC
163. Qi R, Bai Y, Li K, et al. Cancer-associated fibroblasts suppress ferroptosis and induce gemcitabine resistance in pancreatic cancer cells by secreting exosome-derived ACSL4-targeting miRNAs. *Drug Resist Updat* 2023;68:100960. DOI PubMed
164. Liang J, Bi G, Huang Y, et al. MAFF confers vulnerability to cisplatin-based and ionizing radiation treatments by modulating ferroptosis and cell cycle progression in lung adenocarcinoma. *Drug Resist Updat* 2024;73:101057. DOI PubMed
165. Zhang X, Xu Y, Ma L, et al. Essential roles of exosome and circRNA_101093 on ferroptosis desensitization in lung adenocarcinoma. *Cancer Commun* 2022;42:287-313. DOI PubMed PMC
166. Jiang M, Jike Y, Liu K, et al. Exosome-mediated miR-144-3p promotes ferroptosis to inhibit osteosarcoma proliferation, migration, and invasion through regulating ZEB1. *Mol Cancer* 2023;22:113. DOI PubMed PMC
167. Xi Y, Shen Y, Wu D, et al. CircBCAR3 accelerates esophageal cancer tumorigenesis and metastasis via sponging miR-27a-3p. *Mol Cancer* 2022;21:145. DOI PubMed PMC
168. Zhang C, Liu X, Jin S, Chen Y, Guo R. Ferroptosis in cancer therapy: a novel approach to reversing drug resistance. *Mol Cancer* 2022;21:47. DOI PubMed PMC
169. Zhang C, Xu C, Gao X, Yao Q. Platinum-based drugs for cancer therapy and anti-tumor strategies. *Theranostics* 2022;12:2115-32. DOI PubMed PMC
170. Huang Y, Kou Q, Su Y, et al. Combination therapy based on dual-target biomimetic nano-delivery system for overcoming cisplatin resistance in hepatocellular carcinoma. *J Nanobiotechnology* 2023;21:89. DOI PubMed PMC
171. Siemer S, Bauer TA, Scholz P, et al. Targeting cancer chemotherapy resistance by precision medicine-driven nanoparticle-formulated cisplatin. *ACS Nano* 2021;15:18541-56. DOI PubMed
172. Zhang R, You X, Luo M, et al. Poly(β -cyclodextrin)/platinum prodrug supramolecular nano system for enhanced cancer therapy: synthesis and in vivo study. *Carbohydr Polym* 2022;292:119695. DOI PubMed
173. Chen J, Wang X, Yuan Y, et al. Exploiting the acquired vulnerability of cisplatin-resistant tumors with a hypoxia-amplifying DNA repair-inhibiting (HYDRI) nanomedicine. *Sci Adv* 2021;7:eabc5267. DOI PubMed PMC
174. Chen Y, Fang L, Zhou W, et al. Nitric oxide-releasing micelles with intelligent targeting for enhanced anti-tumor effect of cisplatin in hypoxia. *J Nanobiotechnology* 2021;19:246. DOI PubMed PMC
175. Pan WL, Tan Y, Meng W, et al. Microenvironment-driven sequential ferroptosis, photodynamic therapy, and chemotherapy for targeted breast cancer therapy by a cancer-cell-membrane-coated nanoscale metal-organic framework. *Biomaterials* 2022;283:121449. DOI PubMed
176. Zhang Z, Ji Y, Hu N, et al. Ferroptosis-induced anticancer effect of resveratrol with a biomimetic nano-delivery system in colorectal cancer treatment. *Asian J Pharm Sci* 2022;17:751-66. DOI PubMed PMC
177. Li K, Lin C, Li M, et al. Multienzyme-like reactivity cooperatively impairs glutathione peroxidase 4 and ferroptosis suppressor protein 1 pathways in triple-negative breast cancer for sensitized ferroptosis therapy. *ACS Nano* 2022;16:2381-98. DOI PubMed
178. Li K, Xu K, He Y, et al. Oxygen self-generating nanoreactor mediated ferroptosis activation and immunotherapy in triple-negative breast cancer. *ACS Nano* 2023;17:4667-87. DOI PubMed
179. Chen D, Liang C, Qu X, et al. Metal-free polymer nano-photosensitizer actuates ferroptosis in starved cancer. *Biomaterials* 2023;292:121944. DOI PubMed
180. Liu J, Ren L, Li S, et al. The biology, function, and applications of exosomes in cancer. *Acta Pharm Sin B* 2021;11:2783-97. DOI PubMed PMC
181. Wu J, Li S, Zhang P. Tumor-derived exosomes: immune properties and clinical application in lung cancer. *Cancer Drug Resist* 2022;5:102-13. DOI PubMed PMC
182. Wang Z, He J, Bach DH, et al. Induction of m⁶A methylation in adipocyte exosomal lncRNAs mediates myeloma drug resistance. *J Exp Clin Cancer Res* 2022;41:4. DOI PubMed PMC
183. Shi L, Zhu W, Huang Y, et al. Cancer-associated fibroblast-derived exosomal microRNA-20a suppresses the PTEN/PI3K-AKT pathway to promote the progression and chemoresistance of non-small cell lung cancer. *Clin Transl Med* 2022;12:e989. DOI PubMed PMC
184. Pan Z, Zheng J, Zhang J, et al. A novel protein encoded by exosomal circATG4B induces oxaliplatin resistance in colorectal cancer by promoting autophagy. *Adv Sci* 2022;9:e2204513. DOI PubMed PMC
185. Jing X, Xie M, Ding K, et al. Exosome-transmitted miR-769-5p confers cisplatin resistance and progression in gastric cancer by targeting CASP9 and promoting the ubiquitination degradation of p53. *Clin Transl Med* 2022;12:e780. DOI PubMed PMC
186. Liu X, Guo Q, Gao G, et al. Exosome-transmitted circCABIN1 promotes temozolomide resistance in glioblastoma via sustaining ErbB downstream signaling. *J Nanobiotechnology* 2023;21:45. DOI PubMed PMC
187. Deng J, Pan T, Lv C, et al. Exosomal transfer leads to chemoresistance through oxidative phosphorylation-mediated stemness phenotype in colorectal cancer. *Theranostics* 2023;13:5057-74. DOI PubMed PMC
188. Zhang F, Jiang J, Qian H, Yan Y, Xu W. Exosomal circRNA: emerging insights into cancer progression and clinical application

- potential. *J Hematol Oncol* 2023;16:67. DOI PubMed PMC
189. Guo X, Gao C, Yang DH, Li S. Exosomal circular RNAs: a chief culprit in cancer chemotherapy resistance. *Drug Resist Updat* 2023;67:100937. DOI PubMed
190. Li K, Crews CM. PROTACs: past, present and future. *Chem Soc Rev* 2022;51:5214-36. DOI PubMed PMC
191. Marei H, Tsai WK, Kee YS, et al. Antibody targeting of E3 ubiquitin ligases for receptor degradation. *Nature* 2022;610:182-9. DOI PubMed PMC
192. Xiong Y, Zhong Y, Yim H, et al. Bridged proteolysis targeting chimera (PROTAC) enables degradation of undruggable targets. *J Am Chem Soc* 2022;144:22622-32. DOI PubMed PMC
193. Ghidini A, Cléry A, Halloy F, Allain FHT, Hall J. RNA-PROTACs: degraders of RNA-binding proteins. *Angew Chem Int Ed Engl* 2021;60:3163-9. DOI PubMed PMC
194. Li X, Pu W, Zheng Q, Ai M, Chen S, Peng Y. Proteolysis-targeting chimeras (PROTACs) in cancer therapy. *Mol Cancer* 2022;21:99. DOI PubMed PMC
195. Wu Y, Chang X, Yang G, et al. A physiologically responsive nanocomposite hydrogel for treatment of head and neck squamous cell carcinoma via proteolysis-targeting chimeras enhanced immunotherapy. *Adv Mater* 2023;35:e2210787. DOI PubMed
196. Jin J, Wu Y, Zhao Z, et al. Small-molecule PROTAC mediates targeted protein degradation to treat STAT3-dependent epithelial cancer. *JCI Insight* 2022;7:e160606. DOI PubMed PMC
197. Kuismi CC, Nabet B. Lighting the way to tumor destruction. *Trends Pharmacol Sci* 2023;44:750-2. DOI PubMed PMC
198. Yang F, Luo Q, Wang Y, et al. Targeted biomolecule regulation platform: a split-and-mix PROTAC approach. *J Am Chem Soc* 2023;145:7879-87. DOI PubMed
199. Collins R. What makes UK Biobank special? *Lancet* 2012;379:1173-4. DOI PubMed
200. Zhang C, Xu J, Tang R, et al. Novel research and future prospects of artificial intelligence in cancer diagnosis and treatment. *J Hematol Oncol* 2023;16:114. DOI PubMed PMC
201. Tayebi Arasteh S, Han T, Lotfinia M, et al. Large language models streamline automated machine learning for clinical studies. *Nat Commun* 2024;15:1603. DOI PubMed PMC